

Convex Aggregation Certificates for Quadratic Systems

1 Quadratic systems and the certificate question

Let $n, m \geq 1$, and let

$$f_i(x) = x^\top A_i x + 2b_i^\top x + c_i, \quad S = \{x \in \mathbb{R}^n : f_i(x) < 0 \text{ for } i = 1, \dots, m\}, \quad (1.1)$$

where $A_i \in \mathbb{S}^n$, $b_i \in \mathbb{R}^n$, and $c_i \in \mathbb{R}$. Here $\mathbb{S}^n$ denotes the real symmetric $n \times n$ matrices; $A \succeq 0$ and $A \succ 0$ mean positive semidefinite and positive definite, respectively. We use the Euclidean vector norm and the Frobenius matrix norm, and write $\langle A, B \rangle = \text{tr}(AB)$. The symbol $\text{conv } S$ denotes the ordinary convex hull, without taking its closure. A hull is *proper* if it differs from $\mathbb{R}^n$. Our question is whether a nonempty quadratic system has a proper hull, and how this can be certified by its defining inequalities.

For $\lambda \in \mathbb{R}^m$, write

$$(A_\lambda, b_\lambda, c_\lambda) = \sum_{i=1}^m \lambda_i (A_i, b_i, c_i), \quad f_\lambda = \sum_{i=1}^m \lambda_i f_i.$$

A nonzero $\lambda \geq 0$ defines an *aggregation*: $f_\lambda(x) < 0$ is valid for every $x \in S$. We call λ a *convex certificate* if $A_\lambda \succeq 0$, so that f_λ is convex on $\mathbb{R}^n$. It is *nontrivial* if $(A_\lambda, b_\lambda) \neq (0, 0)$. Otherwise the aggregated function is constant. If $S \neq \emptyset$, that constant is strictly negative, as evaluation at any point of S shows. We retain zero in the following cone for convenience:

$$\mathcal{K} = \{\lambda \in \mathbb{R}_+^m : A_\lambda \succeq 0\}, \quad \mathcal{T} = \{\lambda \in \mathbb{R}^m : A_\lambda = 0, b_\lambda = 0\}. \quad (1.2)$$

Thus a nontrivial convex certificate exists precisely when $\mathcal{K} \not\subseteq \mathcal{T}$.

The homogeneous matrices and map associated with (1.1) are

$$Q_i = \begin{pmatrix} A_i & b_i \\ b_i^\top & c_i \end{pmatrix}, \quad f_i^h(x, t) = x^\top A_i x + 2tb_i^\top x + c_i t^2, \quad f^h = (f_1^h, \dots, f_m^h). \quad (1.3)$$

We also write $Q_\lambda = \sum_i \lambda_i Q_i$ and $g_i(x) = x^\top A_i x$. Define the strict homogeneous feasible set $S^h = \{(x, t) : f_i^h(x, t) < 0 \text{ for all } i\}$. It is open and invariant under multiplication by every nonzero real scalar. For $t \neq 0$, $(x, t) \in S^h$ is equivalent to $x/t \in S$. For $\alpha \in \mathbb{R}^n \setminus \{0\}$ and $s \in \mathbb{R}$, put

$$H_{\alpha, s} = \{(x, t) \in \mathbb{R}^{n+1} : \alpha^\top x = st\}, \quad E = \{(x, t) : t = 0\}. \quad (1.4)$$

Definition 1.1 (Hidden hyperplane convexity). A homogeneous quadratic map $\phi : \mathbb{R}^d \rightarrow \mathbb{R}^m$ has *hidden hyperplane convexity* (HHC) if $\phi(H)$ is convex for every linear hyperplane $H \subset \mathbb{R}^d$.

This is the definition of Blekherman, Dey, and Sun [4, Definition 2.1]; their work appeared as [5]. Ordinary *hidden convexity* means only that $\phi(\mathbb{R}^d)$ is convex. The two properties differ. All homogeneous quadratic images contain zero and are invariant under nonnegative scaling, so a convex such image is a convex cone; it need not be closed. HHC implies ordinary hidden convexity when $d \geq 3$: any two points of $\mathbb{R}^d$ lie in some linear hyperplane, whose image contains the segment joining their images.

Question 1.2 (Blekherman–Dey–Sun, Conjecture 3.3). Suppose $S \neq \emptyset$ and f^h has HHC. Is

$$\text{conv } S \neq \mathbb{R}^n \iff \exists \lambda \in \mathbb{R}_+^m : A_\lambda \succeq 0, \quad (A_\lambda, b_\lambda) \neq (0, 0)? \quad (1.5)$$

The formulation in Blekherman et al. [4, Conjecture 3.3] says that the aggregated function is not a negative constant. Under nonemptiness this is equivalent to the displayed condition. Their standing setting has $n \geq 3$ and $m \geq 2$; the formulation here also includes the smaller positive dimensions and numbers of inequalities. Throughout, numbered result locators for that work refer to the explicitly cited May 29, 2023 arXiv version; the earlier author manuscript has different numbering for several background results.

The reverse implication in (1.5) is elementary: a nonconstant convex quadratic aggregation bounds S by a proper convex sublevel set. The unresolved case in the cited conjecture concerns the opposite implication when $\mathcal{K}$ contains nonzero vectors but all of them belong to $\mathcal{T}$. Under convexity of $(g_1, \dots, g_m)(\mathbb{R}^n)$, Blekherman, Dey, and Sun already prove that $\mathcal{K} = \{0\}$ implies $\text{conv } S = \mathbb{R}^n$ [4, Proposition 2.14]. For $n \geq 2$, their separable case, with all Q_i diagonal, also already gives the certificate characterization [4, Theorem 2.12].

It is essential to distinguish this certificate question from a complete description of the hull. In the aggregation literature, a *good aggregation* is a nonnegative nonzero multiplier λ such that $\text{conv } S \subseteq \{x : f_\lambda(x) < 0\}$ and Q_λ has at most one negative eigenvalue. The function f_λ need not be globally convex. Blekherman, Dey, and Sun prove that these good aggregations describe $\text{conv } S$ under HHC, $n \geq 3$, and $\emptyset \neq S$ with proper hull [4, Theorem 2.9]. Thus an existence certificate and a family of hull-defining aggregations address different questions.

The two-constraint case belongs to established theory. Dines’ theorem gives convexity of the joint image of two homogeneous quadratic forms [7], and therefore HHC for a pair of homogeneous forms and their restrictions. Yildiran characterizes the hull of a nonempty set defined by two strict quadratic inequalities using at most two aggregations [13]. For three quadratic inequalities in dimension $n \geq 3$ with a nonempty proper hull, Dey, Muñoz, and Serrano obtain an aggregation description when the homogenized matrices Q_1, Q_2, Q_3 admit a positive definite real linear combination [6]. Here “positive definite linear combination” allows real coefficients of either sign; it does not mean a nonnegative aggregation. More recently, Blekherman and Dunbar use the spectral curve and algebraic topology to obtain further three-quadratic hull descriptions [3]. Their available preprint treats finiteness and related spectral conditions [2]. These results set the context for the certificate question, while neither the classical two-form convexity theorem nor an elementary separation lemma is itself a new contribution of the present work.

2 The certificate theorem

Only a family of hyperplanes approaching E is needed to answer Question 1.2. We first make this weaker assumption precise.

Definition 2.1 (Asymptotic hyperplane convexity). The map f^h has *asymptotic hyperplane convexity* (AHC) if, for every $\alpha \neq 0$ and every $R \in \mathbb{R}$, there exists $s > R$ such that $f^h(H_{\alpha,s})$ is convex.

Equivalently, for each $\alpha \neq 0$ there is a strictly increasing sequence (s_k) tending to $+\infty$ with all these images convex. Indeed, the sequence condition implies the unbounded-level condition; conversely, choose a good $s_1 > 1$ and recursively a good $s_{k+1} > \max\{s_k, k+1\}$. HHC implies AHC because $H_{\alpha,s}$ is a linear hyperplane for every s . This assumption concerns the images themselves and requires no closedness.

Theorem 2.2 (Convex aggregation certificate). *Suppose $S \neq \emptyset$ and f^h has AHC. Then*

$$\text{conv } S \neq \mathbb{R}^n \iff \exists \lambda \geq 0 : \quad A_\lambda \succeq 0, \quad (A_\lambda, b_\lambda) \neq (0, 0). \quad (2.1)$$

The certificate-to-proper-hull implication holds without either assumption.

Corollary 2.3 (Conjecture 3.3 of Blekherman–Dey–Sun). *If $S \neq \emptyset$ and f^h has HHC, then (2.1) holds.*

The corollary follows from HHC implying AHC. It answers the cited conjecture for every finite number of inequalities, including the case where nonzero convex certificates exist but all their quadratic and linear coefficients might vanish. The proof below shows that this case forces a whole-space hull. The theorem also isolates a weaker sufficient hypothesis; we attach no novelty claim to the separation or compactness facts it uses.

2.1 Hyperplanes beyond a supporting halfspace

Lemma 2.4 (No common strict negative quadratic direction). *If $\text{conv } S \neq \mathbb{R}^n$, then there is no $v \in \mathbb{R}^n$ satisfying $v^\top A_i v < 0$ for every i . Equivalently, $E \cap S^h = \emptyset$.*

Proof. If such a v existed, then for any fixed $y \in \mathbb{R}^n$,

$$f_i(y + rv) = r^2 v^\top A_i v + 2r(v^\top A_i y + b_i^\top v) + f_i(y) \longrightarrow -\infty \quad (|r| \longrightarrow \infty).$$

There are finitely many constraints. Hence both $y + rv$ and $y - rv$ belong to S for all sufficiently large positive r , and their midpoint y belongs to $\text{conv } S$. This holds for every y , a contradiction. Finally, $f_i^h(v, 0) = v^\top A_i v$ proves the equivalence. $\square$

This lemma concerns a simultaneous strict sign condition on the leading forms. It is not a characterization of the recession cone of S .

Lemma 2.5 (Sweeping hyperplanes). *Suppose $S \neq \emptyset$ and $\text{conv } S \neq \mathbb{R}^n$. There are $\alpha \neq 0$ and $\beta \in \mathbb{R}$ such that*

$$\alpha^\top x < \beta \quad (x \in S). \quad (2.2)$$

For any such α, β , one has $H_{\alpha, s} \cap S^h = \emptyset$ for every $s \geq \beta$.

Proof. The set S is open, and its convex hull is open as well: in a finite convex combination, one point with positive weight can vary in an open ball, giving an open ball around the combination. Separate a point $y \notin \text{conv } S$ from this nonempty open convex set. Finite-dimensional separation supplies $\alpha \neq 0$ with $\alpha^\top x \leq \alpha^\top y$ throughout $\text{conv } S$. Equality at a point of that open set would be contradicted by a small move in direction α . Thus (2.2) holds with $\beta = \alpha^\top y$.

If $(x, t) \in H_{\alpha, s} \cap S^h$ and $t = 0$, then Lemma 2.4 gives a contradiction. If $t \neq 0$, the identity $f_i^h(x, t) = t^2 f_i(x/t)$ gives $x/t \in S$, whereas $\alpha^\top(x/t) = s \geq \beta$, again a contradiction. This reasoning applies to either sign of t . $\square$

Lemma 2.6 (Separation on a hyperplane). *Suppose $s \neq 0$, the set $f^h(H_{\alpha, s})$ is convex, and $H_{\alpha, s} \cap S^h = \emptyset$. There exists $\lambda \geq 0$ with $\sum_i \lambda_i = 1$ such that, for every $x \in \mathbb{R}^n$,*

$$0 \leq x^\top A_\lambda x + \frac{2}{s}(\alpha^\top x)(b_\lambda^\top x) + \frac{c_\lambda}{s^2}(\alpha^\top x)^2. \quad (2.3)$$

Proof. The convex image $C = f^h(H_{\alpha,s})$ contains zero and is disjoint from the open negative orthant $N = -\mathbb{R}_{++}^m$. Separation of these two convex sets gives a nonzero λ and a real a with $\lambda^\top z \leq a \leq \lambda^\top y$ for $z \in N$, $y \in C$. If some $\lambda_i < 0$, sending z_i to $-\infty$ with the other coordinates fixed contradicts the upper bound. Thus $\lambda \geq 0$, and $\sup_{z \in N} \lambda^\top z = 0$. Since $0 \in C$, necessarily $a = 0$, and $\lambda^\top f^h \geq 0$ on $H_{\alpha,s}$. Divide λ by its positive coordinate sum. Substituting $(x, t) = (x, \alpha^\top x/s)$ gives (2.3). Closedness of C was not used. $\square$

2.2 A nonzero limit in coefficient space

The relevant cone retains the quadratic and linear coefficients together:

$$P = \text{cone}\{(A_i, b_i) : 1 \leq i \leq m\} = \{(A_\lambda, b_\lambda) : \lambda \geq 0\} \subseteq \mathbb{S}^n \times \mathbb{R}^n. \quad (2.4)$$

We equip this space with $\|(A, b)\| = (\|A\|_F^2 + \|b\|^2)^{1/2}$. Closedness of this particular cone is essential: a limit in its closure must correspond to actual aggregation weights. The following elementary fact supplies that property.

Lemma 2.7 (Finite coefficient cones are closed). *The cone generated by a finite family in a finite-dimensional real vector space is closed.*

Proof. Every conic combination can be represented using a linearly independent subfamily. To see this, choose a representation with the fewest positive coefficients. If its active generators are dependent, take a nonzero relation $\sum_i d_i v_i = 0$, reversing its sign so some $d_i > 0$. For the active coefficients $a_i > 0$, set $\tau = \min_{d_i > 0} a_i/d_i$. Then all $a_i - \tau d_i$ are nonnegative, at least one vanishes, and the represented point is unchanged, a contradiction. For a linearly independent subfamily, the coefficient map is a linear homeomorphism onto its span; that span is closed, so the image of the nonnegative orthant is closed. There are only finitely many subfamilies. The original cone is their finite union and is closed. The empty subfamily accounts for the zero vector. $\square$

Proof of Theorem 2.2. First suppose a nontrivial convex certificate λ exists. Its nonconstant coefficient pair implies $\lambda \neq 0$, so $f_\lambda < 0$ on S . Positive semidefiniteness makes f_λ convex, whence $\text{conv } S \subseteq \{f_\lambda < 0\}$. This sublevel set is proper. Indeed, if $A_\lambda \neq 0$, it has a unit eigenvector v with eigenvalue $\mu > 0$, and $f_\lambda(rv) = \mu r^2 + 2rb_\lambda^\top v + c_\lambda \rightarrow +\infty$ as $r \rightarrow +\infty$. If $A_\lambda = 0$, then $b_\lambda \neq 0$ and $f_\lambda(rb_\lambda) = 2r\|b_\lambda\|^2 + c_\lambda \rightarrow +\infty$. This proves the easy implication without using nonemptiness or AHC.

For the converse, suppose $S \neq \emptyset$, f^h has AHC, and $\text{conv } S \neq \mathbb{R}^n$. Fix $x_0 \in S$, and choose α, β from Lemma 2.5. AHC gives positive levels $s_k \rightarrow +\infty$ with $s_k \geq \beta$ and convex images $f^h(H_{\alpha,s_k})$. Lemmas 2.5 and 2.6 give nonzero normalized weights $\lambda^k \geq 0$ satisfying (2.3). Write $(A_k, b_k, c_k) = (A_{\lambda^k}, b_{\lambda^k}, c_{\lambda^k})$. Strict feasibility gives the crucial upper bound

$$c_k < -x_0^\top A_k x_0 - 2b_k^\top x_0 =: L(A_k, b_k), \quad (2.5)$$

where L is a fixed continuous linear functional on coefficient space. The multiplier of c_k in (2.3) is a square. Replacing c_k by its upper bound therefore yields

$$0 \leq x^\top A_k x + \frac{2}{s_k}(\alpha^\top x)(b_k^\top x) + \frac{L(A_k, b_k)}{s_k^2}(\alpha^\top x)^2 \quad (x \in \mathbb{R}^n). \quad (2.6)$$

Moreover, $(A_k, b_k) \neq (0, 0)$: otherwise (2.5) gives $c_k < 0$, and the original inequality (2.3) at $x = \alpha$ reads $0 \leq c_k \|\alpha\|^4 / s_k^2 < 0$.

Set $\rho_k = \|(A_k, b_k)\| > 0$ and $(\hat{A}_k, \hat{b}_k) = (A_k, b_k)/\rho_k$. These pairs lie in the unit section of P , which is compact by Lemma 2.7. Passing to a subsequence, let

$$(\hat{A}_k, \hat{b}_k) \longrightarrow (A_*, b_*) \in P, \quad \|(A_*, b_*)\| = 1.$$

Divide (2.6) by ρ_k . By linearity of L , its right-hand side becomes

$$x^\top \hat{A}_k x + \frac{2}{s_k} (\alpha^\top x) (\hat{b}_k^\top x) + \frac{L(\hat{A}_k, \hat{b}_k)}{s_k^2} (\alpha^\top x)^2.$$

For every fixed x the last two terms tend to zero. Consequently $x^\top A_* x \geq 0$ for every x . This conclusion uses one coefficient subsequence for all x ; no point-dependent subsequence is chosen. The symmetric matrix A_* is positive semidefinite. Since $(A_*, b_*) \in P$, there are actual weights $\lambda^* \geq 0$ with $(A_{\lambda^*}, b_{\lambda^*}) = (A_*, b_*)$. Their coefficient pair has norm one, so they give the required nontrivial convex certificate. $\square$

Remark 2.8 (What the limit proves). Normalizing the weights in the simplex alone can give a negative constant aggregation in the limit. Normalizing the nonconstant coefficient pair prevents this loss of information. Bound (2.5) removes the constant before normalization; no bound on c_k/ρ_k is assumed. The final weights need not be a limit of the original weights, or of their rescalings. Their existence follows from closedness of the finitely generated cone P .

Remark 2.9 (Convexity along one normal). The hard implication needs convexity of $f^h(H_{\alpha,s})$ at arbitrarily large levels for only one α satisfying (2.2). AHC is a convenient condition that can be stated without knowing a supporting normal. For fixed α , the hyperplanes $H_{\alpha,s}$ approach E as $s \rightarrow +\infty$; convexity of $f^h(E)$ alone is not the hypothesis used in the proof. There is no restriction beyond $n, m \geq 1$. For one constraint, a negative quadratic direction gives a whole-space hull by Lemma 2.4; the remaining cases follow from the easy implication or a negative constant. For two constraints, Dines' theorem makes HHC automatic, recovering the known certificate case discussed in Section 1.

2.3 A quantitative form of the obstruction

For completeness, we give a second way to exclude degeneration to constant certificates. It supplies a uniform bound on the sweep level under the hypothesis that all convex certificates are trivial. The cone estimate itself is a standard compactness argument.

Lemma 2.10 (Uniform cone separation). *Let P, D be closed cones containing zero in a finite-dimensional normed space, with $P \cap D = \{0\}$. Then some $\kappa > 0$ satisfies*

$$\text{dist}(p, D) \geq \kappa \|p\| \quad (p \in P).$$

Neither cone is required to be convex.

Proof. If $P = \{0\}$, any positive κ works. Otherwise the continuous function $p \mapsto \text{dist}(p, D)$ has a positive minimum κ on the compact set $P \cap \{\|p\| = 1\}$: a zero distance would put a unit point of P in the closed set D . Scaling both cones proves the bound for $p \neq 0$, and the assertion at zero follows from $0 \in D$. $\square$

Apply the lemma to P in (2.4) and $D = \mathbb{S}_+^n \times \mathbb{R}^n$. The condition $P \cap D = \{0\}$ is exactly the assertion that every convex certificate is trivial. If A_- is the positive semidefinite negative part of a symmetric matrix A , then

$$\text{dist}((A, b), D) = \|A_-\|_F \leq \sqrt{n} \max\{0, -\lambda_{\min}(A)\}. \quad (2.7)$$

For the equality, diagonalize A orthogonally. In that basis every positive semidefinite approximant has nonnegative diagonal entries; the sum of squared errors is at least the sum of squares of the negative eigenvalues of A . Replacing those eigenvalues by zero achieves this bound, and the b component can be left unchanged. The inequality follows by bounding each negative eigenvalue in absolute value by $\max\{0, -\lambda_{\min}(A)\}$.

Proposition 2.11 (Bound on a sweep certificate). *Suppose every convex certificate is trivial, and let κ be as in Lemma 2.10 for the preceding cones. If $\lambda \geq 0$, $c_\lambda < 0$, $s > 0$, $\alpha \neq 0$, and (2.3) holds, then*

$$s \leq \frac{2\sqrt{n}\|\alpha\|}{\kappa}. \quad (2.8)$$

Proof. The inequality at $x = \alpha$ shows that $\rho = \|(A_\lambda, b_\lambda)\|$ cannot vanish. The cone estimate and (2.7) give $\lambda_{\min}(A_\lambda) \leq -\kappa\rho/\sqrt{n}$. Choose a unit eigenvector u for this eigenvalue. Evaluating (2.3) at u , discarding the nonpositive constant term, and using $\|b_\lambda\| \leq \rho$ gives

$$0 \leq -\frac{\kappa}{\sqrt{n}}\rho + \frac{2\|\alpha\|}{s}\rho.$$

Division by $\rho > 0$ proves (2.8). $\square$

This gives an alternative completion of the hard implication. Under its hypotheses, suppose every convex certificate is trivial and choose the normalized sweep weights λ^k as before. Compactness of the simplex gives a subsequence converging to $\bar{\lambda}$, with $\bar{\lambda} \geq 0$ and $\sum_i \bar{\lambda}_i = 1$. The coefficients b_k, c_k are bounded, so letting $k \rightarrow \infty$ in (2.3) shows $A_{\bar{\lambda}} \succeq 0$. Triviality then gives $A_{\bar{\lambda}} = 0$, $b_{\bar{\lambda}} = 0$, and $c_{\bar{\lambda}} < 0$ by strict feasibility. Thus $c_k < 0$ eventually along this subsequence. Proposition 2.11 contradicts $s_k \rightarrow +\infty$. The constant κ depends only on the original quadratic and linear coefficients; this argument neither bounds aggregation weights nor provides a procedure for verifying AHC.

3 Consequences for closed systems and semidefinite relaxations

The certificate theorem determines whether a hull is the whole space. Its consequences below concern that dichotomy; they do not identify every supporting inequality of a proper hull.

3.1 Closed inequalities and the three feasibility regimes

Write $T = \{x : f_i(x) \leq 0 \text{ for all } i\}$.

Corollary 3.1. *Suppose $S \neq \emptyset$ and f^h has asymptotic hyperplane convexity. Then*

$$\text{conv } T \neq \mathbb{R}^n \iff \text{conv } S \neq \mathbb{R}^n \iff \mathcal{K} \not\subseteq \mathcal{T}.$$

Proof. The inclusion $S \subseteq T$ gives one implication. A nontrivial convex certificate bounds T by the closed convex set $\{f_\lambda \leq 0\}$. That set is proper because the nonconstant convex quadratic f_λ is unbounded above. The remaining implication is Theorem 2.2. $\square$

Strict feasibility is essential here; Example 4.5 gives a counterexample even with HHC and a closed-system hull with interior. No equality between $\text{conv } T$ and $\text{cl conv } S$ is asserted.

Corollary 3.2 (Three regimes under HHC). *Suppose f^h has HHC and $n \geq 2$. Exactly one of the following holds:*

1. $S = \emptyset$. This is equivalent to the existence of $\lambda \geq 0$, $\lambda \neq 0$, with $Q_\lambda \succeq 0$.
2. $S \neq \emptyset$ and $\text{conv } S = \mathbb{R}^n$. This is equivalent to nonemptiness together with $\mathcal{K} \subseteq \mathcal{T}$.
3. $S \neq \emptyset$ and $\text{conv } S \neq \mathbb{R}^n$. This is equivalent to nonemptiness together with $\mathcal{K} \not\subseteq \mathcal{T}$. If also $n \geq 3$, the good aggregations defined in Section 1 describe $\text{conv } S$ exactly.

Proof. For the first statement, HHC implies convexity of $f^h(\mathbb{R}^{n+1})$ because $n+1 \geq 3$. Moreover, $S = \emptyset$ if and only if $S^h = \emptyset$: a strict homogeneous point with $t = 0$ can be perturbed to one with $t \neq 0$, and then dehomogenized. Separation of the convex homogeneous image from the open negative orthant gives a nonzero $\lambda \geq 0$ for which $\lambda^\top f^h \geq 0$, equivalently $Q_\lambda \succeq 0$. The converse follows by evaluation at a hypothetical strict point. This is the classical separation argument used in Blekherman et al. [4, Proposition 2.13]. Theorem 2.2 supplies the second and third statements. The final hull description is Blekherman et al. [4, Theorem 2.9]. $\square$

The proper-hull certificate itself remains valid for $n = 1$ by Theorem 2.2; the dimension restrictions above belong to the additional emptiness and full-description statements.

3.2 A finite semidefinite characterization

The certificate cone can be examined on the compact slice

$$\mathcal{D} = \{\lambda \geq 0 : \sum_i \lambda_i = 1, A_\lambda \succeq 0\}.$$

If $\mathcal{D}$ is nonempty, let

$$\tau = \max_{\lambda \in \mathcal{D}} \text{tr } A_\lambda, \quad \beta_j^\pm = \max_{\lambda \in \mathcal{D}} (\pm (b_\lambda)_j) \quad (j = 1, \dots, n). \quad (3.1)$$

All maxima are attained. These are $2n+1$ semidefinite programs, with one $n \times n$ positive semidefinite block and m nonnegative weights.

Corollary 3.3. *The inclusion $\mathcal{K} \subseteq \mathcal{T}$ holds exactly when $\mathcal{D} = \emptyset$, or $\mathcal{D} \neq \emptyset$ and all the values in (3.1) are zero. Under the hypotheses of Theorem 2.2, these conditions characterize $\text{conv } S = \mathbb{R}^n$.*

Proof. Every nonzero element of $\mathcal{K}$ has a positive multiple in $\mathcal{D}$. For a positive semidefinite matrix, zero trace is equivalent to the matrix being zero. Thus $\tau = 0$ forces $A_\lambda = 0$ throughout $\mathcal{D}$. The two coordinate objectives force $(b_\lambda)_j \leq 0$ and $(b_\lambda)_j \geq 0$ there. The converse is immediate. If $\mathcal{D}$ is empty, $\mathcal{K} = \{0\}$. $\square$

The reduction uses trace in place of all matrix coordinates. It is an exact characterization by feasibility and optimal values, not a bit-complexity result or a floating-point decision procedure. In particular, small computed objective values do not establish the exact zeros needed for a whole-space conclusion. Feasibility of $\mathcal{D}$ can be checked as the common feasibility part of these programs.

3.3 What the Shor projection preserves

Define the projection of the standard semidefinite relaxation by

$$P_{\text{SDP}} = \left\{ x \in \mathbb{R}^n : \begin{array}{l} \exists X \in \mathbb{S}^n, \quad X - xx^\top \succeq 0, \\ \langle A_i, X \rangle + 2b_i^\top x + c_i \leq 0 \quad (i = 1, \dots, m) \end{array} \right\}. \quad (3.2)$$

The matrix condition is equivalent to $\begin{pmatrix} 1 & x^\top \\ x & X \end{pmatrix} \succeq 0$, so P_{SDP} is convex. It contains T , by taking $X = xx^\top$. Let

$$C_{\text{agg}} = \bigcap_{\lambda \in \mathcal{K}} \{x : f_\lambda(x) \leq 0\}. \quad (3.3)$$

This is a closed convex set. For any Shor witness X and $\lambda \in \mathcal{K}$,

$$f_\lambda(x) = \sum_i \lambda_i (\langle A_i, X \rangle + 2b_i^\top x + c_i) - \langle A_\lambda, X - xx^\top \rangle \leq 0. \quad (3.4)$$

Consequently $P_{\text{SDP}} \subseteq C_{\text{agg}}$. This is the classical semidefinite/convex-aggregation relationship studied by Fujie and Kojima [9], Kojima and Tunçel [10]. We prove the precise closure statement needed here; projection closedness must not be assumed.

Proposition 3.4. *If $S \neq \emptyset$, then $\text{cl } P_{\text{SDP}} = C_{\text{agg}}$. Under the same assumption, without any hidden-convexity hypothesis,*

$$P_{\text{SDP}} = \mathbb{R}^n \iff \mathcal{K} \subseteq \mathcal{T}. \quad (3.5)$$

Proof. Put $g(v) = (v^\top A_i v)_i$ and introduce the convex cone

$$\mathcal{C} = \{(\langle A_i, Y \rangle)_i : Y \succeq 0\} + \mathbb{R}_+^m.$$

It contains $\mathbb{R}_+^m$ and is therefore full dimensional. It need not be closed. Directly from (3.2),

$$x \in P_{\text{SDP}} \iff -f(x) \in \mathcal{C}, \quad \mathcal{C}^* = \mathcal{K}, \quad x \in C_{\text{agg}} \iff -f(x) \in \text{cl } \mathcal{C}.$$

The dual identity follows from the self-duality of the positive semidefinite cone; the last equivalence is the finite-dimensional bipolar theorem.

We will use $\text{cl } \mathcal{C} + \text{int } \mathcal{C} \subseteq \text{int } \mathcal{C}$. For completeness, if $B(c, r) \subseteq \mathcal{C}$ and $d_k \in \mathcal{C}$ tends to $d \in \text{cl } \mathcal{C}$, then for large k the ball $B(c + d, r/2)$ lies in $B(c + d_k, r)$, which lies in $\mathcal{C} + \mathcal{C} = \mathcal{C}$. This also proves that adding another element of $\mathcal{C}$ preserves the interior.

Choose $x_0 \in S$, so $-f(x_0) \in \mathbb{R}_{++}^m \subseteq \text{int } \mathcal{C}$. For $x \in C_{\text{agg}}$ and $0 < t < 1$, put $x_t = (1 - t)x + tx_0$. The quadratic identity

$$-f(x_t) = (1 - t)(-f(x)) + t(-f(x_0)) + t(1 - t)g(x - x_0) \quad (3.6)$$

puts $-f(x_t)$ in $\text{int } \mathcal{C}$, since $g(v) = (\langle A_i, vv^\top \rangle)_i \in \mathcal{C}$. Thus $x_t \in P_{\text{SDP}}$, and $x_t \rightarrow x$ proves $C_{\text{agg}} \subseteq \text{cl } P_{\text{SDP}}$. The opposite inclusion follows from (3.4) and closedness of C_{agg} .

A nontrivial convex certificate makes C_{agg} proper, so $P_{\text{SDP}} = \mathbb{R}^n$ implies $\mathcal{K} \subseteq \mathcal{T}$. Conversely, suppose every certificate is trivial. For any $\lambda \in \mathcal{K} \setminus \{0\}$ strict feasibility gives $f_\lambda \equiv c_\lambda < 0$. Hence $C_{\text{agg}} = \mathbb{R}^n$ and $-f(z) \in \text{cl } \mathcal{C}$ for every z . For an arbitrary target x , apply (3.6) with $z = 2x - x_0$ in place of x and $t = 1/2$. Its left side is $-f(x)$; its right side belongs to $\text{int } \mathcal{C}$. Thus $x \in P_{\text{SDP}}$. $\square$

The closure conclusion also follows from Fujie and Kojima [9, Theorem 2.1]: strict feasibility provides their semidefinite Slater condition by choosing $X = x_0 x_0^\top + \varepsilon I$ for sufficiently small $\varepsilon > 0$. The elementary cone proof above records explicitly where strict feasibility enters and why no closedness of a cone image is required.

Corollary 3.5. *If $S \neq \emptyset$ and f^h has asymptotic hyperplane convexity, then*

$$P_{\text{SDP}} = \mathbb{R}^n \iff \text{conv } S = \mathbb{R}^n \iff \text{conv } T = \mathbb{R}^n.$$

Proof. Combine Proposition 3.4, Theorem 2.2, and Corollary 3.1. $\square$

Thus the relaxation preserves the existence of at least one nonconstant valid affine bound under these hypotheses. It need not preserve each particular bound; Example 4.6 makes the distinction explicit.

Example 3.6 (A nonclosed Shor projection with strict feasibility). In $\mathbb{R}^2$, let

$$f_1(x) = 2x_1x_2 + 1, \quad f_2(x) = x_1^2 - x_1.$$

The point $(1/2, -2)$ is strictly feasible. A nonnegative aggregate has quadratic matrix $\begin{pmatrix} \lambda_2 & \lambda_1 \\ \lambda_1 & 0 \end{pmatrix}$, which is positive semidefinite exactly when $\lambda_1 = 0$. Therefore

$$C_{\text{agg}} = [0, 1] \times \mathbb{R}, \quad P_{\text{SDP}} = ((0, 1) \times \mathbb{R}) \cup (\{1\} \times (-\infty, -1/2]).$$

To verify the latter equality, write $Y = X - xx^\top \succeq 0$. The second lifted row gives $0 \leq Y_{11} \leq x_1 - x_1^2$, so $0 \leq x_1 \leq 1$. At either endpoint $Y_{11} = 0$, and positive semidefiniteness forces $Y_{12} = 0$. At $x_1 = 0$ the first row then fails; at $x_1 = 1$ it requires $x_2 \leq -1/2$, and $Y = 0$ suffices. For $0 < x_1 < 1$, take

$$X_{11} = \frac{x_1^2 + x_1}{2}, \quad X_{12} = -\frac{1}{2}, \quad Y_{22} = \frac{(-1/2 - x_1x_2)^2}{(x_1 - x_1^2)/2}, \quad X_{22} = x_2^2 + Y_{22}.$$

Then $Y \succeq 0$ has zero determinant and the lifted rows hold. The displayed description proves nonclosedness even with strict feasibility and, by Dines' theorem, HHC.

Compactness of the original closed feasible set does not in general restore projection closedness. Add the two rows $f_3 = -2x_1x_2 - 2$ and $f_4 = 1/4 - x_1$ to this example. The point $(1/2, -3/2)$ is strictly feasible and

$$T = \{x : 1/4 \leq x_1 \leq 1, -1 \leq x_1x_2 \leq -1/2\}$$

is closed and bounded. A nonnegative aggregate is convex exactly when $\lambda_1 = \lambda_3$, and then equals

$$\lambda_2(x_1^2 - x_1) + \lambda_4(1/4 - x_1) - \lambda_1.$$

It follows that $C_{\text{agg}} = [1/4, 1] \times \mathbb{R}$. In contrast,

$$P_{\text{SDP}} = ([1/4, 1) \times \mathbb{R}) \cup (\{1\} \times [-1, -1/2]).$$

Indeed the linear fourth row requires $x_1 \geq 1/4$; for $x_1 < 1$ the preceding lift works with $X_{12} = -3/4$ and $Y_{22} = (-3/4 - x_1x_2)^2 / ((x_1 - x_1^2)/2)$. At $x_1 = 1$, $Y_{11} = Y_{12} = 0$, so the first and third rows require $-1 \leq x_2 \leq -1/2$, and a rank-one lift suffices. This compact variant is used only to demonstrate that a closure qualification is necessary; no HHC claim is needed for it.

In particular, our use of the classical SDP/aggregation relationship is the closure equality in Proposition 3.4 and Fujie and Kojima [9, Theorem 2.1]. We do not use the unqualified projection equality stated in Kojima and Tunçel [10, Theorem 4.2]; Example 3.6 explains the distinction, including when the original feasible set is compact. For that compact variant, the cone generated by its four defining quadratic functions is a closed convex cone by finite generation (Lemma 2.7). Replacing the finite list by this cone leaves the original feasible set, the semidefinite projection, and the convex-aggregation intersection unchanged: the original rows belong to the cone, and every cone member is a nonnegative combination of them. Thus it also meets the closed-convex-cone representation hypothesis in that theorem.

4 Checkable hypotheses and their limits

4.1 Stable convexity of the quadratic parts

A tuple $(A_1, \dots, A_m)$ is *stably convex* if it has a neighborhood in $(\mathbb{S}^n)^m$ such that every tuple $(M_1, \dots, M_m)$ in that neighborhood has a convex joint image $\{(x^\top M_i x)_i : x \in \mathbb{R}^n\}$. This is stable convexity in the sense of Sheriff [12], rather than convexity of each component function.

Proposition 4.1. *If the quadratic-part tuple $(A_i)_i$ is stably convex, then f^h has asymptotic hyperplane convexity, for arbitrary b_i, c_i . Consequently the certificate equivalence holds whenever $S \neq \emptyset$ and either*

1. $m = 2$, in any dimension $n \geq 1$; or
2. $m = 3$, $n \geq 3$, and some real combination $\sum_i \theta_i A_i$ is positive definite.

Proof. For $s \neq 0$, parameterizing $H_{\alpha, s}$ by $(x, \alpha^\top x/s)$ makes its image the joint image of

$$M_i(s) = A_i + s^{-1}(\alpha b_i^\top + b_i \alpha^\top) + s^{-2} c_i \alpha \alpha^\top. \quad (4.1)$$

For each fixed α , these matrices tend to A_i , so stable convexity gives convexity for all sufficiently large s . Dines' theorem gives convexity for every two-form tuple, including every perturbation and hyperplane restriction. For three forms on $\mathbb{R}^n$, $n \geq 3$, Polyak's theorem states that a positive definite real combination implies convexity of the joint image [11, Theorem 2.1]. Positive definiteness persists under small perturbations, so this gives stable convexity and the claimed conclusion. $\square$

The first case is the known two-constraint certificate setting. The second is a convenient criterion on the quadratic blocks, not a new three-form convexity theorem. One must also distinguish positive definiteness in $\text{span}\{A_i\}$ from positive definiteness in $\text{span}\{Q_i\}$. The latter implies the former, but in the delicate case of a trivial certificate they are equivalent. Indeed, if $\lambda^0 \geq 0$, $\lambda^0 \neq 0$, and $A_{\lambda^0} = b_{\lambda^0} = 0$, strict feasibility gives $c_{\lambda^0} < 0$. Thus $e_{n+1} e_{n+1}^\top = Q_{\lambda^0} / c_{\lambda^0}$ belongs to the real span of the Q_i . If $\sum_i \theta_i Q_i = \begin{pmatrix} A & b \\ b^\top & c \end{pmatrix}$ has $A \succ 0$, adding $r e_{n+1} e_{n+1}^\top$ with $r > b^\top A^{-1} b - c$ makes it positive definite by the Schur complement. If no nonzero trivial certificate exists, either a nontrivial one already exists or $\mathcal{K} = \{0\}$; under convexity of the quadratic-part image, the latter case was covered in Blekherman et al. [4, Proposition 2.14]. Thus the block formulation should not be read as resolving an additional previously open PDLC case beyond the main certificate theorem.

Example 4.2 (Asymptotic hyperplane convexity is strictly weaker than HHC). In $\mathbb{R}^3$, set

$$\begin{aligned} f_1 &= x_1^2 + x_2^2 + x_3^2 - 1, \\ f_2 &= x_1^2 - x_2^2 + x_3^2 - 1, \\ f_3 &= 2x_1 x_2 + x_3^2 - 1. \end{aligned}$$

The origin is strictly feasible. Since $A_1 = I$, the tuple is stably convex by the three-form result, and Proposition 4.1 gives asymptotic hyperplane convexity. On the homogeneous hyperplane $x_3 = t$, however, the image is

$$\{(x_1^2 + x_2^2, x_1^2 - x_2^2, 2x_1 x_2) : x_1, x_2 \in \mathbb{R}\}.$$

Every image point (y_1, y_2, y_3) satisfies $y_1^2 = y_2^2 + y_3^2$. The image contains $(1, 1, 0)$ and $(1, -1, 0)$ but not their midpoint $(1, 0, 0)$. Thus HHC fails. This also proves that stable convexity of the quadratic parts does not imply HHC of the homogenization.

Example 4.3 (HHC need not give stable convexity of the parts). Let $f_1 = f_2 = f_3 = x_1^2 - x_2^2 - 1$ in $\mathbb{R}^3$. On every homogeneous hyperplane the three components agree. A single real homogeneous quadratic form has image $\{0\}$, a ray, or $\mathbb{R}$, as follows by scaling and, in the indefinite case, evaluating both signs. Hence every such hyperplane image is convex. For any $\varepsilon > 0$, perturb the three quadratic parts to

$$u = x_1^2 - x_2^2, \quad u + \varepsilon(x_1^2 + x_2^2), \quad u + 2\varepsilon x_1 x_2.$$

Their values at e_1, e_2 have midpoint $(0, \varepsilon, 0)$. A preimage would require $x_1^2 = x_2^2 = 1/2$ and $x_1 x_2 = 0$, which is impossible. These arbitrarily small perturbations are nonconvex. Together with Example 4.2, this separates the two sufficient conditions in both directions.

4.2 Why ordinary hidden convexity is insufficient

Example 4.4. Consider, in $\mathbb{R}^3$ with x_3 free,

$$f_1 = x_1^2 - x_2^2 - 1, \quad f_2 = x_2^2 - x_1^2 - 1, \quad f_3 = x_1 x_2 - 1, \quad f_4 = -x_1 x_2 - 1. \quad (4.2)$$

The strict set contains the origin. On S ,

$$(x_1^2 + x_2^2)^2 = (x_1^2 - x_2^2)^2 + 4x_1^2 x_2^2 < 5.$$

It lies in the convex cylinder $\{x : x_1^2 + x_2^2 < \sqrt{5}\}$, so its hull is proper. Every aggregated quadratic matrix is traceless and every linear part is zero. Positive semidefiniteness therefore forces the matrix to vanish, so all convex certificates are trivial. More explicitly, the only conditions are $\lambda_1 = \lambda_2$ and $\lambda_3 = \lambda_4$.

Nevertheless both the full homogeneous image and the image on E are convex. Put $u = x_1^2 - x_2^2$, $v = x_1 x_2$, and $w = t^2$. The complex square map shows that (u, v) ranges over $\mathbb{R}^2$; w ranges independently over $\mathbb{R}_+$. Hence the full image is the linear image of $\mathbb{R}^2 \times \mathbb{R}_+$ under

$$(u, v, w) \mapsto (u - w, -u - w, v - w, -v - w).$$

The image on E is its restriction to $w = 0$ and is convex too.

For $s \neq 0$ restrict instead to $x_1 = st$. The same linear map is injective, and $w = x_1^2/s^2$. The two preimages $(x_1, x_2, x_3, t) = (1, 0, 0, 1/s)$ and $(0, 1, 0, 0)$ give a midpoint whose (u, v, w) coordinates would have to be $(0, 0, 1/(2s^2))$. They would force $x_1^2 = x_2^2 = 1/2$ and $x_1 x_2 = 0$, a contradiction. Thus no nonzero level in this family has a convex image. This is the precise obstruction to asymptotic hyperplane convexity. Proposition 3.4 also gives $P_{\text{SDP}} = \mathbb{R}^3$ here, although the true hull is bounded in two coordinates.

Three inequalities already suffice for a proper-hull obstruction: delete f_4 . The trace argument and both ordinary hidden-convexity statements remain valid. With $p = x_1 + x_2$ and $q = x_1 - x_2$, strict feasibility requires $|pq| < 1$ and $p^2 - q^2 < 4$. If $|p| \geq 3$, then $|q| < 1/3$ and $p^2 < 4 + 1/9$, a contradiction. Thus $|p| < 3$ is a proper affine bound. The set is unbounded, since $x_1 = -x_2$ is feasible for every value of x_1 . The three-coordinate map $(u, v, w) \mapsto (u - w, -u - w, v - w)$ is also injective, so the preceding hyperplane obstruction persists.

4.3 The role of strict feasibility for closed systems

Example 4.5. In $\mathbb{R}^3$, let

$$f_1 = x_1 x_2 - 1, \quad f_2 = -x_1 x_2 + 1, \quad f_3 = x_1^2 - x_2^2.$$

Then $S = \emptyset$, whereas

$$T = \{x : x_1 x_2 = 1, |x_1| \leq |x_2|\}$$

is nonempty and has $|x_1| \leq 1$. Consequently its hull is proper. The convex hull of the four feasible planar points $\pm(1, 1), \pm(1/2, 2)$ contains a neighborhood of the origin: the two positive points are linearly independent and their symmetric convex hull is a nondegenerate parallelogram. Since x_3 is free, $0 \in G := \text{int conv } T$ and G has full dimension.

The quadratic matrices are traceless and the linear parts vanish, so every convex certificate is trivial. HHC holds because $Q_2 = -Q_1$; every hyperplane image is a linear image of a two-form image, which is convex by Dines' theorem. This verifies all claimed properties without a sampled eigenvalue calculation.

This example also shows that the condition $Q_\lambda \neq 0$ for all nonzero $\lambda \geq 0$ in Blekherman et al. [4, Theorem 2.24 and Remark 2.25] cannot simply be removed. In their notation a good closed-system multiplier must satisfy $G \subseteq \{f_\lambda < 0\}$ and give Q_λ at most one negative eigenvalue. Since $0 \in G$, any such multiplier would have $c_\lambda = \lambda_2 - \lambda_1 < 0$. Put $\mu = \lambda_1 - \lambda_2 > 0$. Its homogeneous matrix is

$$Q_\lambda = \text{diag} \left(\begin{pmatrix} \lambda_3 & \mu/2 \\ \mu/2 & -\lambda_3 \end{pmatrix}, 0, -\mu \right).$$

The 2×2 block has determinant $-\lambda_3^2 - \mu^2/4 < 0$, so it contributes one negative eigenvalue; $-\mu$ contributes a second. There are therefore no good closed-system multipliers, whereas $\emptyset \neq G \neq \mathbb{R}^3$. Also $Q_{(1,1,0)} = 0$. Thus even taking an interior of the intersection of the good aggregations would give the whole space, not G .

4.4 A certificate need not describe the hull

Example 4.6. In $\mathbb{R}^3$ with x_2, x_3 free, let

$$f_1 = x_1^2 - 4, \quad f_2 = -x_1^2 + 2x_1 + 3.$$

Dines' theorem gives HHC. Factoring the two inequalities gives $S = \{x : -2 < x_1 < -1\}$, which is already convex. Globally convex nonzero aggregations have $\lambda_1 \geq \lambda_2 \geq 0$, with $\lambda_1 > 0$, and hence

$$f_\lambda(0) = -4\lambda_1 + 3\lambda_2 < 0.$$

Thus even all strict globally convex aggregations admit the origin, although $x_1 < -1$ is valid for S . The Shor projection also admits $x = 0$, with $X = \text{diag}(7/2, 0, 0)$: the two lifted residuals both equal $-1/2$. The relaxation is still proper, since the first convex quadratic inequality confines $|x_1| \leq 2$. Therefore the whole-space equivalence does not give hull exactness.

These examples distinguish the hypotheses, the open and closed systems, and the strength of the resulting certificate. The sufficient conditions in Proposition 4.1 supply concrete classes; no general procedure for verifying HHC or asymptotic hyperplane convexity is furnished by the certificate theorem.

5 Formal verification of the certificate theorem

A Lean development verifies Theorem 2.2 and its HHC specialization, Corollary 2.3, for finite families of real symmetric matrices, real vectors, and real constants. Its strict feasible set, ordinary convex hull, and nontrivial certificate agree with those defined here. The main declaration

is `proper_hull_iff_certificate` in the namespace `QuadraticAggregation.System`; the HHC specialization is `proper_hull_iff_certificate_of_hhc`. The assumptions are nonemptiness and AHC, or nonemptiness and HHC, respectively. No positive definite combination is assumed. The certificate-to-proper-hull implication is proved separately without either assumption.

The formal definitions express HHC through kernels of nonzero linear functionals, which are precisely the linear hyperplanes in this finite-dimensional setting. AHC is stated using arbitrarily large good levels, as in Definition 2.1; its equivalence with the increasing sequence formulation is also proved. Matrix positive semidefiniteness is linked to nonnegativity of the quadratic form, using the symmetry of the original matrices and their aggregations.

The proof follows the constant-elimination and coefficient-normalization argument in Section 2. In particular, it constructs separation weights from the convex hyperplane images and obtains the nonzero limit in the actual finitely generated coefficient cone. These are proved steps, not additional hypotheses of the headline theorem. The implementation uses the entrywise maximum matrix norm and the product maximum norm for coefficient pairs. These are equivalent to the Frobenius and Euclidean norms in finite dimension; the compactness proof works in either choice.

The verified supporting results include Lemmas 2.4, 2.5, 2.6, and 2.7, and the uniform cone-separation lemma, Lemma 2.10. The last lemma is checked independently, although the main formal proof does not use it. The spectral distance calculation and quantitative alternative in Section 2.3 are not formalized as separate results.

The sources consist of eleven modules in `Formal/QuadraticAggregation/`, accompanied by a declaration coverage map and reproducible checks. With the pinned Lean 4.33.1 toolchain, targeted compilation without warnings, a transitive axiom audit of all 178 declarations owned by those modules, and kernel replay of each module passed. The checks were rerun during preparation of this manuscript, with source fingerprints unchanged throughout the run. The only permitted axioms were `propext`, `Classical.choice`, and `Quot.sound`. Kernel replay uses the pinned Lean kernel and imported dependencies; it does not constitute verification by an independently implemented proof assistant or a replay of all dependencies from source.

This formal scope concerns existence of a nontrivial globally convex aggregation. It does not establish that such aggregations describe the entire hull. The closed-system and relaxation consequences, SDP characterizations, examples, applications, further aggregation results, and literature-priority claims are outside the formal package. Formal verification of the stated theorem does not certify the rest of the manuscript.

A An elementary application to local infeasibility certificates

Convex aggregation can also turn a local affine proof into a globally valid convex inequality. We give the exact certificate and a small example to delimit this use. These are applications of established aggregation and convexification principles, not an algorithmic novelty or a performance claim. In related conflict analysis, Berthold and Witzig [1] use local affine proofs and aggregate convex nonlinear relaxation rows. Dong [8] explicitly allows aggregation of original quadratic inequalities or quadratic rows reconstructed from a lift before convexification. The construction below selects nonnegative weights for the original quadratic rows while retaining an affine box margin.

A.1 Validity and a linear selection problem

Let $f_i(x) \leq 0$ be globally active quadratic rows as in (1.1). On a nonempty bounded box $B = [l, u]$, suppose the affine functions $\ell_i(x) = a_i^\top x + d_i$ satisfy $\ell_i(x) \leq f_i(x)$ for all $x \in B$. For $\lambda \geq 0$, put $L_\lambda = \sum_i \lambda_i \ell_i$.

Proposition A.1. *If $\min_{x \in B} L_\lambda(x) \geq \delta > 0$, the box B contains no point satisfying all the original rows. If also $A_\lambda \succeq 0$, then, for every $\bar{x} \in \mathbb{R}^n$,*

$$f_\lambda(\bar{x}) + \nabla f_\lambda(\bar{x})^\top (x - \bar{x}) \leq 0 \quad (\text{A.1})$$

is valid for every original feasible point. It separates every expansion point $\bar{x} \in B$. If $\bar{x}$ minimizes f_λ on B , this one tangent excludes the entire box.

Proof. Original feasibility implies $f_\lambda \leq 0$ everywhere on the feasible set, whereas on B the estimator inequality gives $f_\lambda \geq L_\lambda \geq \delta > 0$. For a convex quadratic, subtraction of its tangent at $\bar{x}$ leaves $(x - \bar{x})^\top A_\lambda (x - \bar{x}) \geq 0$. This proves global validity and separation at any $\bar{x} \in B$. A minimizer on the compact box exists, and the convex first-order condition gives $\nabla f_\lambda(\bar{x})^\top (x - \bar{x}) \geq 0$ for all $x \in B$. Its tangent is therefore at least δ throughout the box. $\square$

An arbitrary tangent separates its expansion point but need not exclude all of B . The quadratic aggregate remains globally valid even though the estimators used to select its weights are local. When several estimators refer to the same original row, their weights must be summed in reconstructing that row's coefficient.

A sufficient positive-semidefiniteness condition is symmetric diagonal dominance with nonnegative diagonal. Let $a(\lambda) = \sum_i \lambda_i a_i$ and $d(\lambda) = \sum_i \lambda_i d_i$. Consider the linear program

$$\begin{aligned} & \max_{\lambda, t, v, \delta} \quad \delta \\ \text{subject to} \quad & \lambda \geq 0, \quad \sum_i \lambda_i = 1, \\ & t_{jk} \geq (A_\lambda)_{jk}, \quad t_{jk} \geq -(A_\lambda)_{jk} \quad (j < k), \\ & (A_\lambda)_{jj} \geq \sum_{k < j} t_{kj} + \sum_{k > j} t_{jk} \quad (j = 1, \dots, n), \\ & v_j \leq 0, \quad v_j \leq a_j(\lambda) \quad (j = 1, \dots, n), \\ & \delta \leq d(\lambda) + l^\top a(\lambda) + \sum_j (u_j - l_j) v_j. \end{aligned} \quad (\text{A.2})$$

The last expression is a lower bound on the exact affine minimum

$$\min_{x \in B} L_\lambda(x) = d(\lambda) + l^\top a(\lambda) + \sum_j (u_j - l_j) \min\{0, a_j(\lambda)\}. \quad (\text{A.3})$$

It attains that minimum by choosing $v_j = \min\{0, a_j(\lambda)\}$. Likewise the absolute-value lifts can be chosen as $t_{jk} = |(A_\lambda)_{jk}|$. Thus (A.2) optimizes exactly the affine margin over the normalized diagonally dominant aggregates. Any feasible solution with certified $\delta > 0$ suffices for validity; no optimality certificate is needed. The elementary identity

$$x^\top A x = \sum_j \left(A_{jj} - \sum_{k \neq j} |A_{jk}| \right) x_j^2 + \sum_{j < k} |A_{jk}| (x_j + \text{sign}(A_{jk}) x_k)^2 \quad (\text{A.4})$$

proves positive semidefiniteness whenever the diagonal dominance conditions hold. Zero off-diagonal terms may be omitted. For rational data and rational weights, all coefficients and the margin can be checked exactly. Diagonal dominance is sufficient, not necessary, and depends on the chosen coordinates.

Affine equalities $h_e(x) = 0$ may be included with signed weights μ_e . Add $\sum_e \mu_e h_e$ to both the original aggregate and its affine estimator. To normalize the selection problem, write $\mu_e = \mu_e^+ - \mu_e^-$, with nonnegative parts, and replace the simplex equation by

$$\sum_i \lambda_i + \sum_e (\mu_e^+ + \mu_e^-) = 1.$$

This permits equality-only certificates and bounds every multiplier. Normalizing only the inequality weights is insufficient: with the row $-1 \leq 0$, equality $x = 0$, and $B = [1, 2]$, the aggregate $-1 + \mu x$ has margin $\mu - 1$ for $\mu \geq 0$. Its margin is unbounded if μ is left unnormalized, although each individual certificate is valid. The equality extension is for affine equalities, whose signed affine estimators are exact.

If original rows are imposed only when specified logical conditions are active, their aggregate is valid under the conjunction of the conditions supporting its nonzero weights. This condition must be retained in any resulting cut. Global unconditional validity in Proposition A.1 uses globally active rows.

A.2 A complete rational example

Let $0 \leq x, y \leq 1$ and impose

$$f_1 = x^2 + 2xy - \frac{3}{50} \leq 0, \quad f_2 = 2y^2 - xy - \frac{9}{100} \leq 0. \quad (\text{A.5})$$

The two quadratic matrices have determinants -1 and $-1/4$, so both are indefinite. The relaxation with square variables s_x, s_y and product variable w imposes the full square envelopes and the four McCormick inequalities,

$$x^2 \leq s_x \leq x, \quad y^2 \leq s_y \leq y, \quad \max\{0, x + y - 1\} \leq w \leq \min\{x, y\},$$

along with $s_x + 2w \leq 3/50$ and $2s_y - w \leq 9/100$. It admits $(x, y, s_x, s_y, w) = (1/5, 1/5, 1/25, 1/25, 0)$. This is the complete termwise convex relaxation on the root box; the failure is not due to missing square tangents.

On the smaller box $B = [1/5, 1]^2$, take

$$\ell_1 = \frac{4}{5}x + \frac{2}{5}y - \frac{9}{50}, \quad \ell_2 = -x + \frac{3}{5}y + \frac{3}{100}.$$

Their validity follows from the polynomial identities

$$\begin{aligned} f_1 - \ell_1 &= (x - 1/5)^2 + 2(x - 1/5)(y - 1/5), \\ f_2 - \ell_2 &= 2(y - 1/5)^2 + (1 - y)(x - 1/5), \end{aligned}$$

whose right sides are nonnegative on B . We have

$$2\ell_1 + \ell_2 = \frac{3}{5}x + \frac{7}{5}y - \frac{33}{100} \geq \frac{7}{100} > 0.$$

The original aggregate is

$$G = 2f_1 + f_2 = 2x^2 + 3xy + 2y^2 - \frac{21}{100} = \frac{1}{2}x^2 + \frac{1}{2}y^2 + \frac{3}{2}(x + y)^2 - \frac{21}{100}.$$

Its quadratic matrix is positive definite, with determinant $7/4$. At $\bar{x} = \bar{y} = 1/5$, the value of G is $7/100$ and its gradient is $(7/5, 7/5)$. Its tangent inequality is exactly

$$x + y \leq \frac{7}{20}. \quad (\text{A.6})$$

It excludes the root relaxation witness and the entire smaller box, where $x + y \geq 2/5 > 7/20$.

The origin and $(1/5, 0)$, $(0, 1/5)$ satisfy both original rows, so the system is globally feasible. With binary variables z_x, z_y and the additional inequalities $x \geq z_x/5$, $y \geq z_y/5$, either single activation is feasible but joint activation is infeasible. Also $\min_B \ell_1 = 3/50 > 0$: a one-row local proof may use only the first, indefinite row, whereas the displayed convex aggregate uses both rows. No general support-selection guarantee follows.

For completeness the selection LP has an exact solution here. Write $\lambda = (a, 1 - a)$, $0 \leq a \leq 1$. Its matrix is

$$A_\lambda = \begin{pmatrix} a & (3a - 1)/2 \\ (3a - 1)/2 & 2(1 - a) \end{pmatrix}.$$

The inequalities $a \geq |(3a - 1)/2|$ and $2(1 - a) \geq |(3a - 1)/2|$ reduce to $1/5 \leq a \leq 5/7$. The affine coefficients of L_λ are $(9a/5 - 1, 3/5 - a/5)$, and its constant is $(3 - 21a)/100$. The second coefficient is positive, while the first changes sign at $a = 5/9$. Formula (A.3) gives

$$\min_B L_\lambda = \begin{cases} (31a - 17)/20, & a \leq 5/9, \\ (11a - 5)/100, & a \geq 5/9. \end{cases}$$

Both pieces are strictly increasing and agree at their common endpoint. Thus the exact optimum is $\lambda = (5/7, 2/7)$ with normalized margin $1/35$. The illustrative weights $(2, 1)$ have normalized margin $7/300$; their unnormalized margin $7/100$ is not comparable to $1/35$.

A.3 Limits of this application

If the quadratic rows are purely bilinear, their quadratic matrices have zero diagonal. Any positive semidefinite aggregate then vanishes: each 2×2 principal minor gives $0 \leq -A_{jk}^2$, hence $A_{jk} = 0$ for every j, k . Signed combinations of purely bilinear equalities have the same obstruction. Cancellation can produce an affine aggregate but cannot produce nonzero positive semidefinite curvature.

Finally every positive semidefinite aggregate is implied by the common Shor lift, by (3.4); its tangents are implied as well. The example demonstrates strength beyond the termwise relaxation, not beyond the full Shor relaxation. The existence theorem does not guarantee a positive-margin diagonally dominant solution on a prescribed local support. No solver implementation or empirical performance conclusion is needed for the validity statements above.

The accompanying script `supplement/check_examples.py` uses rational arithmetic to check the displayed polynomial identities, margins, lifted witnesses, and selected matrix calculations. These checks supplement the proofs; finite test instances do not prove the universal geometric statements.

References

- [1] Timo Berthold and Jakob Witzig. Conflict analysis for MINLP. *INFORMS Journal on Computing*, 33(2):421–435, 2021. doi: 10.1287/ijoc.2020.1050.

- [2] Grigoriy Blekherman and Alex Dunbar. A topological approach to simple descriptions of convex hulls of sets defined by three quadrics. arXiv:2405.18282v1, 2024. URL <https://arxiv.org/abs/2405.18282v1>. Version of May 28, 2024.
- [3] Grigoriy Blekherman and Alex Dunbar. A topological approach to simple descriptions of convex hulls of sets defined by three quadrics. *SIAM Journal on Applied Algebra and Geometry*, 9(2): 310–342, 2025. doi: 10.1137/24M1668445.
- [4] Grigoriy Blekherman, Santanu S. Dey, and Shengding Sun. Aggregations of quadratic inequalities and hidden hyperplane convexity. arXiv:2210.01722v2, 2023. URL <https://arxiv.org/abs/2210.01722v2>. Version of May 29, 2023; numbered result locators refer to this version.
- [5] Grigoriy Blekherman, Santanu S. Dey, and Shengding Sun. Aggregations of quadratic inequalities and hidden hyperplane convexity. *SIAM Journal on Optimization*, 34(1):98–126, 2024. doi: 10.1137/22M1528215.
- [6] Santanu S. Dey, Gonzalo Muñoz, and Felipe Serrano. On obtaining the convex hull of quadratic inequalities via aggregations. *SIAM Journal on Optimization*, 32(2):659–686, 2022. doi: 10.1137/21M1428583.
- [7] Lloyd L. Dines. On the mapping of quadratic forms. *Bulletin of the American Mathematical Society*, 47:494–498, 1941. doi: 10.1090/S0002-9904-1941-07494-X.
- [8] Hongbo Dong. Relaxing nonconvex quadratic functions by multiple adaptive diagonal perturbations. *SIAM Journal on Optimization*, 26(3):1962–1985, 2016. doi: 10.1137/140960657.
- [9] Tetsuya Fujie and Masakazu Kojima. Semidefinite programming relaxation for nonconvex quadratic programs. *Journal of Global Optimization*, 10(4):367–380, 1997. doi: 10.1023/A:1008282830093.
- [10] Masakazu Kojima and Levent Tunçel. Cones of matrices and successive convex relaxations of nonconvex sets. *SIAM Journal on Optimization*, 10(3):750–778, 2000. doi: 10.1137/S1052623498336450.
- [11] Boris T. Polyak. Convexity of quadratic transformations and its use in control and optimization. *Journal of Optimization Theory and Applications*, 99(3):553–583, 1998. doi: 10.1023/A:1021798932766.
- [12] Jamin Lebbe Sheriff. *The Convexity of Quadratic Maps and the Controllability of Coupled Systems*. PhD thesis, Harvard University, 2013. URL <https://nrs.harvard.edu/urn-3:HUL.InstRepos:11030574>.
- [13] Uğur Yildiran. Convex hull of two quadratic constraints is an LMI set. *IMA Journal of Mathematical Control and Information*, 26(4):417–450, 2009. doi: 10.1093/imamci/dnp023.