




NOAA Technical Memorandum NMFS-XXX-##

NOAA quarto book

Mandy Karnauskas

January 2023

U.S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric
Administration
National Marine Fisheries Service
Northwest Fisheries Science Center



**NOAA
FISHERIES**

NOAA quarto book

Mandy Karnauskas¹

1. NOAA Fisheries, Southeast Fisheries Science Center

Table of contents

1	Introduction	1
1.1	Details	1
2	NOAA Fisheries landings data	2
2.1	Data access and upload	2
2.2	Summarize the landings	3
2.3	Plot the summary	5
2.4	Updated Gulf/Atlantic parsing of data	6
2.5	Plots with updated data	9
3	SAU	11
4	Analysis of international landings estimates	12
5	SST	13
6	Analysis of ocean temperature trends	14

List of Figures

List of Tables

1 Introduction

Here we will be displaying various data summaries and analyses related to dolphin management in the U.S. South Atlantic.

1.1 Details

The documentation of this data exploration was developed in R, using R Studio and Quarto.

2 NOAA Fisheries landings data

Here we will explore commercial and recreational landings data accessed through the FOSS (Fisheries One Stop Shop) data portal. The advantage of this portal is that commercial and recreational data are available side-by-side in the same units (pounds landed).

2.1 Data access and upload

Data are downloaded from the NOAA Fisheries FOSS site. For the Data Set, we choose both commercial and recreational. For the Years, we select all years from 1980 to the present. For NMFS Region, we select the "NMFS Regions" button and then select the Gulf, South Atlantic, Middle Atlantic and New England from the box. For the Species, we select "Dolphinfish" and "Dolphinfish**." For Report Format, we choose TOTALS BY YEAR/REGION. Hitting the "RUN REPORT" button then generates the data needed for the subsequent analysis, which can be downloaded as a .csv file under the "Actions" button.

We then read in the data file and clean it for analysis (removing the commas from the numbers so that R recognizes the values as numbers). We output the file header to make sure it was read in and that the numbers were converted correctly.

```
# clear workspace
rm(list = ls())

# read in data file
d <- read.table("data/FOSS_landings.csv", header = T, sep = ",", skip = 1)
head(d)
```

	Year	Region.Name	Pounds	Dollars	Collection	Metric.Tons
1	1980	Gulf	94,984	55,676	Commercial	43
2	1980	Middle Atlantic	400	256	Commercial	0
3	1980	New England	200	54	Commercial	0

2 NOAA Fisheries landings data

4	1980	South Atlantic	76,042	45,852	Commercial	34
5	1981	Gulf	73,075	46,951	Commercial	33
6	1981	Gulf	1,559,544		Recreational	707

```
# take commas out of file -----
d$Pounds <- as.numeric(gsub(",", "", d$Pounds))
d$Dollars <- as.numeric(gsub(",", "", d$Dollars))
d$Metric.Tons <- as.numeric(gsub(",", "", d$Metric.Tons))
head(d)
```

	Year	Region.Name	Pounds	Dollars	Collection	Metric.Tons
1	1980	Gulf	94984	55676	Commercial	43
2	1980	Middle Atlantic	400	256	Commercial	0
3	1980	New England	200	54	Commercial	0
4	1980	South Atlantic	76042	45852	Commercial	34
5	1981	Gulf	73075	46951	Commercial	33
6	1981	Gulf	1559544	NA	Recreational	707

2.2 Summarize the landings

We then summarize the landings (in pounds) by year, sector, and region. The result is a new data frame with each row representing a year and each column representing the total landings for each region and sector.

```
# compile by year for sectors and regions -----
d$Collection <- factor(d$Collection, levels = c("Recreational", "Commercial"))
tab <- tapply(d$Pounds, list(d$Year, d$Collection, d$Region.Name), sum, na.rm = T)
tab[is.na(tab)] <- 0
#head(tab)

dat <- data.frame(cbind(tab[, , "Gulf"], tab[, , "South Atlantic"],
                        tab[, , "Middle Atlantic"], tab[, , "New England"]))
names(dat) <- c("rec_Gulf", "com_Gulf", "rec_SA", "com_SA", "rec_MA", "com_MA", "rec_NE", "com_NE")
dat
```

	rec_Gulf	com_Gulf	rec_SA	com_SA	rec_MA	com_MA	rec_NE	com_NE
1980	0	94984	0	76042	0	400	0	200
1981	1559544	73075	8421898	59037	0	0	0	0

2 NOAA Fisheries landings data

1982	4197503	162457	7573035	143960	2097	200	0	0
1983	2162936	190808	9134022	127986	45927	1400	0	400
1984	1189573	282976	7265430	158901	0	1700	0	400
1985	2542089	269477	6684821	142684	113666	5000	0	4800
1986	5540003	492212	6190187	189747	125653	4200	0	200
1987	4209638	403703	6883546	212905	121433	13400	0	1100
1988	1014312	491610	3134070	212709	130240	26600	2196	17800
1989	12551461	1029628	13273504	426392	1386752	81700	437	15300
1990	13389540	1098819	9283973	514980	624928	69106	2714	14233
1991	13760600	1674124	14756442	650333	722090	90722	6294	9816
1992	12481422	714369	5893700	326831	860112	74423	1411	10315
1993	6873820	596967	7077105	520512	3001335	99420	200927	27069
1994	5448709	679903	10554836	618614	1400635	101044	12586	12328
1995	11142952	1177325	14318412	1164608	1096473	198425	24061	13811
1996	7086006	1047020	9128895	548682	1077364	43581	0	7788
1997	19958088	989648	12284366	877303	1422057	105559	0	13265
1998	9324186	422501	7312090	406368	2920384	86502	120322	11553
1999	7329953	612445	13973364	504263	1005852	98927	4458	5990
2000	5641491	582576	16049049	537077	1433991	36048	0	6545
2001	7355385	368503	14945284	470897	477081	51342	5189	29134
2002	3513226	290601	13794853	436472	1295470	83534	600948	14836
2003	4481937	310836	12112718	459351	929763	46851	129513	21275
2004	2402315	435341	9707678	555973	427145	31916	0	29042
2005	2537132	207695	10299670	376989	437472	19703	0	19982
2006	3290474	224668	12805952	433121	887684	27724	0	19645
2007	3754513	370397	13164377	740255	241212	79832	19542	44707
2008	3937153	384954	10417536	627939	264017	57764	0	16328
2009	3860527	496701	13251102	1007355	1538112	55271	0	92010
2010	1216198	111790	8543542	572031	813738	35869	0	14939
2011	3023914	309324	10774812	543811	2173136	15376	187289	26587
2012	3229090	146256	9983276	704240	755558	44918	26892	46171
2013	5425636	187966	8483569	583556	302421	41936	226988	18564
2014	3074373	201943	11925345	1041856	2129222	58868	183673	61505
2015	6286577	154133	14821772	920454	4501793	36190	1198628	37808
2016	1777348	37411	12357817	828133	2212513	61336	16499	15747
2017	2064106	65141	9457460	532973	1820541	16179	48607	30969
2018	1626095	56188	11713866	435059	3104018	20514	97107	12667
2019	3567170	83956	7967978	563132	1518629	48224	66	16918
2020	1952241	26332	5653294	260311	679556	14040	0	1880
2021	2893817	26570	6061024	178822	1051785	25318	123038	11421
2022	1664771	15906	6845573	182798	708494	17360	466897	9131

2 NOAA Fisheries landings data

2023 1427518 13305 6630236 125302 734866 28467 122550 8310

2.3 Plot the summary


Now we plot the data, using similar color schemes by region. Because the recreational data have high imprecision prior to the 1990s we plot from 1990 to the last year of data.

```
# select years -----
dat <- dat[which(rownames(dat) > 1990), ]

# define colors and plot -----
cols <- c("#0000FF30", 4, "#FF000030", 2, "#00FF0030", 3, "#00000030", 1)

par(mar = c(4, 4, 1, 0), mgp = c(2.5, 1, 0))
yrs <- as.numeric(rownames(dat))

b <- barplot(t(dat)/10^6, names.arg = yrs, col = cols, ylim = c(0, 43),
             border = NA, las = 2, ylab = "total landings (millions of pounds)",
             main = "Dolphin landings in the U.S. EEZ")
legend("top", c("New England - commercial", "New England - recreational",
               "Mid-Atlantic - commercial", "Mid-Atlantic - recreational",
               "South Atlantic - commercial", "South Atlantic - recreational",
               "Gulf of Mexico - commercial", "Gulf of Mexico - recreational"),
       col = cols[8:1], pch = 15, pt.cex = 1.5, bty = "n", y.intersp = 0.9, cex = 0.85)
axis(1, at = b, lab = rep("", length(b)), pos = 0)
abline(h=0)
```



2.4 Updated Gulf/Atlantic parsing of data

There are some issues with the way that the FOSS system splits out Gulf and Atlantic catches by Monroe County, which is on the border between the two jurisdictions. We received data from Mike Larkin (SERO) which has the correct split out of data for Monroe County between the Gulf and Atlantic. We compare the total recreational catch for the Gulf and Atlantic between the FOSS data and the updated data. The numbers are very close; the average difference from the FOSS data is 2%.

```
ml <- read.table("data/RecreationalDolphinLandings_SAandGOM_MLarkin.csv", header = T, sep = ",")
ml <- ml[which(ml$Year > 1990), ]

# select same years for FOSS data set
dat <- dat[which(yrs <= max(ml$Year)), ]

# check that years match
summary(rownames(dat) == ml$Year)
```

Mode TRUE

2 NOAA Fisheries landings data


logical 31

```
yrs <- as.numeric(rownames(dat))

# check that totals match
old <- dat$rec_Gulf + dat$rec_SA + dat$rec_MA + dat$rec_NE
new <- ml$GOM + ml$ATL

# plot totals
par(mar = c(4, 6, 2, 1), mgp = c(2.5, 1, 0))
matplot(yrs, cbind(old, new), xlab = "", lty = 1, col = c(2, 4), type = "b", pch = 19,
        ylab = "total recreational landings\n(Gulf + Atlantic, in pounds)",
        main = "Comparison of FOSS data with data corrected for Monroe County")
legend("topright", c("FOSS data", "corrected data"), col = c(2, 4), pch = 19, lty = 1)
```

Comparison of FOSS data with data corrected for Monroe



```
# average % difference between FOSS and updated data
mean((old - new)/old*100)
```

[1] -2.023886


2 NOAA Fisheries landings data

```
# min and max % difference between FOSS and updated data  
range((old - new)/old*100)
```

```
[1] -13.008552  7.587789
```

```
hist((old - new)/old*100, main = "Histogram of percent difference from FOSS data", breaks =  
      xlab = "(FOSS - update) / FOSS")
```

Histogram of percent difference from FOSS data



```
# replace FOSS data with update data  
datnew <- dat  
datnew$rec_Gulf <- ml$GOM  
datnew$rec_SA <- ml$ATL - (datnew$rec_MA + datnew$rec_NE)  
  
# old data set  
tail(dat)
```

	rec_Gulf	com_Gulf	rec_SA	com_SA	rec_MA	com_MA	rec_NE	com_NE
2016	1777348	37411	12357817	828133	2212513	61336	16499	15747
2017	2064106	65141	9457460	532973	1820541	16179	48607	30969

2 NOAA Fisheries landings data

2018	1626095	56188	11713866	435059	3104018	20514	97107	12667
2019	3567170	83956	7967978	563132	1518629	48224	66	16918
2020	1952241	26332	5653294	260311	679556	14040	0	1880
2021	2893817	26570	6061024	178822	1051785	25318	123038	11421


```
# updated data set  
tail(datnew)
```

	rec_Gulf	com_Gulf	rec_SA	com_SA	rec_MA	com_MA	rec_NE	com_NE
2016	751499	37411	13766851	828133	2212513	61336	16499	15747
2017	868489	65141	10777104	532973	1820541	16179	48607	30969
2018	508570	56188	13603875	435059	3104018	20514	97107	12667
2019	1304865	83956	10410599	563132	1518629	48224	66	16918
2020	555695	26332	7916891	260311	679556	14040	0	1880
2021	1174412	26570	7950716	178822	1051785	25318	123038	11421


2.5 Plots with updated data

We replace the FOSS values for Gulf and South Atlantic recreational catches with these new values and re-plot the full data set. We can also plot the landings as a line graph so that the relative magnitudes of catch across region and sector are more easily compared.

Dolphin landings in the U.S. EEZ -- correct Monroe Coun



Dolphin landings in the U.S. EEZ -- correct Monroe Coun



3 SAU

4 Analysis of international landings estimates

Coming soon

5 SST

6 Analysis of ocean temperature trends

Coming soon