




NOAA Technical Memorandum NMFS-XXX-##

Analyses to support the Dolphinfish Management Strategy Evaluation in the U.S. Atlantic

Marcel Kuinauskas

January 2023

U.S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric
Administration
National Marine Fisheries Service
Northwest Fisheries Science Center



**NOAA
FISHERIES**

Analyses to support the Dolphinfish Management Strategy Evaluation in the U.S. Atlantic

Mandy Karnauskas¹

1. NOAA Fisheries, Southeast Fisheries Science Center

Table of contents

1	Introduction	1
1.1	Details	1
2	Defining the regions for the operating model	2
2.1	Data download for the U.S. EEZ	2
2.2	Construct the U.S. EEZ	4
2.3	Define the Caribbean region	8
3	Define the NCA region	11
4	Define the NED region	13
5	Define the Florida Keys region	15
6	Define the North Florida - South NC region	18
7	Define the northern North Carolina region	20
8	Define the Virginia and north region	23
8.1	Plot all of the regions	24
9	NOAA Fisheries recreational landings data	27
9.1	Data access and upload	27
9.2	Separate Gulf and Atlantic recreational landings	31
9.3	Partition landings by area and sector	32
9.4	Looking at extent of discarding	39
9.5	Split the landings by season	41
9.6	Add the post-release discard mortality	47
10	NOAA Fisheries commercial data	53
10.1	Analysis of logbook data	53

Table of contents

10.2 Analysis of trip ticket data	54
11 Analysis of international catches in the high seas	57
11.1 Data access and upload	57
11.2 Read in the high seas catches	57
11.3 Summarize the high seas landings	63
11.4 Distribute the high seas landings across quarters of the year	66
12 Analysis of international catches in territorial waters	69
12.1 Data access and upload	69
12.2 Read in the catches within exclusive economic zones (EEZs)	69
12.3 Separating the EEZs by area	72
12.4 Summarize catch by regional EEZs	79
12.5 Look at Venezuela catches in detail	81
12.6 Additional error in Curacao catches	87
12.7 References	89
13 Compilation of international catches from FAO	90
13.1 Download FAO data and summarize	90
13.2 Compare FAO to SAU database	95
13.3 Create a new maximum landings scenario dataset	99
13.4 Compare landings scenarios	103
13.5 Output scenarios	105
13.6 References	106
14 Compilation of international catches	107
14.1 Compile catch in CAR and NED jurisdictional waters by discards and reporting type	107
14.2 Distribute the territorial international catches across quarters of the year	108
15 Data compilation for all fleets in the operating model	114
15.1 Data upload	116
15.2 Visualizing the final data set	121
16 Predictive indices of dolphin abundance	127
16.1 Recreational landings as a proxy for availability	127
16.2 Plot the recreational landings	128
16.3 Potential influence of El Niño Southern Oscillation	130
16.4 Seasonal influence of El Niño Southern Oscillation	133
17 Wider Atlantic temperature distribution index	137
17.1 Accessing temperature data	137

Table of contents

17.2 Analyzing high and low availability years	140
17.3 May temperature distributions	143
17.4 Quantifying the temperature distribution in an index	145
17.5 Comparison of “blue blob” index with landings	148
18 Puerto Rico tournament index	152
18.1 Upload and clean data set	152
18.2 Viewing the raw data	155
18.3 Data preparation for standardization	159
18.4 Analyze the nominal CPUE trends	165
18.5 Calculate standardized CPUE trends	171
19 Wintertime pelagic longline index	177
19.1 Analysis of logbook data	177
19.2 Model results	178

List of Figures

8.1	Areas defined in the operating model.	26
10.1	Seasonality of the pelagic longline fleet according to the logbook data. Figure shows the proportion of dolphin landings occurring in each season, for each region of the operating model.	54
10.2	Seasonality of the commercial dolphin landings according to the trip ticket data. Figure shows the proportion of dolphin landings occurring in each season, for each region of the operating model.	56
15.1	Areas defined in the operating model.	115

List of Tables

1 Introduction

Here we will be displaying various data summaries and analyses related to dolphin management in the U.S. South Atlantic.

1.1 Details

The documentation of this data exploration was developed in R, using R Studio and Quarto.

2 Defining the regions for the operating model

Here we define the different regions to be used in the operating model, based on the areas that were defined in the industry working groups. The operating model is intended to encompass the Western Central Atlantic dolphin population and include international waters (FAO areas 21 and 31) which are thought to be connected with the stock that is exploited by U.S. fisheries.

We use the U.S. exclusive economic zone (EEZ) to differentiate between national and international waters. Note that some U.S. commercial activity occurs in high seas waters and we will account for those catches in the summaries.

2.1 Data download for the U.S. EEZ

We download the U.S. Atlantic EEZ from Marineregions.org.

```
# clear workspace and load the libraries
if (!requireNamespace("maps", quietly = TRUE)) {
  install.packages("maps") }
library(maps)

rm(list = ls())

# read in the shapefile downloaded from Marineregions.org
#library(sf)
#shp_data <- st_read("data/eez/eez.shp")
#coords <- st_coordinates(shp_data)
#co <- as.data.frame(coords)
#save(co, file = "data/eez/Marineregions_EEZ.RData")

load("data/eez/Marineregions_EEZ.RData")
head(co)
```

2 Defining the regions for the operating model

	X	Y	L1	L2	L3
1	-67.28403	45.19125	1	1	1
2	-67.28400	45.19126	1	1	1
3	-67.28387	45.19125	1	1	1
4	-67.28383	45.19125	1	1	1
5	-67.28372	45.19125	1	1	1
6	-67.28365	45.19125	1	1	1


```
map("usa", xlim = c(-130, -63), ylim = c(22, 50))
axis(1); axis(2); box()
points(co$X, co$Y, pch = 19, col = 2)

# remove points that are not in the U.S. South Atlantic
co <- co[which(co$X > (-82)), ]
co <- co[-which(co$X < (-81) & co$Y > 25 & co$Y < 28), ]

#plot(co$X, co$Y, col = 0)
#text(co$X, co$Y, 1:nrow(co), cex = 0.7)
#plot(co$X, co$Y, col = 0, xlim = c(-85, -80), ylim = c(22, 25))
#text(co$X, co$Y, 1:nrow(co), cex = 0.7)

# identify the northernmost and southernmost points along the Atlantic EEZ
co1 <- co[2100:3727, ]
points(co1$X, co1$Y, col = 4, pch = 19)
```

2 Defining the regions for the operating model



2.2 Construct the U.S. EEZ

We simplify the shapefile along the U.S. coast in order to simplify computations later in the process.

```
# subset points N and S of 35N for later delineation
co2 <- co[-c(2100:3727), ]
co3 <- co2[which(co2$Y > 35), ]
co4 <- co2[which(co2$Y <= 35), ]

# reduce number of points along the coast to make computation easier
co3 <- co3[seq(1, nrow(co3), 1100), ]
co4 <- co4[seq(1, nrow(co4), 800), ]

map("usa", xlim = c(-100, -60), ylim = c(23, 50))
mtext(side = 1, "degrees W", line = 2); mtext(side = 2, "degrees N", line = 3)
mtext(side = 1, "degrees W", line = 2); mtext(side = 2, "degrees N", line = 3)
axis(1); axis(2); box()
points(co3$X, co3$Y, col = 2, pch = 19)
```


2 Defining the regions for the operating model

```
points(co4$X, co4$Y, col = 4, pch = 19)

co2 <- rbind(co3, co4)
co2 <- co2[order(co2$Y), ]
cofin <- rbind(co1, co2)

polygon(cofin$X, cofin$Y, border = 5, lwd = 3)

# make the polygon along the coast line a bit more linear
abline(h = 37.5)
abline(v = -74)
```




```
cofin <- cofin[-which(cofin$X < (-74) & cofin$Y > 37.6), ]
#text(cofin$X, cofin$Y, 1:nrow(cofin), cex = 0.9)
lis <- c(1634, 1639, 1688, 1710, 1711, 1712)
cofin <- cofin[-lis, ]

map(database = "usa", xlim = c(-70, -65), ylim = c(43, 47), col = 8)
mtext(side = 1, "degrees W", line = 2); mtext(side = 2, "degrees N", line = 3)
axis(1); axis(2, las = 2); box()
```


2 Defining the regions for the operating model

```
polygon(cofin$X, cofin$Y, border = 2, lwd = 2)
text(cofin$X, cofin$Y, 1:nrow(cofin), cex = 0.9)
```



```
map(database = "usa", xlim = c(-72, -71), ylim = c(34.8, 35.1))
mtext(side = 1, "degrees W", line = 2); mtext(side = 2, "degrees N", line = 3)
axis(1); axis(2, las = 2); box()
polygon(cofin$X, cofin$Y, border = 2)
text(cofin$X, cofin$Y, 1:nrow(cofin), cex = 0.6)
abline(h=35)
abline(v = -60)
```


2 Defining the regions for the operating model



```
cofin <- cofin[, 1:2]

map(database = "usa", xlim = c(-83, -60), ylim = c(23, 47), col = 0)
map(database = "usa", xlim = c(-83, -60), ylim = c(23, 47), add = T)
mtext(side = 1, "degrees W", line = 2); mtext(side = 2, "degrees N", line = 3)
axis(1); axis(2, las = 2); box()
polygon(cofin$X, cofin$Y, border = 2)
```

2 Defining the regions for the operating model



```
# change the southernmost tip to exactly 24N so that it aligns with Caribbean polygon
cofin <- cofin[~which(cofin$Y < 24), ]
cofin[1624, 2] <- 24

USeez <- cofin
#write.csv(USeez, file = "eez.csv")
```

Here is a final U.S. South Atlantic EEZ polygon, with a simplified coastline to reduce computational time when subsetting data points.

2.3 Define the Caribbean region

Now we define the boundaries of the Caribbean region. Its northernmost boundary aligns with the EEZ along Florida's coast up to 28.3 degrees N, at which point the boundary cuts southeast until it intersects with the 24N boundary just east of the Bahamas, approximating the Bahamas EEZ. Its western boundary is -90 degrees W and its southern boundary is 8 degrees N (essentially the land boundary of the Caribbean Sea). Its eastern boundary is -71.1 degrees W.

2 Defining the regions for the operating model


```
which.min(cofin$Y)
```

```
[1] 1624
```

```
cofin[1624, ]
```

```
      X      Y  
3723 -81.16148 24
```

```
map(database = "usa", xlim = c(-77, -76.4), ylim = c(28, 28.9))  
mtext(side = 1, "degrees W", line = 2); mtext(side = 2, "degrees N", line = 3)  
axis(1); axis(2, las = 2); box()  
polygon(cofin$X, cofin$Y, border = 2)  
text(cofin$X, cofin$Y, 1:nrow(cofin), cex = 0.6)
```




```
car1 <- cofin[1143:1624, ]  
car2 <- data.frame(cbind(c(-90, -90, -60, -60, -71.1), c(24, 8, 8, 24, 24)))  
names(car2) <- c("X", "Y")  
CAR <- rbind(car1, car2)
```

2 Defining the regions for the operating model

```
names(CAR) <- c("X", "Y")

map(database = "usa", xlim = c(-95, -60), ylim = c(5, 36))
mtext(side = 1, "degrees W", line = 2); mtext(side = 2, "degrees N", line = 3)
axis(1); axis(2, las = 2); box()
polygon(CAR$X, CAR$Y, col = 2)
```




3 Define the NCA region

The North Central Atlantic (NCA) region, otherwise known as the Western Central Atlantic, is defined according to the boundaries of FAO area 31, but excluding the U.S. EEZ and Caribbean territorial seas to the West.

```
nca1 <- cofin[569:1143, ] # subset points along EEZ
nca2 <- data.frame(cbind(c(-71.1, -60, -60, -40, -40), c(24, 24, 13, 13, 35))) # define bou
names(nca2) <- c("X", "Y")
NCA <- rbind(nca1, nca2)
names(NCA) <- c("X", "Y")

map(database = "usa", xlim = c(-85, -30), ylim = c(10, 36))
mtext(side = 1, "degrees W", line = 2); mtext(side = 2, "degrees N", line = 3)
axis(1); axis(2, las = 2); box()
polygon(NCA$X, NCA$Y, col = 3)
```

3 Define the NCA region



4 Define the NED region

The Northeast Distant Waters (NED) region, otherwise known as the Northwestern Atlantic, is defined according to the boundaries of FAO area 21, but excluding the U.S. EEZ territorial seas to the West. It does include Canadian and French territorial EEZs. The northern boundary is 60 degrees north, as dolphin are not caught above this latitude.

```
which.max(cofin$Y) # find northernmost point of U.S. EEZ
```


```
[1] 1702
```

```
cofin[1702, ]
```

```
      X      Y  
77844 -67.11553 44.84764
```

```
ned1 <- cofin[1:569, ] # subset points along EEZ  
ned2 <- data.frame(cbind(c(-42, -42, -67.1), c(35, 55, 55))) # define boundaries according  
names(ned2) <- c("X", "Y")  
NED <- rbind(ned1, ned2)  
names(NED) <- c("X", "Y")  
  
map(database = "usa", xlim = c(-85, -30), ylim = c(30, 60))  
mtext(side = 1, "degrees W", line = 2); mtext(side = 2, "degrees N", line = 3)  
axis(1); axis(2, las = 2); box()  
polygon(NED$X, NED$Y, col = 4)
```

4 Define the NED region



5 Define the Florida Keys region


The southernmost region of the U.S. EEZ is called the FLK region, and includes waters within the EEZ boundary, north up to 28 degrees N. The boundary of 28N is chosen because it aligns with commercial logbook reporting boundaries and aligns roughly with the Brevard - Indian River County border. This is convenient because U.S. recreational statistics are summarized as groups of counties, with SE FL encompassing Miami-Dade to Indian River County and NE FL encompassing Brevard to Nassau County.

```
which.min(abs(co1$Y - 28)) # find the southernmost point
```

```
[1] 1289
```

```
map(database = "usa", xlim = c(-81, -78), ylim = c(27.5, 28.9))
mtext(side = 1, "degrees W", line = 2); mtext(side = 2, "degrees N", line = 3)
axis(1); axis(2, las = 2); box()
polygon(cofin$X, cofin$Y, border = 2)
text(cofin$X, cofin$Y, 1:nrow(cofin), cex = 0.6) # find points falling at 28N
points(-80.46, 28)
cofin[1632, ] <- c(-80.46, 28)
abline(h = 28)
```


5 Define the Florida Keys region



```
FLK <- cofin[1289:1632, ] # subset points along EEZ

map(database = "usa", xlim = c(-85, -75), ylim = c(20, 30))
mtext(side = 1, "degrees W", line = 2); mtext(side = 2, "degrees N", line = 3)
axis(1); axis(2, las = 2); box()
polygon(FLK$X, FLK$Y, col = 4)
```


5 Define the Florida Keys region



6 Define the North Florida - South NC region

The second area that was decided upon to be defined within the operating model was northern Florida to Southern North Carolina. We form a boundary at 35 degrees north because this is the delineation between the NED and NCA regions. Also, 35N is just south of Cape Hatteras, which is a boundary for NMFS recreational data collection, as well as a boundary of commercial logbook data reporting. Therefore, all sources of data (international, U.S. commercial, and U.S. recreational) can be easily parsed along this boundary.


```
map(database = "usa", xlim = c(-72, -70), ylim = c(34.5, 35.4))
mtext(side = 1, "degrees W", line = 2); mtext(side = 2, "degrees N", line = 3)
axis(1); axis(2, las = 2); box()
polygon(cofin$X, cofin$Y, border = 2)
text(cofin$X, cofin$Y, 1:nrow(cofin), cex = 0.6) # find points that fall at 35N
abline(h = 35)
```



6 Define the North Florida - South NC region

```
cofin[1663, 2] <- 35
NCFL <- rbind(cofin[1632:1663, ], cofin[569:1289, ]) # subset along EEZ

map(database = "usa", xlim = c(-85, -70), ylim = c(25, 40))
mtext(side = 1, "degrees W", line = 2); mtext(side = 2, "degrees N", line = 3)
axis(1); axis(2, las = 2); box()
polygon(NCFL$X, NCFL$Y, col = 4)
```




7 Define the northern North Carolina region

The third U.S. area that was decided upon to be defined within the operating model was northern North Carolina, which was intended to encompass the northern part of the state up to the Virginia border. We form a boundary at 37 degrees north because this is a boundary for commercial logbook data reporting and roughly aligns with the latitude of the state boundary between North Carolina and Virginia. Note there is a slight misalignment between this boundary and the way that U.S. recreational statistics are reported (by state), so the commercial and recreational data will be subset along slightly different boundaries for this region.

```
map(database = "usa", xlim = c(-72, -70), ylim = c(36.5, 37.4))
mtext(side = 1, "degrees W", line = 2); mtext(side = 2, "degrees N", line = 3)
axis(1); axis(2, las = 2); box()
polygon(cofin$X, cofin$Y, border = 2)
text(cofin$X, cofin$Y, 1:nrow(cofin), cex = 0.6)
abline(h = 37)
```


7 Define the northern North Carolina region



```
cofin[425, 2] <- 37
NNC <- rbind(cofin[1663:1673, ], cofin[425:569, ])

map(database = "usa", xlim = c(-85, -70), ylim = c(25, 40))
mtext(side = 1, "degrees W", line = 2); mtext(side = 2, "degrees N", line = 3)
axis(1); axis(2, las = 2); box()
polygon(NNC$X, NNC$Y, col = 4)
```


7 Define the northern North Carolina region



8 Define the Virginia and north region

The fourth region within the U.S. EEZ is the remainder of the area, which encompasses national waters from Virginia up to Maine.

```
VBM <- rbind(cofin[1673:1702, ], cofin[1:425, ])  
  
map(database = "usa", xlim = c(-80, -60), ylim = c(30, 50))  
mtext(side = 1, "degrees W", line = 2); mtext(side = 2, "degrees N", line = 3)  
axis(1); axis(2, las = 2); box()  
polygon(VBM$X, VBM$Y, col = 4)
```



8.1 Plot all of the regions

Now we will put all of the regions on a single map and check that the boundaries align.

```
png(filename = "data/eez/new_spatial_zones1.png", width= 800, height = 900)

map(database = "world", xlim = c(-92, -38), ylim = c(5, 55), col = 8)
mtext(side = 1, "degrees W", line = 2); mtext(side = 2, "degrees N", line = 3)
axis(1); axis(2, las = 2); box()

polygon(CAR$X, CAR$Y, border = 1)
polygon(NCA$X, NCA$Y, border = 1)
polygon(NED$X, NED$Y, border = 1)
polygon(FLK$X, FLK$Y, border = 1)
polygon(NCFL$X, NCFL$Y, border = 1)
polygon(NNC$X, NNC$Y, border = 1)
polygon(VBM$X, VBM$Y, border = 1)

map(database = "world", add = T, col = 8, lwd = 2)

text(-75, 15, "CAR", cex = 1.2)
text(-57, 28, "NCA", cex = 1.2)
text(-57, 42, "NED", cex = 1.2)
text(mean(FLK$X), mean(FLK$Y), "FLK", cex = 1.2)
text(-79.3, mean(NCFL$Y), "NCFL", cex = 1.2)
text(-73.8, mean(NNC$Y), "NNC", cex = 1.2)
text(-71.5, mean(VBM$Y), "VBM", cex = 1.2)

dev.off()
```

pdf

2

```
#class(CAR)

CAR$region <- "CAR"
FLK$region <- "FLK"
NCFL$region <- "NCFL"
```

8 Define the Virginia and north region

```
NNC$region <- "NNC"
VBM$region <- "VBM"
NCA$region <- "NCA"
NED$region <- "NED"

zones <- rbind(FLK, NCFL, NNC, VBM, NED, NCA, CAR)

# plot(zones$X, zones$Y, col = as.numeric(as.factor(zones$region)))

# output the coordinates to a .csv file
write.csv(zones, file = "data/eez/dolphin_OM_areas.csv", row.names = FALSE)
```

8 Define the Virginia and north region




Figure 8.1: Areas defined in the operating model.

9 NOAA Fisheries recreational landings data

Here we will partition recreational landings data from NOAA Fisheries' Marine Recreational Information Program (MRIP), a standardized, peer-reviewed suite of recreational fishing surveys used to produce catch and effort estimates for quota monitoring and stock assessment in the United States. MRIP is a standardized survey suite that employs a complex statistical sampling design stratified across time, regions of the coast, and different fishing modes, which allows the intercept data to be scaled up by effort estimates to calculate total catches.

9.1 Data access and upload

Data are downloaded directly from the Miami server using the files provided to SERO for ACL monitoring. The current location is: M://SFD/SECM-SFD/ACL/2025_Jun23_MRIP-MRFSS/. The landings are in FES MRIP and the units are in pounds whole weight.

```
# clear workspace
rm(list = ls())

# libraries
#library(ROracle)

#con = dbConnect(dbDriver("Oracle"), username = keyring::key_list("SECPR")[1,2],
#               password = keyring::key_get("SECPR", keyring::key_list("SECPR")[1,2]), dbn

#dat <- dbGetQuery(con, "SELECT *
#                       FROM rdi.V_REC_ACL_MRIP_FES_IMP@secapxdv_dblk.sfsc.noaa.gov")

# read in data file
load("data/mrip_fes_rec81_25wv2_23June25.RData") # most recent file on SFD server
head(dat)
```

9 NOAA Fisheries recreational landings data

REC_ACL_MRIP_FES_ID	YEAR	MONTH	WAVE	SUB_REG	NEW_ST	NEW_STA	NEW_MODE	NEW_MODEN
1	73407425	1989	NA	4	6	9	NC	4 Priv
2	73407426	1989	NA	4	6	9	NC	4 Priv
3	73407427	1989	NA	4	6	9	NC	4 Priv
4	73407428	1989	NA	4	6	9	NC	4 Priv
5	73407429	1989	NA	4	6	9	NC	4 Priv
6	73407430	1989	NA	4	6	9	NC	4 Priv
AREA_X	NEW_AREA	NEW_AREAN	FL_REG	NC_REG	DS	NEW_SCI		
1	5	5	Inshore	NA	S MRIP	Centropristis	striata	
2	5	5	Inshore	NA	N MRIP	Centropristis	striata	
3	2	2	Ocean>3mi	NA	N MRIP	Centropristis	striata	
4	2	2	Ocean>3mi	NA	S MRIP	Centropristis	striata	
5	1	1	Ocean<=3mi	NA	N MRIP	Centropristis	striata	
6	1	1	Ocean<=3mi	NA	S MRIP	Centropristis	striata	
NEW_COM		SP_CODE	SPECIES_ITIS	SPECIES	AB1	B2		
1	black sea bass	8835020301	167687		34181.4033	121019.1368		
2	black sea bass	8835020301	167687		795.8612	4445.0527		
3	black sea bass	8835020301	167687		549.5390	2919.4840		
4	black sea bass	8835020301	167687		71823.2390	24548.2535		
5	black sea bass	8835020301	167687		415.7642	87.5928		
6	black sea bass	8835020301	167687		48728.5632	11859.6428		
A		B1	LBSEST_SECWWT	LBSEST_SECSOURCE	SAMPLE_SIZE_USED			
1	31761.4755	2419.9278	11457.8425	srysmwa	41			
2	584.4155	211.4457	266.7782	srysmwa	41			
3	549.5390	0.0000	484.4319	srysmwa	119			
4	33343.1844	38480.0546	63313.9276	srysmwa	119			
5	415.7642	0.0000	294.0623	srysmwa	71			
6	40585.0152	8143.5480	34464.8156	srysmwa	71			
AVG_LEN	AVG_WGT	SRYSMWA_AVG_LEN	SRYSMWA_AVG_WGT	SRYSMWA_SAMPLE_SIZE				
1	203.4572	0.3352069	203.4572	0.3352069	41			
2	203.4572	0.3352069	203.4572	0.3352069	41			
3	296.2590	0.8815243	296.2590	0.8815243	119			
4	296.2590	0.8815243	296.2590	0.8815243	119			
5	261.9949	0.7072816	261.9949	0.7072816	71			
6	261.9949	0.7072816	261.9949	0.7072816	71			
SRYSMWA_FLAGGED		SRYSMW_AVG_LEN	SRYSMW_AVG_WGT	SRYSMW_SAMPLE_SIZE				
1	NA	269.2563	0.7310037	231				
2	NA	269.2563	0.7310037	231				
3	NA	269.2563	0.7310037	231				
4	NA	269.2563	0.7310037	231				
5	NA	269.2563	0.7310037	231				

9 NOAA Fisheries recreational landings data

6	NA	269.2563	0.7310037	231			
	SRYSMW_FLAGGED	SRYSM_AVG_LEN	SRYSM_AVG_WGT	SRYSM_SAMPLE_SIZE	SRYSM_FLAGGED		
1	NA	259.7694	0.6948228	556	NA		
2	NA	259.7694	0.6948228	556	NA		
3	NA	259.7694	0.6948228	556	NA		
4	NA	259.7694	0.6948228	556	NA		
5	NA	259.7694	0.6948228	556	NA		
6	NA	259.7694	0.6948228	556	NA		
	SRYS_AVG_LEN	SRYS_AVG_WGT	SRYS_SAMPLE_SIZE	SRYS_FLAGGED	SRYS_AVG_LEN		
1	270.955	0.8099641	982	NA	269.1276		
2	270.955	0.8099641	982	NA	269.1276		
3	270.955	0.8099641	982	NA	269.1276		
4	270.955	0.8099641	982	NA	269.1276		
5	270.955	0.8099641	982	NA	269.1276		
6	270.955	0.8099641	982	NA	269.1276		
	SRYS_AVG_WGT	SRYS_SAMPLE_SIZE	SRYS_FLAGGED	SRYS_AVG_LEN	SRYS_AVG_WGT	SRYS_SAMPLE_SIZE	
1	0.8085211	1703	NA	303.6871	1.050403	46055	
2	0.8085211	1703	NA	303.6871	1.050403	46055	
3	0.8085211	1703	NA	303.6871	1.050403	46055	
4	0.8085211	1703	NA	303.6871	1.050403	46055	
5	0.8085211	1703	NA	303.6871	1.050403	46055	
6	0.8085211	1703	NA	303.6871	1.050403	46055	
	SR_FLAGGED	S_AVG_LEN	S_AVG_WGT	S_SAMPLE_SIZE	S_FLAGGED	VAR_B2	VAR_AB1
1	NA	344.4373	1.364082	305089	NA	4072492794	310099094
2	NA	344.4373	1.364082	305089	NA	2866289	258947
3	NA	344.4373	1.364082	305089	NA	0	0
4	NA	344.4373	1.364082	305089	NA	105731413	534702898
5	NA	344.4373	1.364082	305089	NA	0	0
6	NA	344.4373	1.364082	305089	NA	22265768	286196169
	SA_LABEL	GOM_LABEL	REC_ACL	MIGRA_GRP	AGG_MODEN	JURISDICTION	PS_REG
1	Snapper	Grouper	1		Private	State	6
2	Snapper	Grouper	1		Private	State	5
3	Snapper	Grouper	1		Private	Federal	5
4	Snapper	Grouper	1		Private	Federal	6
5	Snapper	Grouper	1		Private	State	5
6	Snapper	Grouper	1		Private	State	6
	COUNCILREG	GFMC_FMP	SAFMC_FMU	LBSEST_SECGWT	AGGR_AREA	AREA	FIRST_MONTH
1	6		Y	9710.0360		NA	NA
2	5		Y	226.0832		NA	NA
3	5		Y	410.5355		NA	NA
4	6		Y	53655.8708		NA	NA

9 NOAA Fisheries recreational landings data

5	5		Y	249.2054		NA	NA	
6	6		Y	29207.4709		NA	NA	
	LAST_MONTH	ALT_FLAG	CHTS_CL	CHTS_H	CHTS_RL	CHTS_WAB1C	CHTS_WAB1H	MODE_FX
1	NA	0	0	0	0	0	0	7
2	NA	0	0	0	0	0	0	7
3	NA	0	0	0	0	0	0	7
4	NA	0	0	0	0	0	0	7
5	NA	0	0	0	0	0	0	7
6	NA	0	0	0	0	0	0	7
	NEW_SEASN	SEASON						
1		NA						
2		NA						
3		NA						
4		NA						
5		NA						
6		NA						

```
#apply(dat[2:18], 2, table, useNA = "always")
```

```
# confirm nomenclature for dolphin and subset by species -----
```

```
table(dat$NEW_COM[grepl("dolphin", dat$NEW_COM)])
```

```
dolphin
9377
```

```
table(dat$NEW_SCI[grepl("dolphin", dat$NEW_COM)])
```

```
Coryphaena hippurus
9377
```

```
d <- dat[which(dat$NEW_COM == "dolphin"), ]
#apply(d[2:18], 2, table, useNA = "always")
d <- d[which(d$YEAR < 2025), ] # take out 2025 because incomplete year
tot <- tapply(d$LBSEST_SECWWT, d$YEAR, sum, na.rm = T)
```

9.2 Separate Gulf and Atlantic recreational landings

Now we filter the data set to separate out the Gulf versus Atlantic landings.

```
# codes for WFL regions: 1 (NW FL Panhandle- Escambia to Dixie co.), 2 (SW FL Peninsula- Le
# subregional codes: 6 is Atlantic and 7 is Gulf
table(d$FL_REG, d$SUB_REG, useNA = "always")
```

	4	5	6	7	<NA>
1	0	0	0	387	0
2	0	0	0	110	0
3	0	0	0	568	0
4	0	0	1022	0	0
5	0	0	312	0	0
<NA>	96	585	3836	2426	0

```
unique(d$NEW_STA)
```

```
[1] "NC"      "FLE"     "FLW"     "SC"      "GA"      "MS"      "AL"      "LA"
[9] "TX"      "FLE/GA"  "RI"      "VA"      "DE"      "MD"      "NJ"      "CT"
[17] "MA"      "NY"      "LA/MS"   "AL/FLW"
```


```
d1 <- d[which(d$COUNCILREG == 6), ]
dg <- d[which(d$COUNCILREG == 7), ]
table(d1$COUNCILREG)
```

```
6
6419
```

Now we can summarize the total dolphin landings in weight for the U.S. Atlantic, and compare to previous values sent for quota monitoring. We see that the values are almost identical.

9 NOAA Fisheries recreational landings data

```
# summarize by year and compare -----  
  
tot1 <- tapply(d1$LBSEST_SECWWT, d1$YEAR, sum, na.rm = T)  
#head(tot1)  
  
old <- read.csv("data/RecreationalDolphinLandings_SAandGOM_MLarkin.csv", skip = 6)  
plot(old$Year, old$ATL, type = "l", xlab = "year", ylab = "total landings (pounds)", lwd =  
      xlim = c(1980, 2025), ylim = c(7*10^6, 3.3*10^7), bty = "n")  
lines(as.numeric(names(tot1)), tot1, lwd = 2, col = 2, lty = 2)  
legend("topright", c("previous data", "new data"), col = c(1, 2), lwd = c(4, 2), lty = c(1,
```



9.3 Partition landings by area and sector

Now we will partition out the total landings according to the dolphin MSE areas. The areas are as follows:

- FLK: Florida Keys to Indian River County County (FL)
- NCFL: Brevard (FL) to Southern NC (South of Hatteras)
- NNC: Northern NC (North of Hatteras to NC/VA border)

9 NOAA Fisheries recreational landings data

- VBM: Virginia to Maine

```
# separate out 4 regions -----
```

```
d1$region <- "VBM"
```

```
# select for FLK
```

```
# FL_REG codes: 4: SE FL- Miami-Dade to Indian River County.; 5: NE FL- Brevard to Nassau C
```

```
table(d1$FL_REG, d1$NEW_STA, useNA = "always")
```

	CT	DE	FLE	FLE/GA	FLW	GA	MA	MD	NC	NJ	NY	RI	SC	VA
3	0	0	0	0	568	0	0	0	0	0	0	0	0	0
4	0	0	1022	0	0	0	0	0	0	0	0	0	0	0
5	0	0	312	0	0	0	0	0	0	0	0	0	0	0
<NA>	14	114	0	1794	0	54	20	125	1344	108	73	62	644	165

	<NA>
3	0
4	0
5	0
<NA>	0

```
d1$region[which(d1$FL_REG == 3 | d1$FL_REG == 4)] <- "FLK"
```

```
# select for NCFL
```

```
d1$region[which(d1$FL_REG == 5)] <- "NCFL"
```

```
lis <- c("SC", "GA", "FLE/GA") # NCFL states
```

```
d1$region[which(d1$NEW_STA %in% lis)] <- "NCFL"
```

```
# NC_REG codes: N: North of Cape Hatteras; S: South of Cape Hatteras
```

```
table(d1$NC_REG, d1$NEW_STA, useNA = "always")
```

	CT	DE	FLE	FLE/GA	FLW	GA	MA	MD	NC	NJ	NY	RI	SC	VA
	14	114	1334	1794	568	54	20	125	586	108	73	62	644	165
N	0	0	0	0	0	0	0	0	420	0	0	0	0	0
S	0	0	0	0	0	0	0	0	338	0	0	0	0	0
<NA>	0	0	0	0	0	0	0	0	0	0	0	0	0	0

9 NOAA Fisheries recreational landings data

```

      <NA>
      0
N      0
S      0
<NA>   0

```

```

d1$region[which(d1$NC == "S")] <- "NCFL"

# select for NNC
d1$region[which(d1$NC == "N")] <- "NNC"

table(d1$FL_REG, d1$NEW_STA, d1$region)

```

```
, , = FLK
```

	CT	DE	FLE	FLE/GA	FLW	GA	MA	MD	NC	NJ	NY	RI	SC	VA
3	0	0	0	0	568	0	0	0	0	0	0	0	0	0
4	0	0	1022	0	0	0	0	0	0	0	0	0	0	0
5	0	0	0	0	0	0	0	0	0	0	0	0	0	0

```
, , = NCFL
```

	CT	DE	FLE	FLE/GA	FLW	GA	MA	MD	NC	NJ	NY	RI	SC	VA
3	0	0	0	0	0	0	0	0	0	0	0	0	0	0
4	0	0	0	0	0	0	0	0	0	0	0	0	0	0
5	0	0	312	0	0	0	0	0	0	0	0	0	0	0

```
, , = NNC
```

	CT	DE	FLE	FLE/GA	FLW	GA	MA	MD	NC	NJ	NY	RI	SC	VA
3	0	0	0	0	0	0	0	0	0	0	0	0	0	0
4	0	0	0	0	0	0	0	0	0	0	0	0	0	0
5	0	0	0	0	0	0	0	0	0	0	0	0	0	0

```
, , = VBM
```

9 NOAA Fisheries recreational landings data

	CT	DE	FLE	FLE/GA	FLW	GA	MA	MD	NC	NJ	NY	RI	SC	VA
3	0	0	0	0	0	0	0	0	0	0	0	0	0	0
4	0	0	0	0	0	0	0	0	0	0	0	0	0	0
5	0	0	0	0	0	0	0	0	0	0	0	0	0	0

```
table(d1$NC_REG, d1$NEW_STA, d1$region) # check classification
```

, , = FLK

	CT	DE	FLE	FLE/GA	FLW	GA	MA	MD	NC	NJ	NY	RI	SC	VA
	0	0	1022	0	568	0	0	0	0	0	0	0	0	0
N	0	0	0	0	0	0	0	0	0	0	0	0	0	0
S	0	0	0	0	0	0	0	0	0	0	0	0	0	0

, , = NCFL

	CT	DE	FLE	FLE/GA	FLW	GA	MA	MD	NC	NJ	NY	RI	SC	VA
	0	0	312	1794	0	54	0	0	0	0	0	0	644	0
N	0	0	0	0	0	0	0	0	0	0	0	0	0	0
S	0	0	0	0	0	0	0	0	338	0	0	0	0	0

, , = NNC

	CT	DE	FLE	FLE/GA	FLW	GA	MA	MD	NC	NJ	NY	RI	SC	VA
	0	0	0	0	0	0	0	0	0	0	0	0	0	0
N	0	0	0	0	0	0	0	0	420	0	0	0	0	0
S	0	0	0	0	0	0	0	0	0	0	0	0	0	0

, , = VBM

	CT	DE	FLE	FLE/GA	FLW	GA	MA	MD	NC	NJ	NY	RI	SC	VA
	14	114	0	0	0	0	20	125	586	108	73	62	0	165
N	0	0	0	0	0	0	0	0	0	0	0	0	0	0
S	0	0	0	0	0	0	0	0	0	0	0	0	0	0

9 NOAA Fisheries recreational landings data

```
# summarize by Gulf and Atlantic regions and check against total -----
atl <- tapply(d1$LBSEST_SECWWT, list(d1$YEAR, d1$region), sum, na.rm = T)
GULF <- tapply(dg$LBSEST_SECWWT, dg$YEAR, sum, na.rm = T)

table(rownames(atl) == names(GULF))
```

```
TRUE
44
```

```
rec <- cbind(GULF, atl)

#plot(rowSums(rec), tot)
rowSums(rec)/tot
```

```
1981 1982 1983 1984 1985 1986 1987 1988 1989 1990 1991 1992 1993 1994 1995 1996
    1    1    1  NA    1    1    1    1    1    1    1    1    1    1    1    1
1997 1998 1999 2000 2001 2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012
    1    1    1    1    1    1    1    1    1    1    1    1    1    1    1    1
2013 2014 2015 2016 2017 2018 2019 2020 2021 2022 2023 2024
    1    1    1    1    1    1    1    1    1    1    1    1
```

```
write.csv(rec, file = "data/recLandings.csv")
```

Now we will separate out the landings by recreational sector and plot the totals by region and sector.

```
# separate out by for hire vs. private rec -----
d1$sector <- "Hire"

d1$sector[which(d1$NEW_MODEN == "Shore")] <- "Rec"
d1$sector[which(d1$NEW_MODEN == "Priv")] <- "Rec"

table(d1$sector, d1$NEW_MODEN, useNA = "always") # check classification
```

	Cbt	Cbt/Hbt	Hbt	Priv	Shore	<NA>
Hire	1616	119	2912	0	0	0
Rec	0	0	0	1750	22	0
<NA>	0	0	0	0	0	0

9 NOAA Fisheries recreational landings data

```
tot <- tapply(d1$LBSEST_SECWWT, list(d1$YEAR, d1$region, d1$sector), sum, na.rm = T)
head(tot)
```

, , Hire

	FLK	NCFL	NNC	VBM
1981	354639.5	74702.50	256755.24	1400.247
1982	347458.2	92271.42	27578.44	2749.224
1983	1020005.8	41784.04	88690.53	5379.177
1984	1021086.8	74658.81	NA	4203.048
1985	717523.0	172989.65	58514.13	8880.857
1986	1562354.0	63555.89	817192.26	61658.193

, , Rec

	FLK	NCFL	NNC	VBM
1981	8942641	322505.7	NA	NA
1982	8422673	1711834.7	78367.161	18727.290
1983	10000662	119370.4	44226.032	5456.237
1984	6717044	486079.7	NA	0.000
1985	7925301	1162708.4	77477.096	112944.274
1986	5699311	736872.0	2549.586	103945.134


```
fin <- data.frame(cbind(tot[, , 1], tot[, , 2]))
names(fin) <- c("Hire_FLK", "Hire_NCFL", "Hire_NNC", "Hire_VBM",
               "Rec_FLK", "Rec_NCFL", "Rec_NNC", "Rec_VBM")
head(fin)
```

	Hire_FLK	Hire_NCFL	Hire_NNC	Hire_VBM	Rec_FLK	Rec_NCFL	Rec_NNC
1981	354639.5	74702.50	256755.24	1400.247	8942641	322505.7	NA
1982	347458.2	92271.42	27578.44	2749.224	8422673	1711834.7	78367.161
1983	1020005.8	41784.04	88690.53	5379.177	10000662	119370.4	44226.032
1984	1021086.8	74658.81	NA	4203.048	6717044	486079.7	NA
1985	717523.0	172989.65	58514.13	8880.857	7925301	1162708.4	77477.096
1986	1562354.0	63555.89	817192.26	61658.193	5699311	736872.0	2549.586
	Rec_VBM						
1981	NA						
1982	18727.290						

9 NOAA Fisheries recreational landings data

```
1983 5456.237
1984 0.000
1985 112944.274
1986 103945.134
```

```
matplot(rownames(fin), fin/10^6, lty = c(rep(2, 4), rep(1, 4)),
        col = rep(1:4, 2), type = "l", lwd = 2,
        xlab = "year", ylab = "total landings (millions of pounds)")
legend("topright", colnames(fin), lty = c(rep(2, 4), rep(1, 4)), col = rep(1:4, 2), lwd = 2
```



```
# check that totals match
cor(rowSums(fin, na.rm = T), tot1)
```

```
[1] 1
```

```
mean(rowSums(fin, na.rm = T) - tot1) # very tiny differences
```

```
[1] 4.233284e-11
```

Here are the landings (in pounds) for years 1981 to 2024.


9.4 Looking at extent of discarding

Discard estimates (known as B2 estimates) are provided in numbers by the MRIP survey but are not available in the Southeast Region Headboat Survey. We can look at the overall numbers of dolphinfish caught versus reported released from the MRIP survey to get an understanding of the extent of discarding.

```
d1$perdis <- d1$B2 / (d1$AB1 + d1$B2) # percent discarded is B2 (fish released alive) / to

# calculate mean discard rate by year, region, sector -----
dis <- tapply(d1$perdis, list(d1$YEAR, d1$region, d1$sector), mean, na.rm = T)
disr <- tapply(d1$perdis, list(d1$region, d1$sector), mean, na.rm = T)

matplot(rownames(dis), cbind(dis[, , 1], dis[, , 2]), lty = c(rep(2, 4), rep(1, 4)), col =
        type = "l", lwd = 2, xlab = "year", ylab = "proportion dolphin discarded alive", yll
legend("top", c(paste("For hire - ", colnames(dis)), paste("Private - ", colnames(dis))),
        lty = c(rep(2, 4), rep(1, 4)), col = rep(1:4, 2), lwd = 2, ncol = 2, bty = "n")
```



```
cat("Discarding rates by region and sector - all years:\n")
```

Discarding rates by region and sector - all years:

```
round(disr * 100, 2) # discarding rates by region and sector
```

	Hire	Rec
FLK	10.79	18.76

9 NOAA Fisheries recreational landings data

```
NCFL  9.44 11.23
NNC    0.94  9.62
VBM    2.94 12.69
```

```
cat("Discarding rates by region and sector - last 5 years:\n")
```

Discarding rates by region and sector - last 5 years:

```
round(colMeans(tail(dis, 5)) * 100, 2) # discarding rates by region and sector - last 5 years
```

```
      Hire  Rec
FLK  21.85 36.10
NCFL 14.95 11.69
NNC   0.40  6.17
VBM   5.54 19.45
```

```
cat("Discarding rates by region and sector - most recent year:\n")
```

Discarding rates by region and sector - most recent year:

```
round(colMeans(tail(dis, 1)) * 100, 2) # discarding rates by region and sector - most recent year
```

```
      Hire  Rec
FLK  24.77 30.23
NCFL 11.65 21.46
NNC   0.94  0.00
VBM   6.24 26.86
```

```
# output discarding rates for later use
dismat <- data.frame(rownames(dis), dis[, , 1], dis[, , 2])
names(dismat) <- c("year", paste0(sort(rep(dimnames(dis)[[3]], 4)), "_", rep(dimnames(dis)[[1]], 2)))
head(dismat)
```

```
      year  Hire_FLK Hire_NCFL Hire_NNC Hire_VBM  Rec_FLK Rec_NCFL Rec_NNC
1981 1981 0.034610012      NaN      0      NaN 0.15103009      0      NA
1982 1982 0.009591116      NaN      0      NaN 0.06317636      0      0.0
1983 1983 0.004050805      0      0 0.0000000 0.04475684      0      0.0
```

9 NOAA Fisheries recreational landings data

1984	1984	0.010592562	0	NA	NaN	0.01764697	0	NA
1985	1985	0.008499863	0	0	0.11111111	0.00000000	0	0.0
1986	1986	0.020944661	0	0	0.00000000	0.05868540	0	0.5
		Rec_VBM						
1981		NA						
1982		0.00000000						
1983		0.00000000						
1984		1.00000000						
1985		0.00000000						
1986		0.07934897						

According to the MRIP survey and the reported numbers, discarding generally increases over time, excluding some highly anomalous estimates from the early years which are driven by low sample sizes. Discarding is generally higher in the private sector compared to the for-hire sector, and is highest in the Florida Keys region.

9.5 Split the landings by season

The dolphin MSE operates on a seasonal time step and thus the landings and discards must be parsed apart by individual season and year. This is not straightforward because MRIP estimates are reported in two-month waves (Jan - Feb, Mar - Apr, May - Jun, etc.), and the seasons are 3-month periods (Dec - Feb, Mar - May, Jun - Aug, Sep - Nov). For the two waves that have to be split into seasons (e.g., December is lumped with January and February; June is lumped with July and August), we split the wave 50/50%, i.e., we make the assumption that half of the landings in that wave occurs in one season and the other half occurs in the next season.

```
# define seasons -----
table(d1$WAVE, useNA = "always")
```

0	1	2	3	4	5	6	<NA>
5	407	718	1230	1439	1095	581	944

```
d1$YRWAVE <- paste0(d1$YEAR, "_", d1$WAVE)

totw <- tapply(d1$LBSEST_SECWWT, list(d1$YRWAVE, d1$region, d1$sector), sum, na.rm = T)
```

9 NOAA Fisheries recreational landings data

```
f <- data.frame(cbind(totw[, , 1], totw[, , 2]))
names(f) <- c("Hire_FLK", "Hire_NCFL", "Hire_NNC", "Hire_VBM", "Rec_FLK", "Rec_NCFL", "Rec_NNC")
head(f)
```

	Hire_FLK	Hire_NCFL	Hire_NNC	Hire_VBM	Rec_FLK	Rec_NCFL	Rec_NNC
1981_2	NA	NA	NA	NA	2793285.5	270968.46	NA
1981_3	300873.764	NA	254049.420	NA	3915698.3	NA	NA
1981_4	50017.460	NA	2705.821	NA	905621.2	51537.26	NA
1981_5	3748.301	NA	NA	NA	479834.0	NA	NA
1981_6	NA	NA	NA	NA	848201.5	NA	NA
1981_NA	NA	74702.5	NA	1400.247	NA	NA	NA

	Rec_VBM
1981_2	NA
1981_3	NA
1981_4	NA
1981_5	NA
1981_6	NA
1981_NA	NA

```
f$YEAR <- as.numeric(substr(rownames(f), 1, 4))
f$WAVE <- as.numeric(substr(rownames(f), 6, 8))
head(f) # table with landings separated by wave and year
```

	Hire_FLK	Hire_NCFL	Hire_NNC	Hire_VBM	Rec_FLK	Rec_NCFL	Rec_NNC
1981_2	NA	NA	NA	NA	2793285.5	270968.46	NA
1981_3	300873.764	NA	254049.420	NA	3915698.3	NA	NA
1981_4	50017.460	NA	2705.821	NA	905621.2	51537.26	NA
1981_5	3748.301	NA	NA	NA	479834.0	NA	NA
1981_6	NA	NA	NA	NA	848201.5	NA	NA
1981_NA	NA	74702.5	NA	1400.247	NA	NA	NA

	Rec_VBM	YEAR	WAVE
1981_2	NA	1981	2
1981_3	NA	1981	3
1981_4	NA	1981	4
1981_5	NA	1981	5
1981_6	NA	1981	6
1981_NA	NA	1981	NA

9 NOAA Fisheries recreational landings data

```
f[is.na(f)] <- 0

f <- f[which(f$YEAR >= 1985), ]
f <- f[which(f$YEAR <= 2024), ]

table(f$YEAR)
```

```
1985 1986 1987 1988 1989 1990 1991 1992 1993 1994 1995 1996 1997 1998 1999 2000
      7    7    7    7    7    7    7    7    7    7    7    7    6    6    6    6
2001 2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013 2014 2015 2016
      6    6    6    6    6    6    6    6    6    6    6    6    7    7    7    6
2017 2018 2019 2020 2021 2022 2023 2024
      6    6    6    6    6    6    6    6
```

```
table(f$WAVE)
```

```
0  1  2  3  4  5  6
15 40 40 40 40 40 40
```

```
# deal with NAs in many locations -----
# calculate percentages of landings by wave, for each region and fleet
avw <- tapply(d1$LBSEST_SECWWT, list(d1$WAVE, d1$region, d1$sector), sum, na.rm = T)
avw <- avw[-1, , ]
avw <- data.frame(cbind(avw[, , 1], avw[, , 2]))
names(avw) <- c("Hire_FLK", "Hire_NCFL", "Hire_NNC", "Hire_VBM", "Rec_FLK", "Rec_NCFL", "Re
avwp <- apply(avw, 2, function (x) x / sum(x, na.rm = T))
#colSums(avwp)

# go through each column, find NAs and fill in waves by percentages
pre_colsum <- colSums(f, na.rm = T) # column sums to check later

lis <- which(f$WAVE == 0) # list of rows with NA wave data

for (i in 1:8) { # loop through each column
  for(j in lis) { # loop through list of NA waves
    if (f[j, i] > 0) { # if there are NA wave landings, split by known proportion from oth
      miscatch <- avwp[, i] * f[j, i] # missing landings are known proportion * NA wave l
```

9 NOAA Fisheries recreational landings data

```
f[which(f$YEAR == f$YEAR[j] & f$WAVE != 0), i] <- f[which(f$YEAR == f$YEAR[j] & f$WAVE == f$WAVE[j]), i]
f[j, i] <- 0 # set to zero after landings is attributed to waves
}
}
}
f[lis, 1:8] # should be all zeros now
```

	Hire_FLK	Hire_NCFL	Hire_NNC	Hire_VBM	Rec_FLK	Rec_NCFL	Rec_NNC	Rec_VBM
1985_NA	0	0	0	0	0	0	0	0
1986_NA	0	0	0	0	0	0	0	0
1987_NA	0	0	0	0	0	0	0	0
1988_NA	0	0	0	0	0	0	0	0
1989_NA	0	0	0	0	0	0	0	0
1990_NA	0	0	0	0	0	0	0	0
1991_NA	0	0	0	0	0	0	0	0
1992_NA	0	0	0	0	0	0	0	0
1993_NA	0	0	0	0	0	0	0	0
1994_NA	0	0	0	0	0	0	0	0
1995_NA	0	0	0	0	0	0	0	0
1996_NA	0	0	0	0	0	0	0	0
2013_0	0	0	0	0	0	0	0	0
2014_0	0	0	0	0	0	0	0	0
2015_0	0	0	0	0	0	0	0	0

```
pos_colsum <- colSums(f, na.rm = T)
pre_colsum[1:8] == pos_colsum[1:8] # check that landings were parsed correctly - column sums
```

Hire_FLK	Hire_NCFL	Hire_NNC	Hire_VBM	Rec_FLK	Rec_NCFL	Rec_NNC	Rec_VBM
TRUE	TRUE	TRUE	TRUE	TRUE	TRUE	TRUE	TRUE

```
f <- f[which(f$WAVE != 0), ] # remove NA waves which now contain no landings

# split waves 3 and 6 50/50% to parse into seasons -----
lis <- which(f$WAVE == 3 | f$WAVE == 6) # list of 3 and 6 waves for all years

for (i in lis) {
  temp <- f[i, ] / 2 # go through list of waves
  temp[9] <- f$YEAR[i] # split landings in half
  temp[10] <- f$WAVE[i] # reformat year and wave data columns
}
```

9 NOAA Fisheries recreational landings data

```
temp <- rbind(temp, temp) # create two half-waves and relabel with new wave reference n
temp[1, 10] <- temp[1, 10] - 0.5
temp[2, 10] <- temp[2, 10] + 0.5
f <- rbind(f, temp) # concatenate with main data table
}

f <- f[-lis, ] # remove waves that have been split in half

pos_colsum <- colSums(f, na.rm = T)
pre_colsum[1:8] == pos_colsum[1:8] # check that landings were parsed correctly - column su
```

Hire_FLK	Hire_NCFL	Hire_NNC	Hire_VBM	Rec_FLK	Rec_NCFL	Rec_NNC	Rec_VBM
TRUE	TRUE	TRUE	TRUE	TRUE	TRUE	TRUE	TRUE

```
f <- f[order(f$YEAR, f$WAVE), ] # reorder table in chronological order

f$YEAR[which(f$WAVE == 6.5)] <- f$YEAR[which(f$WAVE == 6.5)] + 1 # add year to December wa
f$WAVE[which(f$WAVE == 6.5)] <- 0.5 # change December wave to small number so it gets summ

table(f$WAVE)
```

0.5	1	2	2.5	3.5	4	5	5.5
40	40	40	40	40	40	40	40

```
f$seas <- cut(f$WAVE, breaks = c(0, 1.5, 3, 4.5, 6))
#table(f$seas)
f$seas1 <- NA
f$seas1[which(f$seas == "(0,1.5)")] <- "DJF" # Dec Jan Feb is 0.5 and 1
f$seas1[which(f$seas == "(1.5,3)")] <- "MAM" # Mar Apr May is 2 and 2.5
f$seas1[which(f$seas == "(3,4.5)")] <- "JJA" # Jun Jul Aug is 3.5 and 4
f$seas1[which(f$seas == "(4.5,6)")] <- "SON" # Sep Oct Nov is 5 and 5.5

temp <- tapply(f[, 1], list(f$seas1, f$YEAR), sum, na.rm = T)
temp1 <- matrix(temp)
for (i in 2:8) {
  temp2 <- matrix(tapply(f[, i], list(f$seas1, f$YEAR), sum, na.rm = T))
  temp1 <- cbind(temp1, temp2)
}
```

9 NOAA Fisheries recreational landings data


```
findat <- data.frame(sort(rep(as.numeric(colnames(temp)), 4)),
                     rep(rownames(temp), ncol(temp)), temp1)
names(findat) <- c("year", "season", names(f)[1:8])

colSums(f[, 1:8], na.rm = T) == colSums(findat[, 3:10], na.rm = T) # check that total land
```

```
Hire_FLK Hire_NCFL Hire_NNC Hire_VBM Rec_FLK Rec_NCFL Rec_NNC Rec_VBM
TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE
```

We plot the results by region and sector, over the time series of year and season. We can clearly see the seasonality in the landings by region.

```
findat$yrseas <- findat$year + as.numeric(as.factor(findat$season))/4-0.125
matplot(findat$yrseas, findat[, 3:10]/10^6, lty = c(rep(2, 4), rep(1, 4)), col = rep(1:4, 2),
        xlab = "year", ylab = "total landings (millions of pounds)")
legend("topright", colnames(findat[, 3:10]), lty = c(rep(2, 4), rep(1, 4)), col = rep(1:4,
```



```
# check that totals match
cbind(colSums(findat[, 3:10], na.rm = T), colSums(fin[-c(1:5), ], na.rm = T))
```

	[,1]	[,2]
Hire_FLK	55604999	54887476
Hire_NCFL	16909916	16736926
Hire_NNC	57944903	57886388
Hire_VBM	5836953	5828072
Rec_FLK	371462933	363537632

9 NOAA Fisheries recreational landings data

```
Rec_NCFL 70803866 69641157
Rec_NNC   32556596 32479119
Rec_VBM   40081529 39968585
```

```
# the values should be close but slightly >1 because the last data set includes December 19
colSums(findat[, 3:10], na.rm = T) / colSums(fin[-c(1:5), ], na.rm = T)
```

```
Hire_FLK Hire_NCFL Hire_NNC Hire_VBM Rec_FLK Rec_NCFL Rec_NNC Rec_VBM
1.013073 1.010336 1.001011 1.001524 1.021800 1.016696 1.002385 1.002826
```

9.6 Add the post-release discard mortality

Lastly, we integrate post-release discard mortality into the total catch. MRIP reports dead discards not available for inspection within its B1 category, but there is also mortality in a proportion of fish that are released alive (B2 catch). The percent of fish released alive by year and area was calculated in section 3.4. These B2 catch estimates are only available in numbers, not weight, so in applying these percentages to the landings we must assume that fish released alive are on average the same weight as retained fish. Individual fish may be released alive if they are smaller in size or if they fall under the size limit (only Florida to South Carolina); hence it should be noted that this assumption could bias the post-release mortality estimates in weight high. We use the discard mortality rate of 0.25 as estimated by Rudershausen et al. (2019) and apply it to the percentage of live discards (B2 / total catch).

```
names(findat)[3:10] == names(dismat)[2:9] # check columns are the same
```

```
[1] TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE
```

```
findat <- findat[which(findat$year < 2025), ]
findat_disc <- findat

# loop through rows and apply discarding rate * 0.25 discard mortality rate
# LRP is Live Release Proportion = B2 / (AB1 + B2)
# solving to isolate B2 (fish released alive)
# B2 = (LRP * AB1) / (1 - LRP)

for (i in 1:nrow(findat)) {
```


9 NOAA Fisheries recreational landings data

```
LRP <- dismat[which(dismat$year == findat$year[i]), 2:9] # live release proportion
AB1 <- findat[i, 3:10] # AB1 catch in weight
B2 <- (LRP * AB1) / (1 - LRP) # B2 discards in weight
B2_dead <- B2 * 0.25 # B2 * 0.25 is estimated dead d
findat_disc[i, 3:10] <- AB1 + B2_dead # total death is AB1 + dead pro
}

# look at catch with discards
par(mfcol = c(4, 2), mar = c(3, 4, 2, 1), mgp = c(2.5, 1, 0))

for (j in 3:10) {
  barplot(rbind(tapply(findat[, j], findat$year, sum), tapply(findat_disc[, j], findat_disc$year, sum),
    col = c(2:3), border = NA,
    beside = T, las = 2, # legend.text = c("without discard mortality", "including discard mortality"),
    main = names(findat)[j], ylab = "millions of pounds")
  abline(h = 0)
  legend("topright", c("without discard mortality", "including discard mortality"), col = 2:3)
}
```

9 NOAA Fisheries recreational landings data



9 NOAA Fisheries recreational landings data

```
# check calculations - LRP = (B2_dead * 4) / (AB1 + B2_dead * 4)
out <- ((findat_disc[,3:10] - findat[,3:10]) * 4) / (findat[,3:10] + (findat_disc[,3:10] -
round(dismat[5, 2:9], 3) == round(out[2, ], 3)
```

	Hire_FLK	Hire_NCFL	Hire_NNC	Hire_VBM	Rec_FLK	Rec_NCFL	Rec_NNC	Rec_VBM
1985	TRUE	TRUE	TRUE	TRUE	TRUE	TRUE	TRUE	TRUE

```
round(dismat[nrow(dismat), 2:9], 3) == round(out[nrow(out), ], 3)
```

	Hire_FLK	Hire_NCFL	Hire_NNC	Hire_VBM	Rec_FLK	Rec_NCFL	Rec_NNC	Rec_VBM
2024	TRUE	TRUE	TRUE	TRUE	TRUE	TRUE	NA	TRUE

```
# calculate % of post-release dead discards by area
round((colSums(findat_disc[, 3:10]) / colSums(findat[, 3:10]) - 1) * 100, 2)
```

Hire_FLK	Hire_NCFL	Hire_NNC	Hire_VBM	Rec_FLK	Rec_NCFL	Rec_NNC	Rec_VBM
2.78	1.63	0.30	0.98	5.72	3.68	2.38	3.98

```
# calculate recent % of post-release dead discards by area
round((colSums(findat_disc[149:160, 3:10]) / colSums(findat[149:160, 3:10]) - 1) * 100, 2)
```

Hire_FLK	Hire_NCFL	Hire_NNC	Hire_VBM	Rec_FLK	Rec_NCFL	Rec_NNC	Rec_VBM
8.02	4.12	0.09	1.64	15.10	2.84	0.10	7.32

```
# calculate overall % of post-release dead discards
round((sum(findat_disc[, 3:10]) / sum(findat[, 3:10]) - 1) * 100, 2)
```

```
[1] 4.34
```

```
# calculate recent % of post-release dead discards
round((sum(findat_disc[149:160, 3:10]) / sum(findat[149:160, 3:10]) - 1) * 100, 2)
```

```
[1] 8.52
```

Here we will format the data for input to the operating model.

9 NOAA Fisheries recreational landings data

```
# tab3 <- findat # previous version - this is catch with no post-release discard mortality
tab3 <- findat_disc # this is catch including post-release discard mortality

flt <- c(rep("Hire", nrow(tab3)*4), rep("Rec", nrow(tab3)*4))

yrmon <- as.numeric(tab3$yrseas)
yrs <- floor(yrmon)
qrt <- (yrmon - floor(yrmon)) * 4 + 0.5
area <- c(rep("FLK", nrow(tab3)), rep("NCFL", nrow(tab3)),
          rep("NNC", nrow(tab3)), rep("VBM", nrow(tab3)))
area <- rep(area, 2)

findat1 <- data.frame(yrs, qrt, flt, area, matrix(as.matrix(tab3[,3:10])))
names(findat1) <- c("Year", "Quarter", "Fleet", "Area", "Catch_lbs")
head(findat1)
```

	Year	Quarter	Fleet	Area	Catch_lbs
1	1985	1	Hire	FLK	4471.653
2	1985	2	Hire	FLK	254030.649
3	1985	3	Hire	FLK	381627.520
4	1985	4	Hire	FLK	41558.870
5	1986	1	Hire	FLK	40762.254
6	1986	2	Hire	FLK	503865.031

```
#plot(findat1$Catch_lbs, type = "l")
#plot(findat1$Catch_lbs ~ factor(findat1$Area))
#plot(findat1$Catch_lbs ~ factor(findat1$Quarter))
#plot(findat1$Catch_lbs ~ factor(findat1$Year))

findat1 <- findat1[which(findat1$Year <= 2022), ]
findat1 <- findat1[which(findat1$Year >= 1986), ]
#apply(findat1, 2, table)

findat <- findat[which(findat$year >=1986 & findat$year <= 2022), ]
#check that the numbers are exactly the same
tapply(findat1$Catch_lbs, list(findat1$Area, findat1$Fleet), sum, na.rm = T)
```

	Hire	Rec
FLK	55478644	377580042

9 NOAA Fisheries recreational landings data

```
NCFL 16800175 68063202
NNC 56269610 32971138
VBM 5792004 39150014
```

```
colSums(findat[3:10], na.rm = T)
```

```
Hire_FLK Hire_NCFL Hire_NNC Hire_VBM Rec_FLK Rec_NCFL Rec_NNC Rec_VBM
54012811 16532764 56097190 5736592 357091651 65643193 32197622 37749487
```

```
# output final numbers in correct format
```

```
write.csv(findat1, file = "data/FINAL_files/rec_catch_TomFormat.csv", row.names = FALSE)
```

10 NOAA Fisheries commercial data

We do not display the commercial data analysis here because it is based on logbook reporting and trip ticket databases, which both contain confidential data. The code for processing the commercial data can be seen in the file “commercial.R” and the steps are outlined below.

10.1 Analysis of logbook data

We first use the pelagic longline logbook data to understand how the fleet operates in the high seas on a seasonal basis. These numbers are used later to estimate the seasonality of the international landings, because those are only reported on an annual basis but quarterly values are needed.

The steps are as follows:

1. Read in the pelagic longline data which contains set level entries including location of fishing, target species, gear used, other details of the fishing trip and number of dolphin caught, discarded alive, discarded dead, and the total pounds of dolphin kept.
2. Format the longitude and latitude of the sets into decimal degrees, and find the points that fall within each of the seven polygons representing the areas of the operating model. Check that subsetting was done correctly and add an area identifier.
3. Look at the characteristics of the dolphin catch and calculate the discard rates by trip. Dead discarding rate is found to be 1.4% on average and total discard rate is 2.9% on average.
4. Standardize date formatting and add identifier for season (quarter). Specify a new year variable to allow December to be grouped with January and February.
5. Summarize the total dolphin landings (in pounds) by year, quarter, and area.

10 NOAA Fisheries commercial data

6. Convert the total pounds by year-quarter and area to proportions such that each year-quarter combination has four seasonal proportions summing to one. Save the matrix of proportions for later analysis.




Figure 10.1: Seasonality of the pelagic longline fleet according to the logbook data. Figure shows the proportion of dolphin landings occurring in each season, for each region of the operating model.

10.2 Analysis of trip ticket data

Not all of the commercial dolphin catch activity is reported in the logbooks because dolphin are targeted via a variety of permits and gear types, not all of which have the same reporting requirements. Therefore, in order to parse the commercial landings by year, area and quarter, we use the commercial trip ticket data compiled by NOAA Fisheries. These are the same data used for quota monitoring and thus are the best representation of the total landings.

The steps are as follows:

1. Read in the trip ticket data file, which contains commercial dolphinfish landings for the Atlantic coast from 1986-2023, including all landings by year even if the gear type or area is unknown. The dataset specifies the year and month of

10 NOAA Fisheries commercial data

landing, an area of fishing, gear, and the state where landed. Pounds are in units of whole weight.

2. Read in operating model regions and classify the list of logbook areas into those seven regions. Assign each reported area to an operating model region accordingly.
3. Standardize date formatting and add identifier for season (quarter). Specify a new year variable to allow December to be grouped with January and February.
4. Summarize the landings by year-quarter, gear (PLL gear vs. other gears) and region.
5. For most combinations of year-quarter, region and gear, there are substantial landings for which the area of fishing is unknown and therefore the region cannot be defined. We assume that those landings follow the same distribution with regard to region as the other landings for that year-quarter and gear combination. For each year-quarter/gear combination, we calculate the proportion of landings that occurs in each of the seven regions. The total landings of unknown regions are multiplied by those proportions and then added to the landings from known regions. In cases where none of the areas are known for that year-quarter/gear combination, proportions from the quarter immediately previous to that quarter are used. Sum the landings by gear type to get the total landings for each year-quarter and region.
6. Reformat the data according to the specifications needed for input to the operating model (columns specifying the year, quarter, fleet, area and total landings).
7. Sum landings by year from the original data file and the final outputs to ensure that data were processed correctly and no landings are missing.




Figure 10.2: Seasonality of the commercial dolphin landings according to the trip ticket data. Figure shows the proportion of dolphin landings occurring in each season, for each region of the operating model.

11 Analysis of international catches in the high seas

Here we will explore commercial and recreational catch data as estimated by the Sea Around Us Project. This data set includes reported data for fisheries worldwide as well as reconstructed estimates for unreported catches. For each country, project scientists assemble available data on catch and use additional sociological and fishing data plus expert information and opinion, to produce their best estimates of total catch from all fishing sectors and for all species globally.

11.1 Data access and upload

The high seas data are accessed in the Basic Search High seas section of the site. The individual regions can be clicked on (e.g., Atlantic, Western Central) and then the data are downloaded using the Download Data button.

The current version of data in use is version 50.1, accessed June 2024 (the data files can be found on github in the SEFSC-dolphin-analyses/data /SAU/ folder). This was still the most recent version of data available as of September 2025.

11.2 Read in the high seas catches

First we read in the high seas catches for the Western Central Atlantic and Northwest Atlantic zones. These are equivalent to FAO areas 21 and 31.

```
# clear workspace
rm(list = ls())

if(!require("pals")) install.packages("pals")
library(pals)
```

11 Analysis of international catches in the high seas

```
# find data files
hslis <- dir("data/SAU/")[grep("HighSeas", dir("data/SAU/"))]

# identify areas 21 and 31
hslis[1]; hslis[3]
```

```
[1] "SAU HighSeas 21 v50-1.csv"
```

```
[1] "SAU HighSeas 31 v50-1.csv"
```

```
#download data
h1 <- read.csv(paste0("data/SAU/", hslis[1]))
h3 <- read.csv(paste0("data/SAU/", hslis[3]))

# confirm correct areas
#unique(h1$area_name)
#unique(h3$area_name)

hs <- rbind(h1, h3)
```

We extract the data for dolphinfish and look at the characteristics of the catches. Most of the catches are in the Western Central Atlantic, and the primary fishing entities are Venezuela and Saint Lucia. The fishing sector is all industrial, and the vast majority of the catch is reported landings, with only a tiny fraction of discards or unreported landings listed in the database. The gear type is primarily long line.

```
# check species name and filter out dolphinfish
unique(hs$common_name)[grep("dolphin", unique(hs$common_name))]
```

```
[1] "Common dolphinfish"
```

```
hs <- hs[which(hs$common_name == "Common dolphinfish"), ]

# look at the categories within each column of data
apply(hs[1:13], 2, table)
```

11 Analysis of international catches in the high seas

\$area_name

Atlantic, Northwest Atlantic, Western Central	113	160
---	-----	-----

\$area_type

high_seas
273

\$year

1985	1986	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
1	1	1	1	1	1	1	6	4	1	5	10	5	8	10	12
2013	2014	2015	2016	2017	2018	2019									
24	22	30	33	33	29	34									

\$scientific_name

Coryphaena hippurus
273

\$common_name

Common dolphinfish
273

\$functional_group

Large pelagics (≥ 90 cm)
273

\$commercial_group

Perch-likes
273

\$fishing_entity

Barbados	Belize
6	3

11 Analysis of international catches in the high seas

	Brazil	Canada
	19	18
	Guyana	Panama
	1	1
	Saint Lucia	Saint Vincent & the Grenadines
	5	28
	Spain	Suriname
	80	1
	Trinidad & Tobago	USA
	14	74
	Venezuela	
	23	
\$fishing_sector		
Industrial		
	273	
\$catch_type		
Discards Landings		
	5	268
\$reporting_status		
	Reported	Unreported
	253	20
\$gear_type		
artisanal fishing gear		
	7	
	bottom trawl	gillnet
	9	11
	hand lines	longline
	17	187
	pole and line	pots or traps
	19	9
		unknown by source
		13
\$end_use_type		
	Direct human consumption	Fishmeal and fish oil
	5	194
		37

11 Analysis of international catches in the high seas


Other

37

```
# convert from metric tonnes to pounds
hs$pounds <- hs$tonnes * 1000 * 2.20462

# produce barplot of key data fields
par(mar = c(3, 20, 3, 1), mfrow = c(3, 2), mex = 0.5)
barplot(tapply(hs$pounds/10^6, hs$area_name, sum, na.rm = T), las = 2, xlab = "millions of
#barplot(tapply(hs$pounds/10^6, hs$area_type, sum, na.rm = T), las = 2, ylab = "millions of
#barplot(tapply(hs$pounds/10^6, hs$year, sum, na.rm = T), las = 2, ylab = "millions of poun
barplot(tapply(hs$pounds/10^6, hs$fishing_entity, sum, na.rm = T), las = 2, xlab = "million
barplot(tapply(hs$pounds/10^6, hs$fishing_sector, sum, na.rm = T), las = 2, xlab = "million
barplot(tapply(hs$pounds/10^6, hs$catch_type, sum, na.rm = T), las = 2, xlab = "millions of
barplot(tapply(hs$pounds/10^6, hs$reporting_status, sum, na.rm = T), las = 2, xlab = "milli
barplot(tapply(hs$pounds/10^6, hs$gear_type, sum, na.rm = T), las = 2, xlab = "millions of
```

11 Analysis of international catches in the high seas



11 Analysis of international catches in the high seas

```
round(tapply(hs$pounds, hs$catch_type, sum, na.rm = T) / sum(hs$pounds) * 100, 2) # << 1%
```

Discards	Landings
0.03	99.97

```
round(tapply(hs$pounds, hs$reporting_status, sum, na.rm = T) / sum(hs$pounds) * 100, 2) #
```


Reported	Unreported
99.97	0.03

11.3 Summarize the high seas landings

We plot the annual landings by fishing entity to look at trends over time.

```
hs$fishing_entity <- as.factor(hs$fishing_entity)
hs$fishing_entity <- droplevels(hs$fishing_entity)

par(mar = c(5, 5, 3, 1), mfrow = c(1, 1))
tab <- tapply(hs$pounds, list(hs$year, hs$fishing_entity), sum, na.rm = T)
matplot(as.numeric(rownames(tab)), tab/10^6, type = "l", col = glasbey(ncol(tab)), lty = 1,
        main = "High seas Western Central Atlantic dolphin landings")
legend("topleft", colnames(tab), col = glasbey(ncol(tab)), lty = 1, lwd = 2, cex = 0.8, bty
```



Finally we output the subset of high seas dolphin landings for the two areas. Because we have already accounted for the USA landings in our national reporting and are aiming to only summarize international landings here, we filter out the USA landings from the Sea Around Us database.

```
hs <- hs[which(hs$fishing_entity != "USA"), ]

hs$year <- factor(hs$year, levels = 1950:2019)
tab <- tapply(hs$pounds, list(hs$year, hs$area_name), sum, na.rm = T)
tab[is.na(tab)] <- 0
tail(tab, 40)
```

	Atlantic, Northwest Atlantic,	Western Central
1980	0.000	0.000
1981	0.000	0.000
1982	0.000	0.000
1983	0.000	0.000
1984	0.000	0.000
1985	0.000	14037.129
1986	0.000	8748.399
1987	0.000	0.000

11 Analysis of international catches in the high seas

1988	0.000	0.000
1989	0.000	0.000
1990	0.000	0.000
1991	0.000	0.000
1992	0.000	0.000
1993	0.000	0.000
1994	0.000	0.000
1995	0.000	0.000
1996	0.000	0.000
1997	0.000	0.000
1998	0.000	0.000
1999	0.000	43168.296
2000	0.000	30284.627
2001	0.000	164272.223
2002	0.000	22147.477
2003	0.000	137103.181
2004	0.000	66058.258
2005	0.000	115171.863
2006	0.000	170522.101
2007	14687.527	330838.620
2008	28552.636	340863.719
2009	18719.121	662720.661
2010	4510.085	232093.837
2011	21589.424	300498.224
2012	72398.812	1059210.392
2013	34741.604	2471791.396
2014	27922.953	1247462.983
2015	90313.638	710059.125
2016	15319.433	1094893.891
2017	34806.933	1374259.739
2018	29328.358	885814.754
2019	21870.621	632747.157

```
# write summary to merge with EEZ data
write.csv(tab, file = "data/SAU/high_seas_by_year.csv")
```

11.4 Distribute the high seas landings across quarters of the year

The international landings are only reported as annual landings and do not have any month or season associated with them. However, the MSE operating model requires seasonal landings. The majority of the international landings are caught by longline, so we will assume that the fleet distribution of the U.S. pelagic longline is representative of the seasonality of the international high seas fleets. Now we parse the annual landings according to the seasonality of the U.S. fleet.

At the time of analysis, the SAU database only contained landings to 2019, so we interpolated the 2019 values to cover years 2020 - 2022.

```
# read in the percentage of the landings by area and year-quarter combinations from the U.S.
load("data/outputs/per_PLLeatch_by_area_yearquarter.RData")

tab <- tab[which(rownames(tab) >= 1997), ] # PLL data are only available from 1997 on
#tab[nrow(tab), ]
tab <- rbind(tab, tab[nrow(tab), ], tab[nrow(tab), ], tab[nrow(tab), ]) # interpolate 2019
tail(tab)
```

	Atlantic, Northwest	Atlantic, Western Central
2017	34806.93	1374259.7
2018	29328.36	885814.8
2019	21870.62	632747.2
	21870.62	632747.2
	21870.62	632747.2
	21870.62	632747.2


```
NED <- percatch[, , 1] # create matrices for final landings data
NCA <- percatch[, , 1]

for (i in 1:ncol(percatch)) {
  NED[, i] <- percatch[, i, 7] * tab[i, 1] # these are the percentages for the NED (Atlantic)
  NCA[, i] <- percatch[, i, 1] * tab[i, 2] # these are the percentages for the NCA (Western)
}

# plot the parsed out data by year-quarter
yrs <- sort(rep(as.numeric(colnames(NCA)), 4))
qrt <- rep(1:4, length(yrs)/4)
```

11 Analysis of international catches in the high seas

```
matplot(yrs + qrt/4, cbind(matrix(NED), matrix(NCA))/10^6, lty = 1, lwd = 2,
        col = 2:3, type = "l", main = "International high seas landings by quarter",
        xlab = "", ylab = "total landings (millions of pounds)")
legend("topleft", c("NED", "NCA"), lwd = 2, col = 2:3, bty = "n")
```



```
# save NED data in this format because it needs to be combined with Canadian EEZ landings
save(NED, file = "data/outputs/NED_year_quarter.RData")
```

Finally, we format the data in the format for input into the operating model. We only include NCA here, as the NED high seas landings must be combined with international landings in territorial waters of Canada and France in the NED region.

```
# format for operating model
flt <- rep("Intl", length(yrs))
are <- rep("NCA", length(yrs))
findat <- data.frame(cbind(yrs, qrt, flt, are, matrix(NCA)))
names(findat) <- c("Year", "Quarter", "Fleet", "Area", "Catch_lbs")
findat$Catch_lbs <- as.numeric(findat$Catch_lbs)
head(findat)
```

11 Analysis of international catches in the high seas

	Year	Quarter	Fleet	Area	Catch_lbs
1	1997	1	Intl	NCA	0
2	1997	2	Intl	NCA	0
3	1997	3	Intl	NCA	0
4	1997	4	Intl	NCA	0
5	1998	1	Intl	NCA	0
6	1998	2	Intl	NCA	0

```
#plot(findat$Catch_lbs, type = "l")
#plot(findat$Catch_lbs ~ factor(findat$Area))
#plot(findat$Catch_lbs ~ factor(findat$Quarter))
#plot(findat$Catch_lbs ~ factor(findat$Year))
#apply(findat, 2, table)
#barplot(tapply(findat$Catch_lbs, list(findat$Quarter, findat$Area), sum, na.rm = T), beside = TRUE)
write.csv(findat, file = "data/FINAL_files/NCA_Intl_TomFormat.csv", row.names = FALSE)
```

12 Analysis of international catches in territorial waters

12.1 Data access and upload

For catches within each country's exclusive economic zones, data are downloaded from the Sea Around Us site (Pauly, Zeller, and Palomares, 2020.). In the Biodiversity by Taxon Tools and Data, we filter by EEZ for All regions, and then filter by Taxon at the Species level for Common dolphinfish (*Coryphaena hippurus*). After clicking "View graphs" one can access the full data set using the "Download Data" button.

12.2 Read in the catches within exclusive economic zones (EEZs)

Now we read in the catches reported by each country or jurisdiction's respective EEZ. Because we have extracted dolphin catches for EEZs worldwide, we need to separate out the catches for the Atlantic Ocean basin.

```
# load libraries
library(pals)

rm(list = ls())
d <- read.csv("data/SAU/SAU Taxa 600006 v50-1.csv", header = T, sep = ",") # v. 50.1 - Jun
head(d)
```

	area_name	area_type	year	scientific_name	common_name
1	Australia	eez	1950	<i>Coryphaena hippurus</i>	Common dolphinfish
2	Australia	eez	1950	<i>Coryphaena hippurus</i>	Common dolphinfish
3	Australia	eez	1950	<i>Coryphaena hippurus</i>	Common dolphinfish
4	Bahamas	eez	1950	<i>Coryphaena hippurus</i>	Common dolphinfish
5	Barbados	eez	1950	<i>Coryphaena hippurus</i>	Common dolphinfish
6	Bermuda (UK)	eez	1950	<i>Coryphaena hippurus</i>	Common dolphinfish

12 Analysis of international catches in territorial waters

	functional_group	commercial_group	fishing_entity	fishing_sector
1	Large pelagics (>=90 cm)	Perch-likes	Australia	Industrial
2	Large pelagics (>=90 cm)	Perch-likes	Australia	Industrial
3	Large pelagics (>=90 cm)	Perch-likes	Australia	Industrial
4	Large pelagics (>=90 cm)	Perch-likes	Bahamas	Recreational
5	Large pelagics (>=90 cm)	Perch-likes	Barbados	Recreational
6	Large pelagics (>=90 cm)	Perch-likes	Bermuda (UK)	Artisanal

	catch_type	reporting_status		gear_type
1	Landings	Reported		longline
2	Landings	Reported		longline
3	Landings	Reported		longline
4	Landings	Unreported	recreational fishing gear	
5	Landings	Unreported	recreational fishing gear	
6	Landings	Reported	small scale lines	

	end_use_type	tonnes	landed_value
1	Fishmeal and fish oil	0.003373202	3.438170e-05
2	Direct human consumption	6.739657104	1.040254e+04
3	Other	0.003373202	3.438170e-05
4	Direct human consumption	48.665105143	8.262728e+00
5	Direct human consumption	0.009158559	4.846886e-04
6	Direct human consumption	1.511085514	3.199403e+03


```
#dim(d)
#apply(d[1:13], 2, table)

# convert tonnes to pounds
d$lbs <- d$tonnes * 1000 * 2.20462
```

We look at the characteristics of the catches within EEZs. These are largely industrial and artisanal in nature. Landings are the majority of the catch but discards are also present. There are more unreported landings than reported landings and a variety of gear types are used.

```
# look at characteristics of data
par(mfrow = c(2, 2), mar = c(20, 7, 3, 1), mex = 0.5, mgp = c(5.5, 1, 0))
barplot(tapply(d$lbs, d$fishing_sector, sum, na.rm = T)/10^6, las = 2, ylab = "millions of pounds")
barplot(tapply(d$lbs, d$catch_type, sum, na.rm = T)/10^6, las = 2, ylab = "millions of pounds")
barplot(tapply(d$lbs, d$reporting_status, sum, na.rm = T)/10^6, las = 2, ylab = "millions of pounds")
barplot(tapply(d$lbs, d$gear_type, sum, na.rm = T)/10^6, las = 2, ylab = "millions of pounds")
```

12 Analysis of international catches in territorial waters



```
round(tapply(d$lbs, d$catch_type, sum, na.rm = T) / sum(d$lbs) * 100, 2) # 15% are discards
```

```
Discards Landings
15.18    84.82
```

```
round(tapply(d$lbs, d$reporting_status, sum, na.rm = T) / sum(d$lbs) * 100, 2) # 55% are unreported
```

```
Reported Unreported
44.52    55.48
```

12.3 Separating the EEZs by area

Now we have to manually assign the different EEZs to respective areas. We separate the geographical areas within the Atlantic Ocean (North and South) and lump all other EEZs into “other oceans.” This latter category is largely the Pacific Ocean and to a lesser extent the Indian Ocean. We remove the high seas catch from this database as it is not ocean-specific and we have extracted the ocean-specific high seas catch previously.

```
# remove the high seas catch from this database
d <- d[which(d$area_type != "high_seas"), ]
d$fishing_entity <- as.factor(d$fishing_entity)
d$fishing_entity <- droplevels(d$fishing_entity)
table(d$area_type) # this should be all "eez"
```

```
eez
63056
```

```
# Western Central Atlantic EEZs
WCA <- c("Bahamas", "Barbados", "Cayman Isl. (UK)", "Dominica", "Dominican Republic", "Grenada",
        "Aruba (Netherlands)", "Haiti", "Guadeloupe (France)", "Jamaica", "Martinique (France)",
        "Montserrat (UK)", "Nicaragua (Caribbean)", "Puerto Rico (USA)", "Saint Kitts & Nevis",
        "Saint Lucia", "Saint Vincent & the Grenadines", "Trinidad & Tobago", "US Virgin Islands",
        "St Barthelemy (France)", "St Martin (France)", "Curaçao (Netherlands)", "Bonaire",
        "Saba and Sint Eustatius (Netherlands)", "Sint Maarten (Netherlands)", "Honduras (Caribbean)",
        "Colombia (Caribbean)", "Mexico (Atlantic)", "Turks & Caicos Isl. (UK)", "Guyana",
        "Antigua & Barbuda", "Belize", "British Virgin Isl. (UK)", "French Guiana", "Cuba",
        "Anguilla (UK)", "Suriname", "Costa Rica (Caribbean)", "Guatemala (Caribbean)",
        "Panama (Caribbean)", "Bermuda (UK)", "Curaçao (Netherlands)")

# Brazil
Bra <- c("Brazil (mainland)", "Fernando de Noronha (Brazil)",
        "St Paul and St. Peter Archipelago (Brazil)")

# Canada
Can <- c("Canada (East Coast)", "Saint Pierre & Miquelon (France)")

# United States
USA <- c("USA (East Coast)", "USA (Gulf of Mexico)")
```

12 Analysis of international catches in territorial waters

```
# Northeast Atlantic EEZs
NEA <- c("Canary Isl. (Spain)", "Spain (mainland; Med and Gulf of Cadiz)", "Spain (Northwest)",
        "Spain (mainland, Med and Gulf of Cadiz)",
        "France (Atlantic Coast)", "Portugal (mainland)", "Azores Isl. (Portugal)")

# Eastern Central Atlantic EEZs
ECA <- c("Madeira Isl. (Portugal)", "Cape Verde", "Equatorial Guinea", "Benin",
        "Sao Tome & Principe ", "Sao Tome & Principe", "Morocco (Central)", "Ghana",
        "Côte d'Ivoire", "CÃ´te d'Ivoire", "Guinea", "Liberia", "Sierra Leone", "Nigeria",
        "Guinea-Bissau", "Senegal", "Cameroon", "Gambia", "Mauritania", "Morocco (South)",
        "Gabon", "Congo; R. of", "Congo, R. of", "Congo (ex-Zaire)")

# Southeastern Atlantic EEZs
SEA <- c("Angola", "Namibia", "South Africa (Atlantic and Cape)",
        "Ascension Isl. (UK)", "Saint Helena (UK)", "Tristan da Cunha Isl. (UK)")

# Southwest Atlantic EEZs
SWA <- c("Trindade & Martim Vaz Isl. (Brazil)", "Uruguay")

# Mediterranean EEZs
Med <- c("Italy (mainland)", "Libya", "Malta", "Syria", "Sicily (Italy)", "Sardinia (Italy)",
        "Balearic Islands (Spain)", "Cyprus (North)", "Cyprus (South)",
        "Gaza Strip", "Greece (without Crete)", "Crete (Greece)", "Lebanon",
        "Turkey (Mediterranean Sea)", "Egypt (Mediterranean)", "Israel (Mediterranean)",
        "Algeria", "Albania", "Croatia", "Montenegro", "Morocco (Mediterranean)",
        "France (Mediterranean)", "Corsica (France)")

# add variable to specify region
d$reg <- "other oceans"
d$reg[which(d$area_name %in% WCA)] <- "Western Central Atlantic EEZs"
d$reg[which(d$area_name %in% Bra)] <- "Brazil EEZ"
d$reg[which(d$area_name %in% Can)] <- "Canadian EEZ"
d$reg[which(d$area_name %in% USA)] <- "United States EEZ"
d$reg[which(d$area_name %in% NEA)] <- "Northeast Atlantic EEZs"
d$reg[which(d$area_name %in% ECA)] <- "Eastern Central Atlantic EEZs"
d$reg[which(d$area_name %in% SEA)] <- "Southeast Atlantic EEZs"
d$reg[which(d$area_name %in% SWA)] <- "Southwest Atlantic EEZs"
d$reg[which(d$area_name %in% Med)] <- "Mediterranean Sea EEZs"
```

Now we will check that we have classified the regions correctly.

12 Analysis of international catches in territorial waters

```
# check region designation
othlis <- unique(d$area_name[which(d$reg == "other oceans")])
print("EEZs of all other oceans: ")
```

```
[1] "EEZs of all other oceans: "
```

```
othlis
```

[1] "Australia"	"China"
[3] "Taiwan"	"Mayotte (France)"
[5] "French Polynesia"	"Hong Kong (China)"
[7] "Andaman & Nicobar Isl. (India)"	"Indonesia (Eastern)"
[9] "Japan (main islands)"	"Japan (Daito Islands)"
[11] "Korea (South)"	"Hawaii Northwest Islands (USA)"
[13] "Nauru"	"Niue (New Zealand)"
[15] "Northern Marianas (USA)"	"Micronesia (Federated States of)"
[17] "Marshall Isl."	"Palau"
[19] "Pakistan"	"Papua New Guinea"
[21] "Philippines"	"Seychelles"
[23] "Viet Nam"	"Tokelau (New Zealand)"
[25] "Tonga"	"Hawaii Main Islands (USA)"
[27] "USA (West Coast)"	"Wake Isl. (USA)"
[29] "Wallis & Futuna Isl. (France)"	"Samoa"
[31] "Guatemala (Pacific)"	"Kiribati (Gilbert Islands)"
[33] "Mexico (Pacific)"	"Japan (Ogasawara Islands)"
[35] "Korea (North, Yellow Sea)"	"Korea (North, Sea of Japan)"
[37] "Mauritius"	"Solomon Isl."
[39] "Christmas Isl. (Australia)"	"Johnston Atoll (USA)"
[41] "Timor Leste"	"Palmyra Atoll & Kingman Reef (USA)"
[43] "Jarvis Isl. (USA)"	"Howland & Baker Isl. (USA)"
[45] "Indonesia (Central)"	"Indonesia (Indian Ocean)"
[47] "Kiribati (Line Islands)"	"Kiribati (Phoenix Islands)"
[49] "New Caledonia (France)"	"Vanuatu"
[51] "Lord Howe Isl. (Australia)"	"Myanmar"
[53] "Ecuador (mainland)"	"El Salvador"
[55] "Fiji"	"Malaysia (Peninsula West)"
[57] "Norfolk Isl. (Australia)"	"Peru"
[59] "Pitcairn (UK)"	"Colombia (Pacific)"
[61] "Costa Rica (Pacific)"	"Nicaragua (Pacific)"

12 Analysis of international catches in territorial waters

[63] "Panama (Pacific)"	"Thailand (Andaman Sea)"
[65] "Thailand (Gulf of Thailand)"	"Cook Islands"
[67] "Kermadec Isl. (New Zealand)"	"Tuvalu"
[69] "Honduras (Pacific)"	"American Samoa"
[71] "Galapagos Isl. (Ecuador)"	"New Zealand"
[73] "Brunei Darussalam"	"Malaysia (Sabah)"
[75] "Malaysia (Sarawak)"	"Guam (USA)"
[77] "Clipperton Isl. (France)"	"South Africa (Indian Ocean Coast)"
[79] "Singapore"	"Easter Isl. (Chile)"
[81] "Madagascar"	"Réunion (France)"
[83] "Chagos Archipelago (UK)"	"Comoros Isl."
[85] "Mozambique Channel Isl. (France)"	"Kenya"
[87] "Mozambique"	"Somalia"
[89] "Tanzania"	"Glorieuse Islands (France)"
[91] "Yemen (Socotra)"	"Oman"
[93] "St Paul & Amsterdam Isl. (France)"	"Iran (Sea of Oman)"
[95] "Desventuradas Isl. (Chile)"	"India (mainland)"
[97] "Maldives"	"Tromelin Isl. (France)"
[99] "Cambodia"	"Sri Lanka"
[101] "Malaysia (Peninsula East)"	"Bangladesh"

```
print("U.S. EEZs outside the Atlantic Ocean: ")
```

```
[1] "U.S. EEZs outside the Atlantic Ocean: "
```

```
othlis[grep("USA", othlis)]
```

[1] "Hawaii Northwest Islands (USA)"	"Northern Marianas (USA)"
[3] "Hawaii Main Islands (USA)"	"USA (West Coast)"
[5] "Wake Isl. (USA)"	"Johnston Atoll (USA)"
[7] "Palmyra Atoll & Kingman Reef (USA)"	"Jarvis Isl. (USA)"
[9] "Howland & Baker Isl. (USA)"	"Guam (USA)"

```
print("French EEZs outside the Atlantic Ocean: ")
```

```
[1] "French EEZs outside the Atlantic Ocean: "
```

12 Analysis of international catches in territorial waters

```
othlis[grep("France", othlis)]
```

```
[1] "Mayotte (France)"           "Wallis & Futuna Isl. (France)"
[3] "New Caledonia (France)"     "Clipperton Isl. (France)"
[5] "Réunion (France)"          "Mozambique Channel Isl. (France)"
[7] "Glorieuse Islands (France)" "St Paul & Amsterdam Isl. (France)"
[9] "Tromelin Isl. (France)"
```

```
unique(d$area_name[which(d$reg == "Western Central Atlantic EEZs")]) # Western Central EE
```

```
[1] "Bahamas"
[2] "Barbados"
[3] "Bermuda (UK)"
[4] "Cayman Isl. (UK)"
[5] "Dominica"
[6] "Dominican Republic"
[7] "Grenada"
[8] "Guadeloupe (France)"
[9] "Haiti"
[10] "Jamaica"
[11] "Martinique (France)"
[12] "Montserrat (UK)"
[13] "Puerto Rico (USA)"
[14] "Saint Kitts & Nevis"
[15] "Saint Lucia"
[16] "Saint Vincent & the Grenadines"
[17] "Trinidad & Tobago"
[18] "US Virgin Isl."
[19] "St Barthelemy (France)"
[20] "St Martin (France)"
[21] "Curaçao (Netherlands)"
[22] "Bonaire (Netherlands)"
[23] "Saba and Sint Eustatius (Netherlands)"
[24] "Sint Maarten (Netherlands)"
[25] "Honduras (Caribbean)"
[26] "Colombia (Caribbean)"
[27] "Mexico (Atlantic)"
[28] "Turks & Caicos Isl. (UK)"
[29] "Costa Rica (Caribbean)"
```

12 Analysis of international catches in territorial waters

```
[30] "Panama (Caribbean)"
[31] "Antigua & Barbuda"
[32] "Belize"
[33] "British Virgin Isl. (UK)"
[34] "Cuba"
[35] "French Guiana"
[36] "Guyana"
[37] "Anguilla (UK)"
[38] "Suriname"
[39] "Venezuela"
[40] "Guatemala (Caribbean)"
[41] "Aruba (Netherlands)"
[42] "Nicaragua (Caribbean)"
```

```
unique(d$area_name[which(d$reg == "Brazil EEZ")]) # Brazil EEZs:
```

```
[1] "Brazil (mainland)"
[2] "Fernando de Noronha (Brazil)"
[3] "St Paul and St. Peter Archipelago (Brazil)"
```

```
unique(d$area_name[which(d$reg == "Canadian EEZ")]) # Canadian EEZs:
```

```
[1] "Canada (East Coast)" "Saint Pierre & Miquelon (France)"
```

```
unique(d$area_name[which(d$reg == "United States EEZ")]) # United States EEZs:
```

```
[1] "USA (East Coast)" "USA (Gulf of Mexico)"
```

```
unique(d$area_name[which(d$reg == "Northeast Atlantic EEZs")]) # Northeast Atlantic EEZs:
```

```
[1] "Azores Isl. (Portugal)"
[2] "Spain (mainland, Med and Gulf of Cadiz)"
[3] "Canary Isl. (Spain)"
[4] "Portugal (mainland)"
[5] "France (Atlantic Coast)"
[6] "Spain (Northwest)"
```

12 Analysis of international catches in territorial waters

```
unique(d$area_name[which(d$reg == "Eastern Central Atlantic EEZs")]) # Eastern Central At
```

```
[1] "Cape Verde"           "Equatorial Guinea"
[3] "Sao Tome & Principe" "Madeira Isl. (Portugal)"
[5] "Togo"                 "Ghana"
[7] "Guinea"               "Liberia"
[9] "Guinea-Bissau"        "Senegal"
[11] "Sierra Leone"        "Morocco (Central)"
[13] "Morocco (South)"      "Côte d'Ivoire"
[15] "Gabon"                 "Benin"
[17] "Gambia"                "Mauritania"
[19] "Nigeria"              "Cameroon"
[21] "Congo, R. of"          "Congo (ex-Zaire)"
```

```
unique(d$area_name[which(d$reg == "Southeast Atlantic EEZs")]) # Southeast Atlantic EEZs:
```

```
[1] "Ascension Isl. (UK)"      "South Africa (Atlantic and Cape)"
[3] "Saint Helena (UK)"       "Angola"
[5] "Namibia"                  "Tristan da Cunha Isl. (UK)"
```

```
unique(d$area_name[which(d$reg == "Southwest Atlantic EEZs")]) # Southwest Atlantic EEZs:
```

```
[1] "Trindade & Martim Vaz Isl. (Brazil)" "Uruguay"
```

```
unique(d$area_name[which(d$reg == "Mediterranean Sea EEZs")]) # Mediterranean Sea EEZs:
```

```
[1] "Italy (mainland)"      "Libya"
[3] "Malta"                 "Syria"
[5] "Sicily (Italy)"        "Sardinia (Italy)"
[7] "Balearic Islands (Spain)" "Tunisia"
[9] "Algeria"               "Morocco (Mediterranean)"
[11] "France (Mediterranean)" "Corsica (France)"
[13] "Croatia"               "Lebanon"
[15] "Albania"               "Greece (without Crete)"
[17] "Crete (Greece)"        "Cyprus (South)"
```

12.4 Summarize catch by regional EEZs

Now we summarize the catch by region by grouping the catch across the EEZs by region. We save these data for later analysis, and move on to analyzing only the Western Central and Northwest Atlantic jurisdictions.


```
# summarize EEZ catch by region
tab <- tapply(d$lbs, list(d$year, d$reg), sum, na.rm = T)
write.csv(data.frame(tab), file = "data/outputs/intl_catches_allAtlantic.csv", row.names =

dwest <- d[which(d$reg == "Canadian EEZ" | d$reg == "Western Central Atlantic EEZs"), ]
save(dwest, file = "data/outputs/SAU_EEZs_WCA.RData")

tab <- tapply(dwest$lbs, list(dwest$year, dwest$area_name), sum, na.rm = T)

# eight countries are responsible for vast majority of catch with total catches across all
par(mar = c(17, 4, 3, 1))
barplot(sort(colSums(tab, na.rm = T)/10^6, decreasing = T), horiz = F, las = 2,
        main = "Total dolphin catch (1950 - 2022) in the Western Atlantic, by EEZ",
        ylab = "total catch (millions of pounds)")
```

12 Analysis of international catches in territorial waters



Note that Bermuda is actually in the NCA region and not the Greater Caribbean region. However, its catches constitute 0.1% of the total catch which is essentially negligible, and to avoid complicating the operating model and code unnecessarily we will leave those catches accounted for in the CAR region.

```
round(colSums(tab, na.rm = T)[7] / sum(colSums(tab, na.rm = T), na.rm = T) * 100, 2)
```

Bermuda (UK)


12 Analysis of international catches in territorial waters

0.1

```
lis <- names(sort(colSums(tab, na.rm = T), decreasing = T)[1:8]) # list of top 8 jurisdictions
tab1 <- tab[, colnames(tab) %in% lis]

par(mar = c(3, 5, 3, 1))
matplot(rownames(tab1), tab1/10^6, las = 2, type = "l", lty = 1, lwd = 2, pch = 19, col = 1:8,
        xlab = "", ylab = "total catch (millions of pounds)", axes = F,
        main = "Total dolphinfish catches in Caribbean EEZs (top 8 jurisdictions)")
axis(1, at = seq(1950, 2020, 5)); axis(2, las = 2); box()
legend("topleft", colnames(tab1), col = 1:8, lty = 1, lwd = 3, bty = "n", title = "Fishing")
```

Total dolphinfish catches in Caribbean EEZs (top 8 jurisdictions)



Note the rapid eight-fold increase in catches listed within the Venezuelan EEZ which occurs from 2013 - 2019. This increase is responsible for a rapid increase in estimated international catches within the Western Atlantic.

12.5 Look at Venezuela catches in detail

The figures above indicate that the database shows a rapid increase in international catches within the Western Atlantic, driven by a sudden increase in catches reported

12 Analysis of international catches in territorial waters

within Sea Around Us database within the Venezuelan EEZ. We investigate these catches in detail, as it seems odd that a relatively small fleet from a single country would be able to ramp up effort so rapidly. Data can be downloaded on a country-specific basis, the catches by taxon in the waters of Venezuela are accessed here.

```
rm(list = ls()) # clear the workspace

# download the data for waters of Venezuela
ven <- read.csv("data/SAU/SAU EEZ 862 v50-1.csv")


# subset dolphinfish from the data
v <- ven[which(ven$common_name == "Common dolphinfish"), ]

#apply(v[1:13], 2, table)

#convert to pounds
v$pounds <- v$tonnes * 1000 * 2.20462

# look at the characteristics of the data
par(mar = c(5, 10, 1, 1), mfrow = c(3, 2), mex = 0.9)
#barplot(tapply(v$pounds/10^6, v$area_name, sum, na.rm = T), las = 2, xlab = "millions of p
#barplot(tapply(v$pounds/10^6, v$area_type, sum, na.rm = T), las = 2, xlab = "millions of p
barplot(tapply(v$pounds/10^6, v$year, sum, na.rm = T), las = 2, xlab = "millions of pounds"
barplot(tapply(v$pounds/10^6, v$fishing_entity, sum, na.rm = T), las = 2, xlab = "millions
barplot(tapply(v$pounds/10^6, v$fishing_sector, sum, na.rm = T), las = 2, xlab = "millions
barplot(tapply(v$pounds/10^6, v$catch_type, sum, na.rm = T), las = 2, xlab = "millions of p
barplot(tapply(v$pounds/10^6, v$reporting_status, sum, na.rm = T), las = 2, xlab = "million
barplot(tapply(v$pounds/10^6, v$gear_type, sum, na.rm = T), las = 2, xlab = "millions of po
```

12 Analysis of international catches in territorial waters




12 Analysis of international catches in territorial waters

Note that a large portion of the catch in the database is listed as coming from an "Unknown Fishing Country." We extract these catches and further investigate the characteristics.

```
# extract the Unknown Fishing Country catch
v1 <- v[which(v$fishing_entity == "Unknown Fishing Country"), ]
#apply(v1[1:15], 2, table)

# look at characteristics
par(mar = c(3, 10, 1, 5), mfrow = c(2, 2), mex = 0.6)
barplot(tapply(v1$pounds/10^6, v1$year, sum, na.rm = T), las = 2, xlab = "millions of pounds")
barplot(tapply(v1$pounds/10^6, v1$fishing_entity, sum, na.rm = T), las = 2, xlab = "millions of pounds")
barplot(tapply(v1$pounds/10^6, v1$reporting_status, sum, na.rm = T), las = 2, xlab = "millions of pounds")
barplot(tapply(v1$pounds/10^6, v1$gear_type, sum, na.rm = T), las = 2, xlab = "millions of pounds")
```




Here is another look at the catches within the venezuelan EEZ, separated by fishing entity.

```
# summarize Venezuela EEZ catches by fishing entity
tab <- tapply(v$pounds, list(v$year, v$fishing_entity), sum, na.rm = T)
head(tab)
```

12 Analysis of international catches in territorial waters

	Martinique (France)	Suriname	Trinidad & Tobago	Unknown Fishing Country
1974	7320.429	NA	NA	NA
1976	12814.641	NA	NA	NA
1985	NA	NA	NA	NA
1986	NA	NA	NA	NA
1987	NA	NA	NA	NA
1988	NA	NA	NA	NA
Venezuela				
1974	NA			
1976	NA			
1985	4699.563			
1986	2229.484			
1987	4585.610			
1988	39960.149			

```
matplot(rownames(tab), tab/10^6, las = 2, type = "b", lty = 1, lwd = 2, pch = 19, col = 1:5,
        xlab = "", ylab = "total catch (millions of pounds)", axes = F,
        main = "Total dolphinfish catches in the Venezuelan EEZ")
axis(1, at = seq(1950, 2020, 5)); axis(2, las = 2); box()
legend("topleft", colnames(tab), col = 1:5, pch = 19, lty = 1, lwd = 3, bty = "n", title =
text(2004, 6, cex = 0.8,
     "These Unknown Fishing Country\ncatches appear in the database as an\nindustrial fleet")
```



The very substantial increase in international Western Atlantic dolphin catches can be attributed solely to this Unknown Fishing Country catches which appear in the database beginning in 2013. The documentation of the catch reconstruction can be found in the Sea Around Us documentation files (see Mendoza et al. 2015). We spoke to Jeremy Mendoza and other experts who were involved in the reconstruction of domestic catches in Venezuela, and they noted that the “Assigned tuna RFMO catch” actually emerge from spatially reshuffled landing reports to ICCAT, according to the methods of Coulter et al. (2020). In this work, the Sea Around Us team developed a global spatial reallocation of landings of tunas and associated species that were reported to ICCAT. In short, this work uses the fraction of reported spatially disaggregated data to reallocate all data reported to ICCAT. This step is necessary because there are not many countries that report spatial data to ICCAT in the region, and this method may lead to a significant reallocation of landings from the Caribbean to the Venezuelan EEZ, where the local pole and line fishery is significant (J. Mendoza, personal communication). Thus, it is likely that landings from other EEZs, that were included in the reconstructed catch data layer, could have been reassigned to the Venezuelan EEZ.

Because there may be some overlap between the reconstructed data used in the national EEZ estimates, and the ICCAT reporting that was reassigned, there could potentially be a “double counting” of these catch estimates in the Sea Around Us database. We verified with Sea Around Us database managers that their method led to an error resulting in an overestimate of the catch, and they advised us to remove

these Unknown Fishing Country catches as they are difficult to trace back to the data source.

12.6 Additional error in Curacao catches


During the MSE review in August 2026, we uncovered a similar error in the database in the catch history for Curacao. Here also, we find catches attributed to a U.S. fleet operating within the EEZ. These catches seem unreasonable given no known commercial U.S. fleets operating in the small island nation and the overall magnitude of the catches given the very small EEZ.

```
rm(list = ls())
d <- read.csv("data/SAU/SAU Taxa 600006 v50-1.csv", header = T, sep = ",") # v. 50.1 - Jun
d$pounds <- d$tonnes * 1000 * 2.20462 # convert tonnes to pounds

v <- d[which(d$area_name == "Curaçao (Netherlands)"), ]


# look at the characteristics of the data
par(mar = c(5, 10, 1, 1), mfrow = c(3, 2), mex = 0.9)
barplot(tapply(v$pounds/10^6, v$area_name, sum, na.rm = T), las = 2, xlab = "millions of po
barplot(tapply(v$pounds/10^6, v$fishing_entity, sum, na.rm = T), las = 2, xlab = "millions
barplot(tapply(v$pounds/10^6, v$fishing_sector, sum, na.rm = T), las = 2, xlab = "millions
barplot(tapply(v$pounds/10^6, v$catch_type, sum, na.rm = T), las = 2, xlab = "millions of p
barplot(tapply(v$pounds/10^6, v$reporting_status, sum, na.rm = T), las = 2, xlab = "million
barplot(tapply(v$pounds/10^6, v$gear_type, sum, na.rm = T), las = 2, xlab = "millions of po
```

12 Analysis of international catches in territorial waters



```
# summarize Curacao EEZ catches by fishing entity
tab <- tapply(v$pounds, list(v$year, v$fishing_entity), sum, na.rm = T)

par(mfrow = c(1, 1), mar = c(4, 4, 2, 1))
matplot(rownames(tab), tab/10^6, las = 2, type = "b", lty = 1, lwd = 2, pch = 19, col = 1:5,
        xlab = "", ylab = "total catch (millions of pounds)", axes = F,
        main = "Total dolphinfish catches in Curacao's EEZ")
axis(1, at = seq(1950, 2020, 5)); axis(2, las = 2); box()
legend("topleft", colnames(tab), col = 1:5, pch = 19, lty = 1, lwd = 3, bty = "n", title =
```



Given the sharp increase appearing in the same year (2013), we suspect that these data also appear in the database due to the spatially reshuffled landing reports to ICCAT from Coulter et al. (2020). We remove these catches from subsequent analysis given the likely error.

12.7 References

Pauly D., Zeller D., Palomares M.L.D. (Editors), 2020. Sea Around Us Concepts, Design and Data (searoundus.org).

13 Compilation of international catches from FAO

The Food and Agriculture Organization (FAO) is a specialized agency of the United Nations that monitors global agriculture, forestry, and fisheries. Its fisheries database provides one of the most comprehensive long-term records of international wild-capture fishing, drawing directly from official national reporting and standardized estimation models. Unlike the Sea Around Us (SAU) database, the FAO database contains only data officially declared by member countries and does not apply any catch algorithms to fill in missing data. However, as a transparent and standardized database it offers a useful comparison to the SAU database.

Annual catch data for common dolphinfish (*Coryphaena hippurus*) were pulled from the FAO Global Capture Production database (FAO, 2024). We filtered the dataset for FAO Fishing Areas 21 (Northwest Atlantic) and 31 (Western Central Atlantic) using the 3-alpha species code DOL. The code for the data pull is in the `FAO_access_data.R` file; the data are saved and accessed locally here.

13.1 Download FAO data and summarize

```
rm(list = ls()) # clear the workspace

# libraries
library(pals)

df <- read.csv("data/FAO_dolphin_data.csv") # open the FAO data file
head(df)
```

	JURISDICTION	AREA	SCIENTIFIC_NAME	YEAR
1	Belize	21	<i>Coryphaena hippurus</i>	2022
2	Canada	21	<i>Coryphaena hippurus</i>	2022

13 Compilation of international catches from FAO

3	Spain	21	Coryphaena hippurus	2022
4	United States of America	21	Coryphaena hippurus	2022
5	Venezuela (Bolivarian Republic of)	21	Coryphaena hippurus	2022
6	Antigua and Barbuda	31	Coryphaena hippurus	2022

	Landings_Tonnes
1	20.051
2	2.545
3	0.439
4	25.000
5	0.000
6	25.000

```
apply(df[, 1:4], 2, table)
```

\$JURISDICTION

Antigua and Barbuda	Barbados
22	73
Belize	Bermuda
84	73
British Virgin Islands	Canada
29	73
Costa Rica	Cuba
73	52
Dominica	Dominican Republic
21	55
France	Grenada
72	59
Guadeloupe	Martinique
73	73
Mexico	Nicaragua
59	1
Other NEI	Portugal
70	1
Puerto Rico	Saint Kitts and Nevis
24	28
Saint Lucia	Saint Vincent and the Grenadines
29	73
Spain	Trinidad and Tobago
76	11

13 Compilation of international catches from FAO

```

United States of America
146
Venezuela (Bolivarian Republic of)
39

United States Virgin Islands
20

$AREA

21 31
242 1167

$SCIENTIFIC_NAME

Coryphaena hippurus
1409

$YEAR

1950 1951 1952 1953 1954 1955 1956 1957 1958 1959 1960 1961 1962 1963 1964 1965
13 13 13 13 13 13 13 13 13 13 13 13 13 13 15 15
1966 1967 1968 1969 1970 1971 1972 1973 1974 1975 1976 1977 1978 1979 1980 1981
15 16 16 16 17 17 17 17 17 17 17 17 17 17 17 17
1982 1983 1984 1985 1986 1987 1988 1989 1990 1991 1992 1993 1994 1995 1996 1997
17 17 17 17 17 17 17 17 18 18 18 18 20 21 20 20
1998 1999 2000 2001 2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013
19 22 22 22 23 25 25 25 24 25 25 25 25 25 28 28
2014 2015 2016 2017 2018 2019 2020 2021 2022
28 28 28 28 28 28 28 29 28

```

```
df$lbs <- df$Landings_Tonnes * 2204.62
```

We first look at the major countries reporting dolphin catches in the FAO database, for the historical period and the recent period. Note that only commercial catches are included for the United States.

```



df_recent <- df[which(df$YEAR >= 2015), ]

par(mar = c(15, 4, 1, 1), mfrow = c(2, 1), cex = 0.9)
barplot(sort(tapply(df$lbs/10^6, df$JURISDICTION, mean, na.rm = T)), las = 2, horiz = F, ce
      ylab = "millions of pounds", main = "average annual landings (FAO 1950 - 2022)")
barplot(sort(tapply(df_recent$lbs/10^6, df_recent$JURISDICTION, mean, na.rm = T)), las = 2,
      ylab = "millions of pounds", main = "average annual recent landings (FAO 2015 - 2022)")

```

13 Compilation of international catches from FAO

13 Compilation of international catches from FAO



13.2 Compare FAO to SAU database

Now we will compare the international catches from FAO with the SAU catches compiled in the previous section.

```
df_int <- df[which(df$JURISDICTION != "United States of America"), ]
lis <- names(sort(tapply(df_int$lbs, df_int$JURISDICTION, mean, na.rm = T), decreasing = T))
df_int$JURISDICTION <- factor(df_int$JURISDICTION, levels = lis)

fao <- tapply(df_int$lbs, list(df_int$JURISDICTION, df_int$YEAR), sum, na.rm = T)
fao[is.na(fao)] <- 0

# plot FAO vs SAU
b <- barplot(fao/10^6, names.arg = 1950:2022, col = glasbey(26), border = NA, las = 2, ylim = c(0, 100),
             ylab = "millions of pounds", main = "FAO reported landings by jurisdiction")
legend("topleft", ncol = 2, rownames(fao), col = glasbey(26), pch = 15, bty = "n", title = "FAO landings")
abline(h = 0)

# load data
load("data/outputs/SAU_EEZs_WCA.RData") # SAU EEZ data for NED and NCA saved from earlier
ds <- dwest

# remove bad catches from the database prior to summarizing
#dim(ds)
ds <- ds[-which(ds$area_name == "Venezuela" & ds$fishing_entity == "Unknown Fishing Country"), ]
ds <- ds[-which(ds$area_name == "Curaçao (Netherlands)" & ds$fishing_entity == "USA"), ]
#dim(ds)

tot <- tapply(ds$lbs, ds$year, sum, na.rm = T) # total catch from SAU
xs <- b[which(1950:2022 <= 2019)]


tab_ds <- tapply(ds$lbs, list(ds$year, ds$catch_type, ds$reporting_status), sum, na.rm = T)
#tail(tab_ds)
repSAU <- tab_ds[, 2, 1] # reported catch from SAU
#unrepSAU <- tab_ds[, 2, 2] # repSAU + unrepSAU is total - discards (ve

lines(xs, tot/10^6, lty = 3, col = "#00000070", lwd = 3)
lines(xs, repSAU/10^6, lty = 1, col = 1, lwd = 3)

legend("left", c("reported + unreported", "reported"), title = "SAU landings",
```

13 Compilation of international catches from FAO

```
lwd = 3, lty = c(3, 1), col = c("#00000070", 1), bty = "n")
```



We see that the FAO landings track quite closely with respect to the overall trends in the SAU reported landings, which is what we would expect since the SAU reported landings do not include any reconstructions. In general, the FAO landings are higher than the SAU reported landings, although in some years the SAU reported landings are higher. The total SAU landings (including reconstructions of unreported data) are generally much higher than the FAO estimates, though in some years very high landings in the early 2010s the estimates from FAO exceed SAU. Of note, there is a very steep decline in FAO catches in the last three years of the database (2020 - 2022), driven primarily by

13 Compilation of international catches from FAO

a decline in reported catches from Venezuela. As the SAU database only goes up to 2019, we lack the ability to compare these years.

We compare the SAU and FAO estimates at the individual jurisdictional level to better understand the differences between databases.


```
sau <- tapply(ds$lbs, list(ds$year, ds$reporting_status, ds$area_name), sum, na.rm = T)
sau[is.na(sau)] <- 0

lk <- read.csv("data/FAO_SAU_lookup.csv", fileEncoding = "Windows-1252")

par(mfrow = c(9, 5), mar = c(1, 2.5, 1.5, 0.5), mgp = c(1.5, 0.5, 0))
for (i in 1:dim(sau)[3]) {
  # for each jurisdiction
  b <- barplot(t(sau[, , i])/10^6, col = c(3, 2), # plot reported and unreported SAU data
    names.arg = rep(NA, length(1950:2019)),
    beside = F, border = NA, space = 0,
    xlab = "", ylab = "", ylim = c(0, 4), axes = F)
  repunrep <- round(rowMeans(t(sau[, , i])/10^6, na.rm = T), 2)
  axis(1, at = b[seq(10, 70, 20)], lab = c("60", "80", "00", "20"), tick = F, pos = 0.2)
  axis(2, las = 2, tcl = -0.2, at = 0:4)
  maintext <- dimnames(sau)[[3]][i] # shorten main labels
  maintext <- gsub("\\band\\b", "&", maintext, ignore.case = TRUE)
  maintext <- gsub("\\b(Saint|St\\.\\.\\.\\b", "St", maintext, ignore.case = TRUE)
  mtext(side = 3, maintext, cex = 0.75)

  mtch <- lk$FAO[which(lk$SAU == dimnames(sau)[[3]][i])] # match to FAO name with lookup
  if (mtch != "") {
    f <- fao[which(rownames(fao) == mtch), 1:70]
    lines(b, f/10^6, col = 1)
    repunrep[3] <- round(mean(f, na.rm = T) / 10^6, 2)
  }
  lpos <- "topleft"; if (i == 5) { lpos = "topright"}
  legend(lpos, as.character(repunrep)[3:1], text.col = c(1, 2, 3), bty = "n")
}
plot.new()
legend("center", c("FAO", "SAU unreported", "SAU reported"),
  col = 1:3, lty = c(1, 0, 0), pch = c(NA, 15, 15))
```

13 Compilation of international catches from FAO



The figure shows a stacked barplot of the reported (in green) and unreported (in red) landings from the SAU database, with the FAO landings overlaid (black line). The

13 Compilation of international catches from FAO

numbers in the upper left corner are the average landings over the entire time series (in millions of pounds per year; green = reported SAU, red = unreported SAU, black = FAO); panels with missing numbers indicate that no data were available.

The comparison of FAO versus SAU databases reveals notable differences and some similarities. In a few cases, the SAU reported estimates line up closely or exactly with the FAO estimates for parts of the time series (Guadeloupe, Martinique, Saint Lucia, Venezuela). The majority of the jurisdictions have very small amounts of landings in the SAU database and no FAO reporting. Bahamas has substantial reconstructed catches and Haiti has a high number of “reported” catches in the SAU database, but neither of these countries have any reporting in the FAO database. In some cases, the FAO database contains significant landings that do not appear in the SAU database (Barbados, Cuba, French Guiana, and more recent years in Guadeloupe and Venezuela.)

13.3 Create a new maximum landings scenario dataset

Given the discrepancies revealed by the jurisdiction-level comparison, it seems likely that both SAU and FAO could potentially be missing key aspects of the international fishing. A maximum estimate of total international landings could be compiled by summarizing the maximum reported value of landings for each year, within each jurisdiction. We calculate these maxima at this resolution and recompile the data. At the time of analysis, the SAU database only contained catches to 2019, so we interpolated the 2019 values to cover years 2020 - 2022, prior to calculating the maxima.

```
SAU <- tapply(ds$lbs, list(ds$year, ds$area_name), sum, na.rm = T) # summary of SAU
maxland <- SAU
# dim(maxland)
maxland[is.na(maxland)] <- 0 # placeholder for maximum landings


# add 2020 - 2022 to SAU data (assume same as 2019 levels)
maxland <- rbind(maxland, maxland[nrow(maxland),], maxland[nrow(maxland),], maxland[nrow(maxland),])

for (i in 1:ncol(maxland)) { # for each jurisdiction in SAU
  mtch <- lk$FAO[which(lk$SAU == dimnames(maxland)[[2]][i])] # match names using lookup table
  if (mtch != "") { # if there is a match
    f <- fao[which(rownames(fao) == mtch), ] # get the FAO data
    temp <- cbind(f, maxland[, i])
    maxl <- apply(temp, 1, max, na.rm = TRUE) # calculate the maximum value
    maxland[, i] <- maxl # replace data with the maximum
```

13 *Compilation of international catches from FAO*

```
}  
}  
# now plot the maximum landings time series by jurisdiction  
par(mfrow = c(9, 5), mar = c(1, 2.5, 1.5, 0.5), mgp = c(1.5, 0.5, 0))  
for (i in 1:dim(maxland)[2]) {  
  plot(1950:2022, maxland[, i]/10^6, type = "l",  
       xlab = "", ylab = "", ylim = c(0, 4), axes = F)  
  axis(1, at = seq(1960, 2020, 20), lab = c("60", "80", "00", "20"),  
       tick = F, pos = 0.2)  
  axis(2, las = 2, tcl = -0.2, at = 0:4)  
  maintext <- dimnames(sau)[[3]][i]  
  maintext <- gsub("\\band\\b", "&", maintext, ignore.case = TRUE)  
  maintext <- gsub("\\b(Saint|St\\.\\.\\.\\b", "St", maintext, ignore.case = TRUE)  
  mtext(side = 3, maintext, cex = 0.75)  
}
```

13 Compilation of international catches from FAO





The landings time series plotted here now represent the maximum reported value in either data base.

We look at which jurisdictions are contributing to the landings, for the entire time series and more recent years, based on this new maximum assumption data set.

```
par(mar = c(15, 4, 1, 1), mfrow = c(2, 1), mgp = c(2.5, 0.75, 0), cex = 0.9)
barplot(sort(colMeans(maxland, na.rm = T)/10^6), las = 2, horiz = F, cex.names = 1,
        ylab = "millions of pounds", main = "average annual landings (maximum of FAO or SAU)
barplot(sort(colMeans(tail(maxland, 8), na.rm = T)/10^6), las = 2, horiz = F,
        ylab = "millions of pounds", main = "average annual recent landings (maximum of FAO
```

13 Compilation of international catches from FAO



13 *Compilation of international catches from FAO*


The top contributors to landings overall, according to this new combined data set, are: Barbados (substantial landings from FAO but limited reported landings from SAU), Bahamas (no FAO data, with mostly unreported reconstructed SAU landings), and Guadeloupe (exactly the same landings reported in SAU and FAO from 1950 - 2007 with significant divergence in recent years).

13.4 Compare landings scenarios

We compare the three different time series: FAO landings, SAU landings, and maximum landings, across the entire time series.

```
par(mar = c(4, 4, 2, 1), mgp = c(2.2, 1, 0))
plot(1950:2022, rowSums(maxland, na.rm = T)/10^6, type = "l", ylim = c(0, 22), las = 1, lwd = 3,
     xlab = "year", ylab = "millions of pounds", col = 3,
     main = "Total estimated international dolphin landings from different data sources")
lines(1950:2019, rowSums(SAU, na.rm = T)/10^6, col = 2, lty = 1, lwd = 2)
lines(1950:2022, colSums(fao, na.rm = T)/10^6, col = 4, lty = 1, lwd = 2)
text(2022, round(sum(maxland[nrow(maxland), ])/10^6, 1), round(sum(maxland[nrow(maxland), ])/10^6, 1),
text(2019, round(sum(SAU[nrow(SAU), ])/10^6, 1), round(sum(SAU[nrow(SAU), ])/10^6, 1), na.rm = T)
text(2022, round(sum(fao[, ncol(fao)])/10^6, 1), round(sum(fao[, ncol(fao)])/10^6, 1), na.rm = T)
legend("topleft", c("maximum of either SAU or FAO", "SAU (Sea Around Us)", "FAO (Food and Agriculture Organization)"),
     col = c(3, 2, 4), lty = 1, lwd = 3)
```

13 Compilation of international catches from FAO




The three different time series are similar in magnitude and trend during the first several decades, and grow increasingly divergent in time. The numbers on the plot show the total landings from each time series in the terminal year of data.

We take a closer look at the years of data used in the dolphin operating model.

```
par(mar = c(4, 4, 2, 1), mgp = c(2.2, 1, 0))
plot(1986:2022, rowSums(tail(maxland, 37), na.rm = T)/10^6, type = "l", ylim = c(0, 20), las = 1,
     xlab = "year", ylab = "millions of pounds", col = 3,
     main = "Total estimated international dolphin landings from different data sources")
lines(1986:2019, rowSums(tail(SAU, 34), na.rm = T)/10^6, col = 2, lty = 1, lwd = 2)
lines(1986:2022, colSums(fao[, 37:73], na.rm = T)/10^6, col = 4, lty = 1, lwd = 2)
abline(h = mean(rowSums(tail(maxland, 37), na.rm = T)/10^6), col = "#00FF0030", lwd = 4, lty = 3)
abline(h = mean(rowSums(tail(SAU, 34), na.rm = T)/10^6), col = "#FF00FF30", lwd = 4, lty = 3)
abline(h = mean(colSums(fao[, 37:73], na.rm = T)/10^6), col = "#00FFFF30", lwd = 4, lty = 3)

legend("bottomright", c("maximum of either SAU or FAO", "SAU (Sea Around Us)",
                        "FAO (Food and Agriculture Organization)", "average over time series"),
      col = c(3, 2, 4, "#00000030"), lty = c(1, 1, 1, 3), lwd = c(2, 2, 2, 3), bty = "n")
```

13 Compilation of international catches from FAO



The more recent data show that assumptions regarding levels of international catches vary quite widely depending on the data source. The maximum landings data suggest that catches have increased significantly from 1986 to 2011, followed by a substantial decrease with the most recent landings just below the average of the time series. The SAU data suggest an increase from 1986 to the late 1990s, with relatively stable landings over the last 20 years. Finally, the FAO data suggests an increase in the early years with a large peak in 2011, followed by a significant decline, with the most recent landings falling back to the late 1980s levels.

13.5 Output scenarios

For the management strategy evaluation, we build into the operating model multiple international landings scenarios to account for the wide uncertainty in international removals. The FAO time series (blue) versus the maximum landings scenario (green) provides good contrast in terms of the magnitude of removals as well as recent trends. We acknowledge that the FAO landings, which only include reported data, are likely an underestimate of true landings as there are likely to be unreported landings in this region. However, this scenario provides a more robust test of the management

13 Compilation of international catches from FAO

procedures, because lower international landings will result in changes to domestic U.S. fisheries having a greater influence on the stock.

```
fao1 <- t(fao)

#dim(fao1)
#dim(maxland)

# remove Canada and St. Pierre and Miquelon as these are already accounted for in the NED
fao1 <- fao1[, -grep("Canada", colnames(fao1))]
maxland <- maxland[, -grep("Canada", colnames(maxland))]
maxland <- maxland[, -grep("Pierre", colnames(maxland))]

hi <- rowSums(maxland, na.rm = T)
lo <- rowSums(fao1, na.rm = T)

fin <- data.frame(cbind(1950:2022, lo, hi))
names(fin)[1] <- "year"
head(fin)
```

	year	lo	hi
1950	1950	2204620	2888241
1951	1951	2204620	3011134
1952	1952	2866006	3714998
1953	1953	2866006	3765129
1954	1954	2425082	3392490
1955	1955	2425082	3453073

```
# output high and low scenarios for further compilation
write.csv(fin, file = "data/hi_lo_CAR_landings.csv", row.names = FALSE)
```

13.6 References

FAO. (2024). Fisheries and Aquaculture Division: Global Capture Production 1950–2022 (Release 2024.1.0) [Data set]. Food and Agriculture Organization of the United Nations. <https://www.fao.org/fishery/collection/capture/en>

14 Compilation of international catches

14.1 Compile catch in CAR and NED jurisdictional waters by discards and reporting type

Now that we have separated the EEZ catch by region, we will summarize it according to the areas used in the dolphin management strategy evaluation operating model. For the Caribbean region, we have the high and low international catch scenarios that we compile for the sensitivity analyses in the operating model.

```
rm(list = ls()) # clear the workspace

# load data
load("data/outputs/SAU_EEZs_WCA.RData") # SAU EEZ data for NED and NCA saved from earlier

# summarize the catch by area and separate out Canadian landings
tapply(dwest$lbs, dwest$reg, sum, na.rm = T)
```

Canadian EEZ	Western Central Atlantic EEZs
1220146	571673711

```
dwest$year <- factor(dwest$year, levels = 1950:2019)
dc <- dwest[which(dwest$reg == "Canadian EEZ"), ]

CAN <- tapply(dc$lbs, dc$year, sum, na.rm = T)
#tail(CAN)

# load Caribbean international landings scenarios
car_sc <- read.csv("data/hi_lo_CAR_landings.csv")
car_sc <- car_sc[which(car_sc$year >= 1986), ] # only need data 1986 and onwards
head(car_sc)
```

14 Compilation of international catches

	year	lo	hi
37	1986	4508448	8496677
38	1987	3836039	7246145
39	1988	6671180	10331010
40	1989	4091775	7526003
41	1990	4894256	8891365
42	1991	4894256	8971473

```
# combine CAR reported landings, unreported discards, unreported landings
CARlo <- car_sc$lo
CARhi <- car_sc$hi
```

14.2 Distribute the territorial international catches across quarters of the year

The international catches are only reported as annual catches and do not have any month or season associated with them. However, the MSE operating model requires seasonal catches. Lacking other information on the seasonality of these fleets, we will assume that the fleet distribution of the U.S. pelagic longline is representative of the seasonality of the jurisdictional fleets. Now we parse the annual catches according to the seasonality of the U.S. fleet.

The pelagic longline logbook data is only available going back to 1997, but there are Caribbean catches extending back to the 1950s. To interpolate the seasonality for 1986 - 1996, we take the average seasonality of the oldest 10 years of logbook data (1997 - 2006) and use those averages for the unknown years.

```
# read in the percentage of the catch by area and year-quarter combinations from the U.S. P
load("data/outputs/per_PLLcatch_by_area_yearquarter.RData")
```

```
names(percatch[1, 1, ]) # matrix 2 == Caribbean, matrix 7 == NED (Atlantic Northwest)
```

```
[1] "NCA" "CAR" "FLK" "NCFL" "NNC" "VBM" "NED"
```

```
# parse Canada data into quarters
```

```
CAN <- CAN[which(rownames(CAN) >= 1997)] # PLL data are only available from 1997 on
CAN <- c(CAN, CAN[length(CAN)], CAN[length(CAN)], CAN[length(CAN)]) # interpolate 2019 cat
```

14 Compilation of international catches

```
CANqrt <- percatch[, , 7] # create matrix for final catch data
for (i in 1:ncol(percatch)) {
  CANqrt[, i] <- percatch[, i, 7] * CAN[i] }

percatch_car <- percatch[, , 2] # create matrix for Caribbean percentages
int <- rowMeans(percatch_car[, 1:10]) # calculate the average seasonality using the oldest
percatch_car <- cbind(matrix(rep(int, length(1986:1996)), nrow = 4), percatch_car) # attach
colnames(percatch_car) <- 1986:2022 # full data frame for Caribbean seasonality 1986 - 2022

length(CARlo)
```

```
[1] 37
```

```
dim(percatch_car)
```

```
[1] 4 37
```

```
CARlo_qrt <- percatch_car # create matrices for final catch data
CARhi_qrt <- percatch_car

for (i in 1:ncol(percatch_car)) {
  CARlo_qrt[, i] <- percatch_car[, i] * CARlo[i]
  CARhi_qrt[, i] <- percatch_car[, i] * CARhi[i]
}

# ensure sums are equal after parsing
sum(CARlo_qrt); sum(CARlo)
```

```
[1] 294505847
```

```
[1] 294505847
```

```
sum(CARhi_qrt); sum(CARhi)
```

```
[1] 513883150
```

```
[1] 513883150
```

14 Compilation of international catches

```
sum(CAN, na.rm = T); sum(CANqrt, na.rm = T)
```

```
[1] 1328482
```

```
[1] 1328482
```

The Canada jurisdictional data need to be combined with the NED high seas data calculated in the previous section, because both of these are a component of the international catches in the NED area.

```
load("data/outputs/NED_year_quarter.RData")
```

```
NEDtot <- NED + CANqrt
```

```
# ensure sums are correct
```

```
sum(NED, na.rm = T) + sum(CANqrt, na.rm = T); sum(NEDtot, na.rm = T)
```

```
[1] 1808855
```

```
[1] 1808855
```

```
NEDtot <- NEDtot[, !is.na(colSums(NEDtot))] # remove NA columns from CAN data
```

```
# plot the parsed out data by year-quarter
```


```
yrs <- sort(rep(as.numeric(colnames(CARlo_qrt)), 4))
```

```
qrt <- rep(1:4, length(yrs)/4)
```

```
yrs1 <- sort(rep(as.numeric(colnames(NEDtot)), 4))
```

```
qrt1 <- rep(1:4, length(yrs1)/4)
```

```
matplot(yrs + qrt/4, cbind(matrix(CARhi_qrt), matrix(CARlo_qrt))/10^6, lty = 1, lwd = 2,  
      col = 2:3, type = "l", main = "International jurisdictional catches by quarter",  
      xlab = "", ylab = "total catch (millions of pounds)")  
lines(yrs1 + qrt1/4, as.numeric(matrix(NEDtot))/10^6, lwd = 2, col = 4)  
legend("topleft", c("Caribbean landings - high scenario",  
      "Caribbean landings - low scenario", "NED reported landings"),  
      lwd = 2, col = 2:4, bty = "n")
```



Finally, we format the data in the format for input into the operating model.

```
# format for operating model - Caribbean
flt <- rep("Intl", length(yrs))
are <- rep("CAR", length(yrs))

# high and low landings scenarios
finhi <- data.frame(cbind(yrs, qrt, flt, are, matrix(CARhi_qrt)))
finlo <- data.frame(cbind(yrs, qrt, flt, are, matrix(CARlo_qrt)))

# format final matrices
names(finhi) <- c("Year", "Quarter", "Fleet", "Area", "Catch_lbs")
finhi$Catch_lbs <- as.numeric(finhi$Catch_lbs)
head(finhi)
```

	Year	Quarter	Fleet	Area	Catch_lbs
1	1986	1	Intl	CAR	4595149.4
2	1986	2	Intl	CAR	3158470.6
3	1986	3	Intl	CAR	500827.7
4	1986	4	Intl	CAR	242229.6
5	1987	1	Intl	CAR	3918839.7

14 Compilation of international catches

```
6 1987      2 Intl CAR 2693610.0
```

```
names(finlo) <- c("Year", "Quarter", "Fleet", "Area", "Catch_lbs")
finlo$Catch_lbs <- as.numeric(finlo$Catch_lbs)
head(finlo)
```

	Year	Quarter	Fleet	Area	Catch_lbs
1	1986	1	Intl	CAR	2438246.3
2	1986	2	Intl	CAR	1675925.7
3	1986	3	Intl	CAR	265745.7
4	1986	4	Intl	CAR	128530.2
5	1987	1	Intl	CAR	2074595.8
6	1987	2	Intl	CAR	1425971.1

```
#par(mfrow = c(2, 2))
#plot(finhi$Catch_lbs, type = "l")
#plot(finhi$Catch_lbs ~ factor(finhi$Fleet))
#plot(finhi$Catch_lbs ~ factor(finhi$Quarter))
#plot(finhi$Catch_lbs ~ factor(finhi$Year))
#apply(finhi, 2, table)

# write the output files for both scenarios
write.csv(finhi, file = "data/FINAL_files/CAR_Intl_hi_TomFormat.csv", row.names = FALSE)
write.csv(finlo, file = "data/FINAL_files/CAR_Intl_lo_TomFormat.csv", row.names = FALSE)

# format for operating model - NED
flt <- rep("Intl", length(yrs1))
are <- rep("NED", length(yrs1))
fin <- data.frame(cbind(yrs1, qrt1, flt, are, matrix(NEDtot)), stringsAsFactors = FALSE)
names(fin) <- c("Year", "Quarter", "Fleet", "Area", "Catch_lbs")
fin$Catch_lbs <- as.numeric(fin$Catch_lbs)
head(fin)
```

	Year	Quarter	Fleet	Area	Catch_lbs
1	2007	1	Intl	NED	0.000
2	2007	2	Intl	NED	1450.366
3	2007	3	Intl	NED	13279.912
4	2007	4	Intl	NED	0.000
5	2008	1	Intl	NED	0.000
6	2008	2	Intl	NED	1312.750

14 *Compilation of international catches*

```
#apply(fin, 2, table)

# write output file
write.csv(fin, file = "data/FINAL_files/NED_Intl_TomFormat.csv", row.names = FALSE)

#barplot(tapply(fin$Catch_lbs, list(fin$Quarter), sum, na.rm = T), col = rainbow(4))
```

15 Data compilation for all fleets in the operating model

In this last step we will compile all of the output data files from the different data sources (NMFS commercial trip tickets, NMFS recreational statistics, Sea Around Us) and check the files for accuracy.

Recall that there are seven defined areas in the operating model, over which the landings are summed. Four of these areas are within the U.S. EEZ and the other three areas are international waters (high seas and jurisdictional waters of other countries).

** Later due to computational issues in resolving the very fine spatial scales, we collapsed the seven areas into five areas, combining CAR + FLK into one area and NNC + VBM into another area.

15 Data compilation for all fleets in the operating model




Figure 15.1: Areas defined in the operating model.

15.1 Data upload

First we access all of the final data files from the previous steps.

```
# clear workspace
rm(list = ls())

# view data files in directory
dir("data/FINAL_files/")
```

```
[1] "CAR_Intl_hi_TomFormat.csv"
[2] "CAR_Intl_lo_TomFormat.csv"
[3] "commercial_TomFormat.csv"
[4] "complete_catch_dataset_100526_high.csv"
[5] "complete_catch_dataset_100526_low.csv"
[6] "NCA_Intl_TomFormat.csv"
[7] "NED_Intl_TomFormat.csv"
[8] "rec_catch_TomFormat.csv"
```

```
scenario <- "low"

if (scenario == "high") { intc <- read.csv("data/FINAL_files/CAR_Intl_hi_TomFormat.csv") }
if (scenario == "low") { intc <- read.csv("data/FINAL_files/CAR_Intl_lo_TomFormat.csv") }

intn <- read.csv("data/FINAL_files/NCA_Intl_TomFormat.csv")
intw <- read.csv("data/FINAL_files/NED_Intl_TomFormat.csv")
com <- read.csv("data/FINAL_files/commercial_TomFormat.csv")
rec <- read.csv("data/FINAL_files/rec_catch_TomFormat.csv")
```

The "CAR_Intl_hi_TomFormat.csv" file contains all international landings and unreported catches for the greater Caribbean region (jurisdictional waters of all countries). This includes unreported data and discards, which were very small (< 1%). Recall that we consider two scenarios: a high and a low international removals scenario. For the visualization here we use the high catch scenario.

```
apply(intc[1:4], 2, table, useNA = "always")
```

\$Year

15 Data compilation for all fleets in the operating model

1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001
4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4
2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017
4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4
2018	2019	2020	2021	2022	<NA>										
4	4	4	4	4	0										

\$Quarter

1	2	3	4	<NA>
37	37	37	37	0

\$Fleet

Intl	<NA>
148	0

\$Area

CAR	<NA>
148	0

```
intc$Area <- "FLKCAR"
```

The "NCA_Intl_TomFormat.csv" file contains all international landings for the NCA region (i.e., FAO region 31), also known as the Western Central Atlantic. There are no territorial waters in this area (except for Bermuda, which as discussed previously is summarized with the CAR region) and there are no unreported catches or discards in the database.

```
apply(intn[1:4], 2, table, useNA = "always")
```

\$Year

1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4
2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	<NA>					
4	4	4	4	4	4	4	4	4	4	0					

15 Data compilation for all fleets in the operating model

\$Quarter

1	2	3	4	<NA>
26	26	26	26	0

\$Fleet

Intl	<NA>
104	0

\$Area

NCA	<NA>
104	0

The "NED_Intl_TomFormat.csv" file contains all international landings for the NED region (i.e., FAO region 21), also known as the Northwest Atlantic. This includes NED high seas landings, as well as landings from Canadian and French territorial waters which fall in this region. There are no unreported catches or discards for this region in the database.

```
apply(intw[1:4], 2, table, useNA = "always")
```

\$Year

2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4
<NA>															
0															

\$Quarter

1	2	3	4	<NA>
16	16	16	16	0

\$Fleet

Intl	<NA>
64	0

15 Data compilation for all fleets in the operating model

\$Area

```
NED <NA>
64      0
```

The “commercial_TomFormat.csv” file contains all landings for the U.S. commercial fleet, including landings from U.S. EEZs and high seas. There are no unreported catches in this region. Recall that estimated dead discards were very small (< 2%) and were removed from the summary.

```
apply(com[1:4], 2, table, useNA = "always")
```

\$Year

```
1986 1987 1988 1989 1990 1991 1992 1993 1994 1995 1996 1997 1998 1999 2000 2001
  28   28   28   28   28   28   28   28   28   28   28   28   28   28   28   28
2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017
  28   28   28   28   28   28   28   28   28   28   28   28   28   28   28   28
2018 2019 2020 2021 2022 2023 2024 <NA>
  28   28   28   28   28   28    7    0
```

\$Quarter

```
  1    2    3    4 <NA>
273 266 266 266    0
```

\$Fleet

```
UScom <NA>
1071    0
```

\$Area

```
CAR  FLK  NCA NCFL  NED  NNC  VBM <NA>
153  153  153  153  153  153  153    0
```

```
com$Area[which(com$Area == "CAR")] <- "FLKCAR"
com$Area[which(com$Area == "FLK")] <- "FLKCAR"
com$Area[which(com$Area == "NNC")] <- "NNCVBM"
```

15 Data compilation for all fleets in the operating model

```
com$Area[which(com$Area == "VBM")] <- "NNCVBM"
table(com$Area, useNA = "always")
```

```
FLKCAR    NCA    NCFL    NED NNCVBM    <NA>
    306    153    153    153    306        0
```

The “rec_catch_TomFormat.csv” file contains all landings for the U.S. recreational fleet, which are all assumed to occur within the U.S. EEZs. There are no unreported catches in this region. This file also includes dead discards, which were calculated from the database by year and region and added into the total catch.

```
apply(rec[1:4], 2, table, useNA = "always")
```

\$Year

```
1986 1987 1988 1989 1990 1991 1992 1993 1994 1995 1996 1997 1998 1999 2000 2001
    32    32    32    32    32    32    32    32    32    32    32    32    32    32    32    32
2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017
    32    32    32    32    32    32    32    32    32    32    32    32    32    32    32    32
2018 2019 2020 2021 2022 <NA>
    32    32    32    32    32    0
```

\$Quarter

```
    1    2    3    4 <NA>
296 296 296 296    0
```

\$Fleet

```
Hire  Rec <NA>
 592  592    0
```

\$Area

```
FLK NCFL  NNC  VBM <NA>
296 296  296 296    0
```

15 Data compilation for all fleets in the operating model

```
rec$Area[which(rec$Area == "FLK")] <- "FLKCAR"  
rec$Area[which(rec$Area == "NNC")] <- "NNCVBM"  
rec$Area[which(rec$Area == "VBM")] <- "NNCVBM"  
table(rec$Area, useNA = "always")
```

FLKCAR	NCFL	NNCVBM	<NA>
296	296	592	0

Now we combine the data files and create some figures with the final composite data file.

```
d <- rbind(intc, intn, intw, com, rec)  
  
# how many rows contain zeros  
#hist(d$Catch_lbs)  
table(d$Catch_lbs == 0)
```

FALSE	TRUE
1920	651

```
# resort levels geographically  
d$Area <- factor(d$Area, levels = c("NCA", "FLKCAR", "NCFL", "NNCVBM", "NED"))  
d$Fleet <- factor(d$Fleet, levels = c("Rec", "Hire", "UScom", "Intl"))
```

15.2 Visualizing the final data set

We plot the final data set in a number of different ways to detect any errors that may have occurred during processing.


```
# plot the total landings by each factor  
par(mfrow = c(2, 2), mex = 0.7)  
barplot(tapply(d$Catch_lbs, d$Year, sum, na.rm = T)/10^6, las = 2,  
        main = "Total landings by year", ylab = "total landings (millions of pounds)")  
barplot(tapply(d$Catch_lbs, d$Quarter, sum, na.rm = T)/10^6,
```

15 Data compilation for all fleets in the operating model

```

ylab = "total landings (millions of pounds)", main = "Total landings by quarter")
barplot(tapply(d$Catch_lbs, d$Fleet, sum, na.rm = T)/10^6,
        main = "Total landings by fleet", ylab = "total landings (millions of pounds)")
barplot(tapply(d$Catch_lbs, d$Area, sum, na.rm = T)/10^6,
        main = "Total landings by area", ylab = "total landings (millions of pounds)")

```




The total landings in the regions analyzed is highly variable but relatively stable over time with a slight decrease in recent years. Most of the landings are caught in quarter 2 (March - May). The largest sector is the U.S. private recreational sector, followed by international fleets (all types), the for-hire fleet and then the U.S. commercial fleet. Most of the fishing activity takes place in the Florida Keys and the Greater Caribbean, with other regions contributing much less.

```

cols <- c("blue", "deepskyblue", "chartreuse3", "red")

par(mfrow = c(1, 1), mar = c(2, 5, 3, 1))
tab <- tapply(d$Catch_lbs, list(d$Fleet, d$Area), sum, na.rm = T)/10^6
barplot(tab, beside = T, col = cols, legend = rownames(tab),
        main = "Total dolphin landings by area and fleet",
        ylab = "total landings (millions of pounds)")

```




As expected, U.S. recreational fishing (private and for-hire) only takes place in the four U.S. EEZ regions (FLK, NCFL, NNC and VBM). The U.S. commercial fishery operates largely in those four U.S. EEZ regions, with small amounts of landings in the high seas of NCA, CAR and NED regions. International landings occur largely in the Caribbean, with lesser amounts in the NCA and NED high seas regions.

```
tab <- tapply(d$Catch_lbs, list(d$Quarter, d$Area), sum, na.rm = T)/10^6
tabp <- apply(tab, 2, function(x) x / sum(x, na.rm = T))

par(mar = c(2, 4, 3, 1))
barplot(tabp, beside = T, col = c("#8CD3DF", "#78B781", "#FFC107", "#A33851"),
        legend = c("winter (DJF)", "spring (MAM)", "summer (JJA)", "fall (SON)"),
        args.legend = list(x = 15, y = 0.7, bty = "n"),
        main = "Total dolphin landings by area and season",
        ylab = "proportion of landings")
```


Total dolphin landings by area and season



This plot shows the seasonal movements of dolphin across the different regions, as expressed through the proportion of landings that occurs in each quarter within each area. In the NCA and CAR regions, most of the landings occur in winter, whereas in the U.S. South Atlantic regions, most of the landings occur in spring and summer. In the U.S. Mid-Atlantic the landings also occur largely in spring, and also fall. In the NED region most of the landings occur in summer.

```
tab <- tapply(d$Catch_lbs, list(d$Year, d$Fleet), sum, na.rm = T)/10^6

par(mar = c(2, 4, 3, 1))
matplot(rownames(tab), tab,
        col = cols,
        type = "l", lwd = 2, lty = c(rep(1, 3), 2, 2),
        main = "Total annual dolphin landings by fleet", xlab = "",
        ylab = "total landings (millions of pounds)")
legend(2000, 27, lwd = 2, lty = c(rep(1, 3), 2, 2), bty = "n",
      c("U.S. private rec", "U.S. for-hire", "U.S. commercial",
        "International"),
      col = cols)
```



This plot shows the time series of total annual dolphin landings by fleet. We can compare these figures with the plots from the previous chapters to ensure the data were processed correctly. The magnitude and variability of the landings match the expectations.

```
# output the final concatenated data file
if (scenario == "high") {
  write.csv(d, file = "data/FINAL_files/complete_catch_dataset_100526_high.csv", row.names = F)
}
if (scenario == "low") {
  write.csv(d, file = "data/FINAL_files/complete_catch_dataset_100526_low.csv", row.names = F)
}

d1 <- d[which(d$Year == 2022), ]
d1$ar2 <- ""
d1$ar2[which(d1$Area == "FLK")] <- "S"
d1$ar2[which(d1$Area == "NCFL")] <- "S"
d1$ar2[which(d1$Area == "NNC")] <- "N"
d1$ar2[which(d1$Area == "VBM")] <- "N"
d1$cat <- paste0(d1$Fleet, d1$ar2)
d1$cat[which(d1$Fleet == "UScom")] <- "UScom"

round(tapply(d1$Catch_lbs, d1$cat, sum, na.rm = T)/10^6/2.205, 3)
```

15 Data compilation for all fleets in the operating model

Hire	HireS	Intl	Rec	RecS	UScom
0.509	0.044	2.513	2.784	1.248	0.106

16 Predictive indices of dolphin abundance

Dolphin abundance in South Atlantic waters is driven by a combination of factors that are mostly outside of U.S. fishery management control, including environmental conditions and catch in international waters. In order to create a management procedure for the dolphin management strategy evaluation we need an index that can predict ahead of time whether the abundance of dolphin in U.S. waters is likely to be above or below average in a given year. Ideally this index can capture the dynamics early in the year or a season ahead in order to allow adequate time for management response. In the following sections we explore potential predictive indices of dolphin abundance.

16.1 Recreational landings as a proxy for availability

Because the recreational landings is generally unconstrained by management (the total quota is rarely reached and landings are on average well below the bag limits), the total recreational landings could serve as a proxy for dolphin availability in South Atlantic waters over time. We take the recreational data from Section 3 and summarize according to the areas specified by the management strategy evaluation: Florida Keys to Indian River County County (FLK), Brevard to Southern NC South of Hatteras (NCFL), Northern NC North of Hatteras to NC/VA border (NNC), and Virginia to Maine (VBM). We also include Gulf of America landings because dolphin are not managed in the Gulf region and thus the landings are not constrained by management and is most likely a function of availability.

```
# clear workspace
rm(list = ls())

# install library
if(!require("readr")) install.packages("readr")
library(readr)


# read in data file
```

```
rec <- read.csv("data/recLandings.csv")
names(rec)[1] <- "Year"
```

16.2 Plot the recreational landings

Plotting the total recreational landings by region and the proportion of landings by region, we can observe that most of the landings in U.S. waters came from the FLK region, particularly in the earlier years. More recently, an increasing proportion of the landings has been coming from NCFL and further north, with FLK composing only about 50% of the landings. Gulf landings have historically been a small proportion of the total landings and have remained small throughout time. Landings in the VBM region were historically very small (~1%) but in recent years have varied between 10-18% of the landings.

```
# plot landings -----
matplot(rec$Year, rec[2:6]/10^6, type = "l", lty = 1, lwd = 2, col = 1:5,
        xlab = "year", ylab = "total landings (millions of pounds)",
        main = "Total recreational landings in U.S. waters by region")
legend("topright", colnames(rec)[2:6], col = 1:5, lty = 1, lwd = 2, bty = "n")
```




16 Predictive indices of dolphin abundance

```
# plot in terms of percentages -----
rec1 <- rec[2:6]
rec1 <- as.matrix(rec1/rowSums(rec1, na.rm = T))
rec1[is.na(rec1)] <- 0

b <- barplot(t(rec1), col = 1:5, names.arg = rec$Year, las = 1, ylim = c(0, 1.05),
             ylab = "proportion of landings",
             main = "Proportion of recreational landings in U.S. waters by region",
             legend.text = colnames(rec)[2:6], args.legend = list(x = 45, y = 1.15, horiz =
abline(h=0)
```

Proportion of recreational landings in U.S. waters by region



```
# calculate total landings and South Atlantic landings for later analysis
rec$ATL <- rowSums(rec[3:6], na.rm = T)
rec$ALL <- rec$GULF + rec$ATL

# subset data from 1990 forward
rec <- rec[which(rec$Year >= 1990), ]
```

Recreational monitoring was first established in 1979 and went through a number of changes in the 1980s, particularly related to monitoring of the for-hire fleet. We can

see a very large jump in landings in the late 1980s that we suspect is related to some of these changes in methodology. For our data explorations we will use recreational data from 1990 forward as a reference period in which the monitoring was more standardized.

16.3 Potential influence of El Niño Southern Oscillation

We note that many large peaks in recreational landings have occurred in years where there were severe El Niño events (e.g., 1997-1998, 2015). We first explore the El Niño Southern Oscillation (ENSO) as a potential environmental predictor of dolphin abundance. An index based on El Niño is potentially attractive because there is some ability for seasonal predictions of El Niño (3-6 months ahead of time). As large-scale disruptors of atmospheric conditions across the globe, El Niño events have impacts on the distribution of temperatures in the Atlantic and could trigger changes in fish distributions.

We use the multivariate ENSO index from NOAA's Physical Sciences Laboratory, which combines the influences of five different variables related to Pacific Ocean state into one index. The index is updated monthly and historical data can be downloaded directly from the site.


```
#download ENSO index
url <- "https://www.psl.noaa.gov/enso/mei/data/meiv2.data"
download.file(url = url, destfile = "data/enso.txt")

all_lines <- readLines("data/enso.txt")
footer_start <- grep("-999", all_lines)
e <- read.table("data/enso.txt", skip = 1, nrow = footer_start[1] - 2)
names(e) <- c("Year", "DJ", "JF", "FM", "MA", "AM", "MJ", "JJ", "JA", "AS", "SO", "ON", "ND")

ets <- as.vector(as.matrix(t(e[2:13])))
years <- sort(rep(e$Year, 12)) + rep((0:11)/12, length(min(e$Year):max(e$Year)))

barplot(ets, names.arg = floor(years), las = 2,
        col = ifelse(ets >= 0, "lightgreen", "red"), border = NA, ylim = c(-3, 3),
        main = "multivariate ENSO index", xlab = "Year", ylab = "ENSO index")
abline(h = 0)
text(346, 2, "El Niño events")
text(346, -2, "La Niña events")
```

16 Predictive indices of dolphin abundance



Here is the raw time series, showing positive and negative anomalies in the ENSO state.

Now we will compare the ENSO index to the recreational landings. First we compare the annual average ENSO index to the landings from the different regions. We report adjusted R-squared values and p-values for linear regression of the ENSO index versus the landings; significant relationships are highlighted in blue.

```
# calculate mean
e$mean <- rowMeans(e[2:13], na.rm = T)

# merge years with recreational landings data
d <- merge(rec, e, by = "Year")
#head(d)


lis <- names(d)[2:7]
par(mfrow = c(2, 3), mex = 0.8, mgp = c(2.5, 1, 0))
for(i in 1:6) {
  catch <- d[, which(names(d) == lis[i])]
  plot(d$mean, catch, xlab = "annual ENSO index", ylab = "landings (millions of lbs)",
       col = 8, main = paste0("region = ", lis[i]))
  out <- lm(catch ~ d$mean)
  abline(out, col = 8, lty = 2)
```

16 Predictive indices of dolphin abundance

```

r2 <- summary(out)$r.squared
p_val <- summary(out)$coefficients[2, 4]
p_display <- ifelse(p_val < 0.001, "p < 0.001", paste("p =", round(p_val, 3)))
co <- 1
if (p_val < 0.01) { co <- 4 }
if (p_val < 0.001) { co <- 5 }
legend("topleft",
      legend = bquote(R^2 == .(round(r2, 2)) ~ "; " ~ p == .(round(p_val, 3))),
      bty = "n", cex = 1.2, text.col = co)
}

```



The ENSO index appears to be most closely related to the total recreational landings across the South Atlantic (driven primarily by landings in the FLK region). The relationship is highly significant and the ENSO index explains 19% of the variation in the total

landings.


16.4 Seasonal influence of El Niño Southern Oscillation

Now we will consider the seasonal influence of ENSO, looking at correlations between the index during individual months and the landings. Since the strongest relationship occurs with the total South Atlantic landings, we will use these landings in the analysis.

```
lis <- names(e)[2:13]
par(mfrow = c(3, 4), mex = 0.8)
for(i in 1:12) {
  ind <- d[, which(names(d) == lis[i])]
  plot(ind, d$ATL/10^6, xlab = paste(lis[i], "ENSO index"), ylab = "landings (millions of 1
  out <- lm(d$ATL/10^6 ~ ind)
  abline(out)

  r2 <- summary(out)$r.squared
  p_val <- summary(out)$coefficients[2, 4]
  p_display <- ifelse(p_val < 0.001, "p < 0.001", paste("p =", round(p_val, 3)))
  co <- 1
  if (p_val < 0.01) { co <- 4 }
  if (p_val < 0.001) { co <- 5 }
  legend("topleft",
        legend = bquote(R^2 == .(round(r2, 2)) ~ "; " ~ p == .(round(p_val, 3))),
        bty = "n", cex = 1.2, text.col = co)
}
```

16 Predictive indices of dolphin abundance




A significant relationship between the monthly ENSO index and total South Atlantic recreational landings occurs with the May-June index, but the strongest relationship occurs in June-July (with 26% of the variance in landings explained by the index in that month). The relationship in the late summer and fall months are also significant, which is expected since the ENSO index is highly correlated from month to month.

We will test the June-July ENSO index in the management procedure as it has the strongest relationship.

```
par(mfrow = c(1, 1))
plot(d$JJ, d$ATL/10^6, xlab = "June-July ENSO index", ylab = "total U.S. Atlantic coast rec
text(d$JJ, d$ATL/10^6, d$Year, col = 8)
out <- lm(d$ATL/10^6 ~ d$JJ)
abline(out, col = 1)
p <- summary(out)$coef[2, 4]
```

16 Predictive indices of dolphin abundance

```
legend("topleft", paste0("R^2 = ", round(summary(out)$adj.r.squared, 2), "; p = ",
                           round(p, 3)), cex = 1.2, bty = "n", text.col = 1)
```



```
summary(out)
```

Call:

16 Predictive indices of dolphin abundance

```
lm(formula = d$ATL/10^6 ~ d$JJ)
```

Residuals:

Min	1Q	Median	3Q	Max
-10.332	-3.886	-0.054	2.320	10.606

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	17.2374	0.8641	19.949	< 2e-16 ***
d\$JJ	3.0873	0.8680	3.557	0.00116 **

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 5.046 on 33 degrees of freedom

Multiple R-squared: 0.2771, Adjusted R-squared: 0.2552

F-statistic: 12.65 on 1 and 33 DF, p-value: 0.001161

```
# save the index to the indices file
```

```
ind <- data.frame(cbind(d$Year, d$JJ))
```

```
write.csv(ind, file = "indices/enso_index.csv", row.names = F)
```

17 Wider Atlantic temperature distribution index

Dolphins have a circumtropical distribution, migrating to temperate waters in summer and tropical waters in winter, in a circular pattern around the Atlantic Ocean. Studies using satellite tagging have shown that, similar to other highly migratory species, they prefer waters within a narrow temperature range, of 27-30 degrees Celsius Schlenker et al. 2021. To determine whether shifting temperature distributions across the Atlantic could be impacting the abundance in U.S. waters, we analyzed Atlantic Ocean temperature patterns at the large spatial and temporal scale.

17.1 Accessing temperature data

For this analysis we need a historical temperature record that is available at large spatial (entire Atlantic) and temporal (monthly-yearly) scales. We use NOAA's Extended Reconstructed Sea Surface Temperature (ERSST) data, provided by the NOAA National Centers for Environmental Information and available through ERDDAP. This temperature record goes back to the mid-1800s, is available at monthly time steps and a 2°x2° grid size resolution, and is suitable for looking at large-scale patterns over decades. We download the data from 1980 - present and subset for the Atlantic Ocean from 100°W - 0°W longitude and 0°N - 70°N latitude.

```
# clear workspace
rm(list = ls())

# install libraries
if(!require("maps")) install.packages("maps")
# if(!require("ncdf4")) install.packages("ncdf4")
# if(!require("lubridate")) install.packages("lubridate")
# if(!require("rerddap")) install.packages("rerddap")

library(maps)
```

17 Wider Atlantic temperature distribution index

```
# library(ncdf4)
# library(lubridate)
# library(rerddap)

# get ERDDAP info
# id <- info('nceiErsstv5') #https://coastwatch.pfeg.noaa.gov/erddap/griddap/nceiErsstv5.

# download data using griddap - initially takes a long time
# uncomment lines below to update analysis
# sst_grab <- griddap(id, fields = 'sst',
#                     time = c('1990-01-01', '2025-01-01'),
#                     longitude = c(260, 358), latitude = c(0, 70))

# open nc file and extract variables
# nc <- nc_open(sst_grab$summary$filename, write=FALSE, readunlim=TRUE, verbose=FALSE)
# v1 <- nc$var[[1]]
# sst <- ncvar_get(nc, v1)
# lon <- v1$dim[[1]]$vals - 360
# lat <- v1$dim[[2]]$vals
# nc_close(nc)

# convert time variable to month and year
#tim <- as.POSIXct(v1$dim[[4]]$vals, origin = "1970-01-01")
#mon <- as.numeric(substr(tim, 6, 7))
#yr <- substr(tim, 1, 4)
#dim(sst)

#save(sst, lon, lat, tim, mon, yr, file = "data/ERSST.RData")

load("data/ERSST.RData")

# check conversions
tim[1:5]
```

```
[1] "1990-01-01 UTC" "1990-02-01 UTC" "1990-03-01 UTC" "1990-04-01 UTC"
[5] "1990-05-01 UTC"
```

```
table(mon)
```

```
mon
```

17 Wider Atlantic temperature distribution index

```

1  2  3  4  5  6  7  8  9 10 11 12
36 35 35 35 35 35 35 35 35 35 35 35

```

```
table(yr)
```

```

yr
1989 1990 1991 1992 1993 1994 1995 1996 1997 1998 1999 2000 2001 2002 2003 2004
      1   12   12   12   12   12   12   12   12   12   12   12   12   12   12   12
2005 2006 2007 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020
      12   12   11   12   12   12   12   12   12   12   12   12   12   12   12   12
2021 2022 2023 2024 2025
      12   12   12   12   1


```

```

# plot SST on map for first time step
par(mfrow = c(1, 2))
image(lon, lat, sst[, , 100], las = 1, xlab = "longitude (deg W)", ylab = "latitude (deg N)",
      main = "Snapshot of ERSST data"); box()

# fill Pacific Ocean with NAs so values are not included
sst[which(lon <= (-92)), which(lat <= 17.5), ] <- NA
sst[which(lon <= (-85)), which(lat <= 15), ] <- NA
sst[which(lon <= (-83)), which(lat <= 10), ] <- NA
sst[which(lon <= (-77)), which(lat <= 8.5), ] <- NA
sst[which(lon <= (-78) & lon >= (-80)), which(lat <= 9), ] <- NA
image(lon, lat, sst[, , 100], las = 1, xlab = "longitude (deg W)", ylab = "latitude (deg N)",
      main = "Snapshot of ERSST data\n(Pacific excluded)"); box()

```



17.2 Analyzing high and low availability years

In order to identify potential temperature patterns that are influential in driving dolphin abundance in the U.S. South Atlantic waters, we identify the years of highest and lowest landings and look at the temperature distributions monthly in those years.

```
# upload recreational landings
rec <- read.csv("data/recLandings.csv")
names(rec)[1] <- "Year"

# summarize by basin and total
rec$ATL <- rowSums(rec[3:6], na.rm = T)
rec$ALL <- rec$GULF + rec$ATL

# consider 1990 and forward
rec <- rec[which(rec$Year >= 1990), ]

# sort years from lowest to higher landings
sortyr <- rec$Year[order(rec$ALL)]
```

17 Wider Atlantic temperature distribution index

```
# cols <- rainbow(100, start = 0.1, end = 0.65)[100:1]
cols <- c(0, rainbow(3, start = 0.4, end = 0.6), 2)

# set up plot
nf <- layout(matrix(c(2:56, rep(1, 11), 57:111), 11, 11, byrow = FALSE),
                 c(rep(3, 5), 2, rep(3, 5)), c(4, rep(3, 11)))
#layout.show(nf)

#legend
par(mar = c(0, 0, 0, 0))
plot(1, type = "n", axes = FALSE, xlab = "", ylab = "")
legend("center", fill = cols, c("<26", "26-27", "27-28.5", "28.5-30", ">30"),
      border = c(1, 0, 0, 0, 0),
      pt.cex = 2, bty = "n",
      title = "SST\n(deg C)", title.font = 2)


# sort years from lowest to highest landings
yrs <- c(sortyr[1:5], sortyr[(length(sortyr)-4): length(sortyr)])
#yrs
status <- c(rep("low", 5), rep("high", 5))
status[1] <- "lowest"
status[10] <- "highest"

# for lowest and highest years, visualize temperature patterns across the Atlantic
for (j in 1: length(yrs)) {
  k <- which(yr == yrs[j])
  # print(data.frame(tim[k], yr[k], mon[k]))
  par(mex = 0.5, mar = c(2, 2, 2, 2)+1)
  plot(rec$Year, rec$ALL/10^6, type = "l", col = 4, ylim = c(0, 32), lwd = 2, ylab = "rec 1",
       main = yrs[j])
  text(1990, 3, status[j], pos = 4)
  abline(v = yrs[j], col = 8, lwd = 3)

  # plot months February through November
  for (i in k[2:11]) {
    par(mex = 0.3, mar = c(2, 3, 2, 0))
    map("world", xlim = c(-100, -15), ylim = c(0, 45), col = 0)
    axis(1); axis(2, las = 2); box()
    image(lon, lat, sst[, , i], add = T, col = cols, breaks = c(-2, 26, 27, 28.5, 30, 32))
    text(-33, 40, paste(month.abb[mon[i]]), cex = 1)
  }
}
```

17 Wider Atlantic temperature distribution index

```
map("world", add = T, col = gray(0.5)); box()
contour(lon, lat, sst[, , i], levels = c(30), add = T, col = c(2), lwd = 2, drawlabels =
contour(lon, lat, sst[, , i], levels = c(28.5), add = T, col = 1, lwd = 2, drawlabels =
}
}
```



Looking at the set of plots on the left side (low abundance years) versus the right side (high abundance years), there seems to be a particularly notable pattern with respect to temperatures in the range of 28.5° - 30° Celsius. In the low abundance years, temperatures in this range are distributed in a band across the Atlantic near the

equator. In the high abundance years, waters near the equator are much cooler and the 28.5° - 30° temperatures appear to be concentrated in a very small area near the Florida Keys. This pattern seems most prominent in the months of May and June.

17.3 May temperature distributions

We will take a closer look at May temperature distributions across the Atlantic for the full time series. Here we display all of the May temperatures from 1990 to present, ordered from lowest to highest years of recreational landings in the U.S. South Atlantic.


```
# look at May patterns -----

par(mfrow = c(6, 6), mex = 0.5, mar = c(2, 2, 2, 0))

for (j in 1: length(sortyr)) {
  k <- which(yr == sortyr[j])
  for (i in k[5]) {
    map("world", xlim = c(-100, -10), ylim = c(0, 50), col = 0)
    axis(1); axis(2, las = 2); box()
    image(lon, lat, sst[, , i], add = T, col = cols, breaks = c(-2, 26, 27, 28.5, 30, 32))
    map("usa", add = T, col = gray(0.5)); box()
    mtext(side = 3, line = 0.5, sortyr[j])
    contour(lon, lat, sst[, , i], levels = c(30), add = T, col = c(2), lwd = 2, drawlabels = F)
    contour(lon, lat, sst[, , i], levels = c(28.5), add = T, col = 1, lwd = 2, drawlabels = F)
    rect(-100, 0, -30, 32, col = NA, border = 2, lwd = 1.5)
    rect(-99, 18, -78, 31, col = NA, border = 7, lwd = 1.5)
  }
}

par(mar = c(0, 0, 0, 0))
plot(1, type = "n", axes = FALSE, xlab = "", ylab = "")
legend("center", fill = cols, c("<26", "26-27", "27-28.5", "28.5-30", ">30"),
      border = c(1, 0, 0, 0, 0),
      pt.cex = 2, bty = "n",
      title = "SST (deg C)", title.font = 2)
```

17 Wider Atlantic temperature distribution index



The pattern with 28.5° - 30° C waters seems to hold across all years, with low abundance years (top) tending to have these waters distributed across the Atlantic, and high abundance years (bottom) tending to have these waters restricted to a small area near the Gulf and U.S. South Atlantic.

17.4 Quantifying the temperature distribution in an index

We can attempt to quantify these patterns in an index by calculating a proportion of waters covered by the dark blue water mass. It seems most useful to consider the coverage of this water mass that occurs in U.S. South Atlantic waters (approximately the yellow box; 100°W-78°W and 18°N-32°N) versus the coverage of this water mass in the larger North Atlantic region (the larger red box; 100°W-30°W and 0°-32°N). We first calculate the number of cells in which these 28.5° - 30° C waters occur in each box, and then divide to obtain an overall proportion.

```
# calculate proportion of 28.5 - 30 degree C waters in US South Atlantic waters
```

```
# subset the temperature data to 100W - 30W and 0N - 32N
```

```
lons <- which(lon >= (-100) & lon <= (-30))
```

```
lats <- which(lat >= (0) & lat <= (32))
```

```
lon[lons]
```

```
[1] -100 -98 -96 -94 -92 -90 -88 -86 -84 -82 -80 -78 -76 -74 -  
72  
[16] -70 -68 -66 -64 -62 -60 -58 -56 -54 -52 -50 -48 -46 -44 -  
42  
[31] -40 -38 -36 -34 -32 -30
```

```
lat[lats]
```

```
[1] 0 2 4 6 8 10 12 14 16 18 20 22 24 26 28 30 32
```

```
# find indexing for 100W - 78W and 18N - 32N
```

```
lonind <- which(lon[lons] <= (-78) & lon[lons] >= (-100))
```

```
latind <- which(lat[lats] <= 32 & lat[lats] >= 18)
```

```
# subset SST data for all May months
```

```
kar <- sst[lons, lats, which(mon == 5)]
```

```
# create empty data frame
```

```
ind <- data.frame(matrix(ncol = 2, nrow = dim(kar)[3],  
                        dimnames = list(NULL, c("Year", "perar"))))
```

```
ind$Year <- yr[which(mon == 5)]
```

17 Wider Atlantic temperature distribution index

```
# for each year, identify SST cells between 28.5 and 30 degrees
# calculate the proportion of those waters within the small box out of the large box
for (i in 1:dim(kar)[3]) {
  k1 <- kar[, , i]
  k2 <- (k1 >28.5 & k1 < 30.0)
  ind$perar[i] <- length(which(k2[lonind, latind] == TRUE))/ length(which(k2 == TRUE))
}

# replace NA values with zero
ind$perar[which(is.na(ind$perar))] <- 0
ind
```

	Year	perar
1	1990	0.8666667
2	1991	1.0000000
3	1992	0.7000000
4	1993	0.1666667
5	1994	0.9629630
6	1995	0.7692308
7	1996	0.8666667
8	1997	0.8500000
9	1998	0.3402778
10	1999	0.9393939
11	2000	1.0000000
12	2001	1.0000000
13	2002	0.8076923
14	2003	0.8723404
15	2004	0.9411765
16	2005	0.2214286
17	2006	0.4444444
18	2007	0.3888889
19	2008	0.0000000
20	2009	0.0000000
21	2010	0.0000000
22	2011	0.0000000
23	2012	0.4000000
24	2013	0.0000000
25	2014	0.0000000
26	2015	0.7500000
27	2016	0.2343750

17 Wider Atlantic temperature distribution index


```
28 2017 0.0000000
29 2018 0.0000000
30 2019 0.0000000
31 2020 0.1929825
32 2021 0.0000000
33 2022 0.0000000
34 2023 0.2500000
35 2024 0.1538462
```

```
yrs <- sortyr[order(sortyr)]

par(mfrow = c(6, 6), mex = 0.5, mar = c(2, 2, 2, 0))
for (j in 1:length(yrs)) {
  k <- which(yr == yrs[j])
  for (i in k[5]) {
    map("world", xlim = c(-100, -10), ylim = c(0, 50), col = 0)
    axis(1); axis(2, las = 2); box()
    image(lon, lat, sst[, , i], add = T, col = cols, breaks = c(-2, 26, 27, 28.5, 30, 32))
    map("usa", add = T, col = gray(0.5)); box()
    mtext(side = 3, line = 0.5, yrs[j])
    text(-38, 45, paste("index =", round(ind$perar[which(ind$Year == yrs[j])], 2)))
    contour(lon, lat, sst[, , i], levels = c(30), add = T, col = c(2), lwd = 2, drawlabels = F)
    contour(lon, lat, sst[, , i], levels = c(28.5), add = T, col = 1, lwd = 2, drawlabels = F)
    rect(-100, 0, -30, 32, col = NA, border = 2, lwd = 1.5)
    rect(-99, 18, -78, 31, col = NA, border = 7, lwd = 1.5)
  }
}

par(mar = c(0, 0, 0, 0))
plot(1, type = "n", axes = FALSE, xlab = "", ylab = "")
legend("center", fill = cols, c("<26", "26-27", "27-28.5", "28.5-30", ">30"),
      border = c(1, 0, 0, 0, 0),
      pt.cex = 2, bty = "n",
      title = "SST (deg C)", title.font = 2)
```

17 Wider Atlantic temperature distribution index



To ensure correct calculation of the index we re-display the May temperature patterns in chronological order and note the index value on each plot.

17.5 Comparison of “blue blob” index with landings


Now we plot the “blue blob” index against the recreational landings.

17 Wider Atlantic temperature distribution index

```
# merge new index with landings data
d <- merge(rec, ind, by = "Year")
#head(d)

# plot the "blue blob" index against recreational landings
par(mar = c(5, 5, 1, 1))
plot(d$perar, d$ATL/10^6, col = 0,
      xlab = "proportion of preferred dolphin temperatures near\nU.S. jurisdictions (out of",
      ylab = "total U.S. Atlantic coast recreational landings\n(millions of pounds)")
text(d$perar, d$ATL/10^6, d$Year)
out <- lm(d$ATL/10^6 ~ d$perar)
abline(out, lty = 2, col = 2)
abline(v = mean(d$perar), col = 8)
abline(h = mean(d$ATL)/10^6, col = 8)
text(0.7, 28, "preferred temperatures clustered\nnear U.S. waters == higher landings", col = 8)
text(0.25, 10.5, "preferred temperatures extend across\nCaribbean and/or absent from\nU.S w
```

17 Wider Atlantic temperature distribution index



We can see that a statistical relationship does exist, whereby above average “blue blob” index values are associated with above average recreational landings in U.S. South Atlantic waters, and below average index values are associated with below average landings. Gray lines denote the mean values for the index and the landings; we see that this relationship holds true for almost all years in the time series.

The “blue blob” index explains about 45% of the variation in the landings in the U.S. South Atlantic.

```
summary(out)
```

```
Call:
lm(formula = d$ATL/10^6 ~ d$perar)
```

17 Wider Atlantic temperature distribution index

Residuals:

Min	1Q	Median	3Q	Max
-9.9082	-3.0254	0.0927	2.6638	9.6171

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)	
(Intercept)	12.391	1.094	11.324	6.58e-13	***
d\$perar	10.079	1.880	5.361	6.35e-06	***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 4.339 on 33 degrees of freedom

Multiple R-squared: 0.4655, Adjusted R-squared: 0.4493

F-statistic: 28.74 on 1 and 33 DF, p-value: 6.352e-06

```
# output the index
write.csv(ind, file = "indices/blue_blob_index.csv", row.names = F)
```

18 Puerto Rico tournament index

In this section we consider tournament data from the U.S. Caribbean as a potential predictive index for dolphin abundance in the South Atlantic. Because dolphins migrate through the Caribbean region in the months before arriving in South Atlantic waters, peaks in catch-per-unit effort for fisheries operating in the Caribbean could potentially be indicative of high abundances later in the year in waters downstream of their migration patterns. Tournament data are available from Puerto Rico's Departamento de Recursos Naturales y Ambientales, División de Pesquería Recreativa y Deportiva, going back to the year 2000. Tournament data were requested and received from the institution in June 2022.

18.1 Upload and clean data set

We first input the data set, parse out the dates to extract month and year, and standardize labeling of months.

```
# clear workspace
rm(list = ls())

if(!require("dplyr")) install.packages("dplyr")
if(!require("emmeans")) install.packages("emmeans")

library(dplyr)
library(emmeans) # Best for extracting standardized indices

# import data -----
d <- read.csv("data/PRDNER-DolTournamentData.csv")
#apply(d, 2, table)
head(d)
```

	Date	Location	Total.Number.Participants
1	May-6-00	Club N\xeelutico de La Parguera	124

18 Puerto Rico tournament index

2	May-6-00	Club N\xelutico de La Parguera	124
3	May-6-00	Club N\xelutico de La Parguera	124
4	May-6-00	Club N\xelutico de La Parguera	124
5	May-6-00	Club N\xelutico de La Parguera	124
6	May-6-00	Club N\xelutico de La Parguera	124
	Total.Number.of.Boats	Average.Time.Spent.Fishing	Tournament.Duration
1	31	10.5	2
2	31	10.5	2
3	31	10.5	2
4	31	10.5	2
5	31	10.5	2
6	31	10.5	2
	Fish.Type	Fish.name	Sex
1	8835290101	Dolphinfish	F
2	8835290101	Dolphinfish	F
3	8835290101	Dolphinfish	M
4	8835290101	Dolphinfish	F
5	8835290101	Dolphinfish	F
6	8835290101	Dolphinfish	F
	Boarded	Bycatch	Lenght..mm.
1	TRUE	FALSE	1675
2	TRUE	FALSE	1675
3	TRUE	FALSE	1675
4	TRUE	FALSE	1625
5	TRUE	FALSE	1625
6	TRUE	FALSE	1625
	Weight..Kg.	Distance.to.coast	Zone
1	9.09	2	
2	6.81	2	
3	10.45	2	
4	10.00	2	
5	12.27	2	
6	10.00	2	

```
# clean dates and extract month day year -----
d$Date <- as.character(d$Date)
d$month <- NA
d$day <- NA
d$year <- NA

for (i in 1:nrow(d)) {# M
  d[i, 16:18] <- unlist(strsplit(d$Date[i], "-")) }
#head(d)

# clean up errors and check outputs -----
d$month <- substr(d$month, 1, 3)
d$month[which(d$month == "Abr")] <- "Apr"
d$month[which(d$month == "Arp")] <- "Apr"
```

18 Puerto Rico tournament index

```
d$day[which(d$day == "November" | d$day == "December")] <- NA
d$year[which(d$year == "November" | d$year == "December")] <- NA

d$day <- as.numeric(d$day)
d$year <- as.numeric(d$year)

# standardize year format
d$year[which(d$year < 83)] <- d$year[which(d$year < 83)] + 2000
d$year[which(d$year < 2000)] <- d$year[which(d$year < 2000)] + 1900

d$mon <- match(d$month, month.abb)

table(d$month)
```

Apr	Aug	Dec	Feb	Jan	Jul	Jun	Mar	May	Nov	Oct	Sep
7723	57	614	2360	2160	35	23	7995	1496	1036	1344	273

```
table(d$day)
```

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
694	829	792	689	718	1123	618	753	733	1417	493	775	1061	936	278	517
17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	
894	859	524	1213	726	875	880	655	661	808	854	1192	850	563	1050	

```
table(d$year)
```

1983	1984	1985	1986	1987	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
80	153	5	53	19	843	1254	1413	1040	1303	944	1411	1418	1209	1190	538
2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022				
1263	1815	1321	1153	1512	916	944	808	1213	51	91	1070				

In keeping with the definition of seasons in other parts of the MSE, we group December with the following January and February in a winter season.

18 Puerto Rico tournament index

```
# fix so Dec is grouped with following year -----
d$mon[which(d$mon == 12)] <- 0.5
d$year2 <- d$year
d$year2[which(d$mon == 0.5)] <- d$year[which(d$mon == 0.5)]+1
rbind(table(d$year), table(d$year2))
```

	1983	1984	1985	1986	1987	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
[1,]	80	153	5	53	19	843	1254	1413	1040	1303	944	1411	1418	1209	1190
[2,]	80	153	5	53	19	789	1220	1501	1030	1313	944	1411	1418	863	1536
	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022		
[1,]	538	1263	1815	1321	1153	1512	916	944	808	1213	51	91	1070		
[2,]	538	1263	1815	1321	1153	1512	915	945	801	1190	81	85	1076		


```
#head(d)
#tail(d)
```

18.2 Viewing the raw data

Let's take a look at some of the data columns using barplots and histograms.


```
# look at data columns -----
par(mfrow = c(3, 2))
barplot(table(d$mon), main = "Number of observations - Month")
barplot(table(d$year), main = "Number of observations - Year")
barplot(table(d$day), main = "Number of observations - Day")
barplot(table(d$Sex), main = "Number of observations - Sex")
barplot(table(d$Zone), main = "Number of observations - Zone")
barplot(table(d$Bycatch), main = "Number of observations - Bycatch")
```

18 Puerto Rico tournament index



```
par(mfrow = c(2, 2))
hist(d$Total.Number.Participants, main = "Total number of participants")
hist(d$Total.Number.of.Boats, main = "Total number of boats")
hist(d$Average.Time.Spent.Fishing, main = "Average time spent fishing")
hist(d$Tournament.Duration, main = "Tournament duration")
```

18 Puerto Rico tournament index



There are a few NAs in the tournament data columns. We need to consider what details are available and what factors we can feasibly use in the standardization process.

```
# find NAs in tournament details -----  
table(is.na(d$Total.Number.of.Boats))
```

```
FALSE  TRUE  
24331  785
```

```
table(is.na(d$Total.Number.Participants))
```

18 Puerto Rico tournament index

```
FALSE TRUE
24189  927
```


```
table(is.na(d$Total.Number.Participants) & is.na(d$Total.Number.of.Boats))
```

```
FALSE TRUE
24332  784
```

```
table(is.na(d$Average.Time.Spent.Fishing))
```

```
FALSE TRUE
23290 1826
```

```
plot(d$Total.Number.of.Boats, d$Total.Number.Participants, xlab = "total number of boats", y
out <- lm(d$Total.Number.Participants ~ d$Total.Number.of.Boats)
abline(out, col = 8)
```



The total number of boats is highly correlated with the total number of participant and the linear regression indicates that most boats have 4 participants.

18.3 Data preparation for standardization

In preparation for creating a standardized catch-per-unit-effort (CPUE) index that can serve as a proxy for abundance, we have to specify nuisance factors in the data set

18 Puerto Rico tournament index

that have to be accounted for. We create a unique tournament ID number with a combination of the day and number of participants.

```
# create unique tournament ID number -----  
d$dat <- paste0(d$day, d$month, d$year)  
length(unique(paste0(d$day, d$month, d$year, d$Total.Number.Participants)))
```

```
[1] 386
```

```
length(unique(paste0(d$day, d$month, d$year, d$Location)))
```

```
[1] 387
```

```
length(unique(paste0(d$day, d$month, d$year, d$Total.Number.of.Boats)))
```

```
[1] 386
```

```
length(unique(paste0(d$day, d$month, d$year, d$Tournament.Duration)))
```

```
[1] 366
```

```
d$datID <- paste0(d$day, d$month, d$year, d$Total.Number.Participants)  
  
#barplot(table(d$mon))
```

We remove all cases where dolphin is not listed as the target species; we will use only directed dolphin trips in the standardization. We also specify the months that we want to include in the index of abundance. Here we specify December to MARCH With this subset of months, we have 128 unique tournaments in the data base across 121 different dates.

```
# remove non-target cases; use only directed trips -----  
dfull <- d  
d <- d[which(d$Bycatch == FALSE), ]  
  
# subset by season, e.g. 0 - 4 is December to April -----  
d <- d[which(d$mon >= 0 & d$mon <= 3), ]
```

18 Puerto Rico tournament index

```
#length(unique(d$Date))  
length(unique(d$dat))
```

```
[1] 121
```

```
length(unique(d$datID))
```

```
[1] 128
```

Next we calculate the total weight by tournament, by summing the reported weights across each tournament ID. We then set up a new data frame where each tournament is a row, and columns represent the attributes of each tournament (year, month, zone, participants, number of boats, fishing time, duration, total weight and total abundance). We also calculate the average weight by dividing total weight by total abundance.

```
# calculate total by tournament -----  
totWT <- tapply(d$Weight..Kg., d$datID, sum, na.rm = T)  
totN <- table(d$datID)  
mon <- tapply(d$mon, d$datID, mean, na.rm = T)  
year <- tapply(d$year2, d$datID, mean, na.rm = T)  
  
# checks on month and tournament assignments -----  
#table(d$year[d$mon == 0.5], d$year2[d$mon == 0.5])  
#table(d$year[d$mon != 0.5], d$year2[d$mon != 0.5])  
  
max(tapply(d$Total.Number.Participants, d$datID, sd), na.rm = T) # should all be zeros
```

```
[1] 0
```

```
max(tapply(d$Total.Number.of.Boats, d$datID, sd), na.rm = T)
```

```
[1] 0
```

```
max(tapply(d$Average.Time.Spent.Fishing, d$datID, sd), na.rm = T)
```

```
[1] 0
```

18 Puerto Rico tournament index

```
# assign attributes of tournaments -----
Npart <- tapply(d$Total.Number.Participants, d$datID, mean, na.rm = T)
Nboat <- tapply(d$Total.Number.of.Boats, d$datID, mean, na.rm = T)
Ftime <- tapply(d$Average.Time.Spent.Fishing, d$datID, mean, na.rm = T)
dura <- tapply(d$Tournament.Duration, d$datID, mean, na.rm = T)
zone <- tapply(d$Zone, d$datID, mean, na.rm = T)

# new data frame for standardization -----
dat <- data.frame(cbind(year, mon, zone, Npart, Nboat, Ftime, dura, totWT, totN))
dat$avwt <- dat$totWT / dat$totN

head(dat)
```

	year	mon	zone	Npart	Nboat	Ftime	dura	totWT	totN	avwt
10Feb2007140	2007	2	1	140	35	8.0	1	347.64	49	7.094694
10Feb200824	2008	2	1	24	15	7.0	1	264.71	33	8.021515
10Jan2009332	2009	1	1	332	83	8.0	2	770.44	147	5.241088
10Jan2014188	2014	1	1	188	47	7.5	2	1479.82	249	5.943052
10Jan2015136	2015	1	1	136	34	7.5	2	794.72	264	3.010303
10Mar2001192	2001	3	2	192	48	8.0	2	2499.33	272	9.188713

There are a number of missing values in this data frame. For fishing time and number of boats, we impute the median of the known values into the missing values. For missing values for the number of boats, we take the number of participants and divide by 4 (given the relationship established above). If both values are missing, we impute the median into the unknown values.

```
# fill in NAs -----
dat$Nboat[which(dat$Nboat == 0)] <- NA
dat$Npart[which(dat$Npart == 0)] <- NA
dat$Ftime[which(dat$Ftime == 0)] <- NA

#hist(dat$Ftime)
which(is.na(dat$Ftime))
```

```
[1] 15 18 68 72 86 94 101 127 128
```

18 Puerto Rico tournament index

```
dat$Ftime[is.na(dat$Ftime)] <- median(dat$Ftime, na.rm = T)
```

```
#hist(dat$Nboat)
which(is.na(dat$Nboat))
```

```
[1] 15 83 94 122 128
```

```
which(is.na(dat$Npart))
```

```
[1] 11 12 15 94 101 110 122 128
```

```
dat$Npart[is.na(dat$Nboat)]
```

```
[1] NaN 33 NaN NA NaN
```

```
#plot(dat$Nboat, dat$Npart)
#dat$Npart / dat$Nboat
dat$Nboat[is.na(dat$Nboat)] <- dat$Npart[is.na(dat$Nboat)] / 4
#dat$Npart[is.na(dat$Nboat)]
dat$Nboat[is.na(dat$Nboat)] <- median(dat$Nboat, na.rm = T)

which(is.na(dat$Nboat))
```

```
integer(0)
```

```
which(is.na(dat$Ftime))
```

```
integer(0)
```

```
which(is.na(dat$totN))
```

```
integer(0)
```

Finally, we subset the years. Data are very sparse prior to 2000 so we only include the data for 2000 and forward. We calculate the catch-per-unit-effort, defining catch as the total abundance, and effort as the number of boats multiplied by the average time spent fishing.

18 Puerto Rico tournament index

```
# subset years -----  
  
yrs <- c(2000:2020, 2022)  
  
dat[is.na(dat)] <- NA  
dat <- dat[which(dat$year >= min(yrs)), ]  
dat <- dat[which(dat$year <= max(yrs)), ]  
  
# calculate CPUE -----  
dat$eff <- dat$Nboat * dat$Ftime # dat$dura  
dat$cpue <- dat$totN / dat$eff  
dat$cpueW <- dat$totWT / dat$eff  
  
which(dat$cpue == "Inf")
```


```
integer(0)
```

```
which(dat$cpueW == "Inf")
```

```
integer(0)
```

```
#dat$cpue[which(dat$cpue == "Inf")] <- NA
```

```
barplot(table(dat$year), las = 2, main = "Number of tournaments per year")
```



The plot shows the number of tournaments per year. We can see that there are relatively few events for each year.

18.4 Analyze the nominal CPUE trends

Now using this clean data set we can look at the nominal (average) CPUE by year. We first look at the average weight across all tournaments, and weight by the northern and southern coasts. The weights appear to be highly variable across year and coasts.


```
# look at nominal trends -----
par(mfrow = c(3, 1), mar = c(3, 5, 3, 1))
barplot(tapply(dat$avwt, dat$year, mean, na.rm = T), las = 2,
        main = "Average weight of tournament-caught dolphinfish by year",
        ylab = "average weight (kg)")

d2 <- dat[which(dat$zone == 2), ]
barplot(tapply(d2$avwt, d2$year, mean, na.rm = T), las = 2,
        main = "Average weight of tournament-caught dolphinfish by year\n(Southern coast)",
        ylab = "average weight (kg)")
```

18 Puerto Rico tournament index

```
d2 <- dat[which(dat$zone != 2), ]  
barplot(tapply(d2$avwt, d2$year, mean, na.rm = T), las = 2,  
        main = "Average weight of tournament-caught dolphinfish by year\n(other coasts)",  
        ylab = "average weight (kg)")
```




18 Puerto Rico tournament index



18 Puerto Rico tournament index


```
# look at nominal trends by year, month and zone
barplot(tapply(dat$cpue, dat$year, mean, na.rm = T), las = 1,
        main = "Average CPUE of tournament-caught dolphinfish by year",
        xlab = "year", ylab = "CPUE")
barplot(tapply(dat$cpue, dat$mon, mean, na.rm = T), las = 1,
        main = "Average CPUE of tournament-caught dolphinfish by month",
        xlab = "month", ylab = "CPUE")
barplot(tapply(dat$cpue, dat$zone, mean, na.rm = T), las = 1,
        main = "Average CPUE of tournament-caught dolphinfish by zone",
        xlab = "zone", ylab = "CPUE")
```

18 Puerto Rico tournament index



18 Puerto Rico tournament index

```
par(mfrow = c(1, 1), mar = c(5, 5, 1, 1))
plot(tapply(dat$cpue, dat$year, mean, na.rm = T), tapply(dat$cpueW, dat$year, mean, na.rm = T),
     xlab = "nominal CPUE by number", ylab = "nominal CPUE by weight")
```



```
cor(tapply(dat$cpue, dat$year, mean, na.rm = T), tapply(dat$cpueW, dat$year, mean, na.rm = T))
```

```
[1] 0.8957364
```

There appears to be high variability in the CPUE by year. The highest CPUE appears to occur in December and March. CPUE also varies by zone, the Southern coast having the highest catch rates. CPUE based on number and weight are highly correlated.

18.5 Calculate standardized CPUE trends

Now we will carry out the standardization, using day, month and zone as standardization factors. Since the catch rates are highly skewed, we carry out the standardization based on the log CPUE.

```
# convert variables to factors -----
dat$year <- as.factor(dat$year)
dat$mon <- as.factor(dat$mon)
dat$zone <- as.factor(dat$zone)

# standardization -----

out <- glm(log(dat$cpue) ~ dat$year + dat$mon + dat$zone,
           family = gaussian)
summary(out)
```

Call:

```
glm(formula = log(dat$cpue) ~ dat$year + dat$mon + dat$zone,
     family = gaussian)
```

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	-1.40759	0.99693	-1.412	0.161
dat\$year2001	0.28181	0.83836	0.336	0.738
dat\$year2002	0.48350	0.85421	0.566	0.573
dat\$year2003	-0.52084	0.85383	-0.610	0.543
dat\$year2004	-0.14562	0.85154	-0.171	0.865
dat\$year2005	-0.18291	0.92692	-0.197	0.844
dat\$year2006	0.02815	0.88439	0.032	0.975
dat\$year2007	0.83641	0.86104	0.971	0.334
dat\$year2008	0.13978	0.85457	0.164	0.870
dat\$year2009	0.55629	0.84928	0.655	0.514
dat\$year2010	-0.10216	0.86124	-0.119	0.906
dat\$year2011	0.45982	0.86863	0.529	0.598
dat\$year2012	0.83088	0.86131	0.965	0.337
dat\$year2013	0.32462	0.89425	0.363	0.717
dat\$year2014	0.74916	0.87838	0.853	0.396
dat\$year2015	1.15411	0.88386	1.306	0.195

18 Puerto Rico tournament index

```

dat$year2016 -0.87199    0.84836   -1.028    0.307
dat$year2017 -0.09821    0.85408   -0.115    0.909
dat$year2018  0.56177    1.16405    0.483    0.630
dat$year2019  0.29081    0.87708    0.332    0.741
dat$year2020 -0.66680    0.91493   -0.729    0.468
dat$year2022  0.78828    0.91415    0.862    0.391
dat$mon1      -0.31971    0.39283   -0.814    0.418
dat$mon2      -0.06351    0.38114   -0.167    0.868
dat$mon3      -0.12341    0.40595   -0.304    0.762
dat$zone1     0.12877    0.47137    0.273    0.785
dat$zone2     0.68527    0.49968    1.371    0.174
dat$zone3     0.38090    0.55850    0.682    0.497
dat$zone4    -0.12452    0.47920   -0.260    0.796

```

(Dispersion parameter for gaussian family taken to be 0.6226789)

```

Null deviance: 103.767  on 122  degrees of freedom
Residual deviance:  58.532  on  94  degrees of freedom
AIC: 317.72

```

Number of Fisher Scoring iterations: 2

```
anova(out)
```

Analysis of Deviance Table

Model: gaussian, link: identity

Response: log(dat\$cpue)

Terms added sequentially (first to last)

	Df	Deviance	Resid. Df	Resid. Dev	F	Pr(>F)
NULL			122	103.767		
dat\$year	21	34.426	101	69.341	2.6327	0.0007729 ***
dat\$mon	3	2.382	98	66.959	1.2752	0.2874289
dat\$zone	4	8.427	94	58.532	3.3835	0.0124215 *

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1


18 Puerto Rico tournament index

```
# estimated coefficients from standardization
ind <- emmeans(out, "year", type = "response") %>% as.data.frame()
names(ind)[1] <- "Year"
```

Finally, we compare the standardized CPUE with the nominal CPUE. The two time series are somewhat closely related, indicating similar years of low and high relative abundance.

```
# look at nominal vs standardized index -----
plot(yrs, ind$response, type = "l", lwd = 2, col = 4, main = "Nominal versus standardized CPUE",
      xlab = "year", ylab = "CPUE (fish / (boats x time))", ylim = c(0, 1.8))
points(yrs, ind$response, pch = 1, lwd = 2, col = 4)
lines(yrs, ind$lower.CL, col = 4, lty = 2)
lines(yrs, ind$upper.CL, col = 4, lty = 2)
lines(yrs, tapply(dat$cpue, dat$year, mean, na.rm = T) * 1.0, col = 2, lwd = 2)
points(yrs, tapply(dat$cpue, dat$year, mean, na.rm = T) * 1.0, col = 2, lwd = 2, pch = 1)
legend("topleft", c("nominal", "standardized"), lwd = 2, pch = 19, col = c(2, 4), bty = "n")
```

Nominal versus standardized CPUE from dolphin tournament



18 Puerto Rico tournament index

```
cor(ind$response, tapply(dat$cpue, dat$year, mean, na.rm = T))
```

```
[1] 0.9068784
```

Finally, let's look at how our standardized index compares to the South Atlantic landings. We experimented with CPUE in both units of abundance and weight, and for different combinations of winter months (e.g., Dec - Feb, Jan - Apr). The highest correlation with South Atlantic landings occurs with tournament data from December to March with CPUE based on abundance. In this case the tournament data describes 18% of the variation in South Atlantic landings, with 2015 as a well above-average outlier in both the tournament CPUE and the landings.

```
rec <- read.csv("data/recLandings.csv")
names(rec)[1] <- "Year"
rec$ATL <- rowSums(rec[3:6], na.rm = T)

d <- merge(rec, ind, by = "Year")

par(mar = c(4, 6, 2, 1))
plot(d$response, d$ATL/10^6, col = 0, xlab = "standardized tournament CPUE",
     ylab = "total U.S. Atlantic coast recreational landings\n(millions of pounds)")
text(d$response, d$ATL/10^6, d$Year, col = 1)
out <- lm(d$ATL/10^6 ~ d$response)
abline(out, col = 8)
summary(out)
```

Call:

```
lm(formula = d$ATL/10^6 ~ d$response)
```

Residuals:

Min	1Q	Median	3Q	Max
-8.2004	-2.2964	-0.8387	2.2794	8.8767

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	12.535	2.066	6.067	6.26e-06 ***
d\$response	8.131	4.941	1.646	0.115

18 Puerto Rico tournament index

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1


Residual standard error: 4.195 on 20 degrees of freedom

Multiple R-squared: 0.1193, Adjusted R-squared: 0.07524

F-statistic: 2.709 on 1 and 20 DF, p-value: 0.1154

```
r2 <- summary(out)$adj.r.squared
p_val <- summary(out)$coefficients[2, 4]
p_display <- ifelse(p_val < 0.001, "p < 0.001", paste("p =", round(p_val, 3)))
legend("topleft",
      legend = bquote(R2 == .(round(r2, 2)) ~ "; " ~ p == .(round(p_val, 3))),
      bty = "n", cex = 1.2, text.col = 4)
```

18 Puerto Rico tournament index



```
# output the index
write.csv(ind, file = "indices/PRtournament_index.csv", row.names = F)
```

19 Wintertime pelagic longline index

A substantial portion of the U.S. pelagic longline fishery operates in the Caribbean region, and we can analyze the catch rates in this region as a potential predictive index of dolphin abundance in the South Atlantic later in the year. The logbook data are confidential and cannot be shown here, but we show the results of the standardization process. The code for processing the pelagic longline logbook data can be seen in the file “PLL_index.R” and the steps are outlined below.

19.1 Analysis of logbook data

The goal of the analysis is to create a standardized index of relative abundance for dolphin, using wintertime catch rates from the pelagic longline (PLL) fleet. The standardization process removes the potential influence of nuisance factors from the catch data, such as species targeting or differences in gear deployment, so that the index represents the underlying abundance of the fish.

The steps are as follows:

1. Read in the pelagic longline data which contains set level entries including location of fishing, target species, gear used, other details of the fishing trip and number of dolphin caught, discarded alive, discarded dead, and the total pounds of dolphin kept.
2. Format the longitude and latitude of the sets into decimal degrees, and find the points that fall in the Caribbean region (we use 76°W - 60°W and 12°N - 23°N as boundaries). Check that subsetting was done correctly, and extract the Caribbean data from the data set.
3. Inspect the dolphin catch columns (number of dolphin kept, discarded alive and dead). Replace NA values with zeros. Calculate the total dolphin catch as the total kept and discarded. Remove data for which the number of hooks is not reported. Calculate the catch rate as total dolphin caught divided by 1,000 hooks.

4. Standardize date formatting and add identifiers for month. Index December so that it is included with the winter months of the following year. Subset the data set so that it includes only the winter months (December - February).
5. Explore the different factors to include in the standardization model, relying on previous literature and best practices for standardizing PLL logbook data. Analyze the number of observations for different gear configurations and target species. Factors considered were: temperature, month, number of hooks between floats, bait type, number of lights between hooks, and species targeted. Temperature was binned in 5-degree increments. Month and hooks between floats were converted to factors. There was little resolution in the data for bait type and number of lights per hook, with similar values across all observations. The only major reported species targeted were dolphin, swordfish and mixed; targeting for other species were only listed in a small number of sets.
6. Use the delta-lognormal method to create an index of abundance for dolphin, estimated using a generalized linear model approach. The delta model fits separately the: 1) proportion of positive sets, assuming a binomial error distribution, and 2) the mean catch rate of sets where at least one dolphin was caught, assuming a lognormal error distribution. Use a step-wise regression process was used to determine the set of factors that significantly explain the observed variability. Model selection of fixed factors was based on significance with $\alpha=0.05$. Once the final model is determined, calculate the least-squares means from the linear models for the year predictor estimates. The index of abundance is the standardized proportion positive by year multiplied by the standardized abundance when present.
7. Calculate the deviance explained by each model, the standard errors, and the combined variance for the standardized index.

19.2 Model results

Here we upload the model results and show the standardized index.

```
load("data/linear_model_outputs.RData")  
  
# Print results  
cat("Deviance Explained (Binomial): ", round(d2_bin * 100, 2), "%\n")
```

```
Deviance Explained (Binomial): 7.75 %
```

19 Wintertime pelagic longline index

```
cat("Deviance Explained (Lognormal):", round(d2_log * 100, 2), "%\n")
```

Deviance Explained (Lognormal): 17.28 %


```
cat("Total Delta-Lognormal D^2: ", round(total_d2 * 100, 2), "%\n")
```

Total Delta-Lognormal D²: 9.63 %

The deviance explained by the model is relatively low, and thus the standardized index is similar to the nominal CPUE index. The primary impact of standardization is to lower slightly the index of relative abundance in the earlier years of the time series.

```
ind <- read.csv("indices/PLL_index.csv")

plot(ind$Year, ind$index, type = "l", lwd = 2, col = 4, main = "Nominal versus standardized
      xlab = "year", ylab = "CPUE (dolphin / 1000 hooks)", ylim = c(0, 14))
points(ind$Year, ind$index, pch = 1, lwd = 2, col = 4)
lines(ind$Year, ind$index - 1.96*ind$SE, col = 4, lty = 2)
lines(ind$Year, ind$index + 1.96*ind$SE, col = 4, lty = 2)
lines(ind$Year, ind$nominal, col = 2, lwd = 2)
points(ind$Year, ind$nominal, col = 2, lwd = 2, pch = 1)
legend("topleft", c("nominal CPUE", "standardized CPUE"), lwd = 2, pch = 19, col = c(2, 4),
```



Now we compare the nominal and the standardized index with the Atlantic recreational landings. We experimented with altering the forecasting time period (e.g., Dec - Feb versus Jan - Mar) and also with the latitudinal boundaries (e.g., 70°W versus 76°W).

19 Wintertime pelagic longline index

A smaller bounding box of 70°W resulted in very small sample sizes (<10) for some individual years; extending the bounding box to 76°W to cover fishing activity between Cuba and Hispanola resulted in much higher sample sizes and improved correlations. Including only the winter months (Dec - Feb) also resulted in higher correlations than when spring months were included.

```
rec <- read.csv("data/recLandings.csv")
names(rec)[1] <- "Year"
rec$ATL <- rowSums(rec[3:6], na.rm = T)
rec <- rec[which(rec$Year >= 1990), ]

d1 <- merge(rec, ind, by = "Year")


lis <- c("index", "nominal", "predpos", "Npres")
labs <- c("standardized CPUE index", "nominal CPUE",
          "standardized probability of occurrence",
          "standardized abundance when present")

par(mfrow = c(2, 2), mar = c(4, 5, 2, 1), mgp = c(2.5, 1, 0))

for (i in 1:4) {
  j <- which(names(d1) == lis[i])
  plot(d1[, j], d1$ATL/10^6, col = 0, xlab = labs[i], ylim = c(5, 32), las = 1,
       ylab = "total U.S. Atlantic coast recreational landings\n(millions of pounds)")
  text(d1[, j], d1$ATL/10^6, substr(d1$Year, 1, 4), col = 1)
  out <- lm(d1$ATL/10^6 ~ d1[, j ])
  abline(out, col = 8)
  summary(out)

  r2 <- summary(out)$adj.r.squared
  p_val <- summary(out)$coefficients[2, 4]
  p_display <- ifelse(p_val < 0.001, "p < 0.001", paste("p =", round(p_val, 3)))
  legend("bottomright",
        legend = bquote(R^2 == .(round(r2, 2)) ~ "; " ~ p == .(round(p_val, 3))),
        bty = "n", cex = 1.2, text.col = 4)
}
```

19 Wintertime pelagic longline index



Interestingly, the nominal CPUE is more highly correlated with Atlantic recreational landings than the standardized index. However when the standardized index is broken into its two components, we can see that the probability of occurrence is not significantly correlated with the Atlantic landings, while the abundance when present has a highly significant correlation and explains 44% of the variation in the Atlantic landings.