Table 1 Clinicopathological features of study participants.

Characteristic	Non-cancer (n=400)	PRC (n=100)	OVC (n=70)	LC (n=300)	PAC (n=70)	CRC (n=300)	p-Value
Age (yr), median [IQR]	58.0 [49.0, 65.0]	72.5 [67.0, 82.0]	56.5 [47.2, 62.8]	66.0 [60.0, 72.0]	68.0 [63.2, 73.8]	67.0 [59.8, 76.0]	**<0.001**
Sex, n (%)							**<0.001**
Female	102 (25.5)	0 (0.0)	70 (100.0)	138 (46.0)	36 (51.4)	107 (35.7)	
Male	298 (74.5)	100 (100.0)	0 (0.0)	162 (54.0)	34 (48.6)	193 (64.3)	
BMI (kg/m²), median [IQR]	24.6 [22.6, 26.9]	24.4 [22.0, 26.1]	23.5 [21.1, 26.7]	23.6 [21.7, 25.4]	21.4 [18.7, 24.3]	24.1 [21.5, 26.4]	**<0.001**
Smoking status, n (%)							
Never	158 (40.9)	49 (51.0)	29 (87.9)	87 (64.9)	—	—	
Former	141 (36.5)	34 (35.4)	0 (0.0)	10 (7.5)	—	—	
Current	87 (22.5)	13 (13.5)	4 (12.1)	37 (27.6)	—	—	
Unknown	14 (3.5)	4 (4.0)	37 (52.9)	166 (55.3)	70 (100.0)	300 (100.0)	
Drinking status, n (%)							
Never	143 (37.1)	74 (77.9)	27 (90.0)	84 (63.2)	—	—	
Former	0 (0.0)	1 (1.1)	0 (0.0)	0 (0.0)	—	—	
Current	242 (62.9)	20 (21.1)	3 (10.0)	49 (36.8)	—	—	
Unknown	15 (3.8)	5 (5.0)	40 (57.1)	167 (55.7)	70 (100.0)	300 (100.0)	
Stage, n (%)							
Early (I–II)	—	24 (24.0)	8 (11.4)	228 (76.0)	39 (55.7)	253 (84.3)	
Advanced (III–IV)	—	51 (51.0)	5 (7.1)	27 (9.0)	29 (41.4)	47 (15.7)	
Unknown	—	25 (25.0)	57 (81.4)	45 (15.0)	2 (2.9)	0	

Abbreviations: BMI, body mass index; IQR, interquartile range; yr, years.
Non-cancer includes: Normal control (n=100), DM (diabetes, n=100), HTN (hypertension, n=100), DM + HTN (n=100).
CRC, colorectal cancer; LC, lung cancer; OVC, ovarian cancer; PAC, pancreatic cancer; PRC, prostate cancer.
Values are presented as median [IQR] or n (%). p-Values from Kruskal–Wallis test (continuous) or χ^2 test (categorical).
Stage: Early (I–II), Advanced (III–IV). Based on AJCC Cancer Staging Manual, 8th Edition.