

Table 1. Baseline Demographics and Disease Stage of Study Participants (N = 1,629)

| Group           | Cancer type | <i>n</i> | Age (yr), median [IQR] | Sex (F:M), n (%) | BMI (kg/m <sup>2</sup> ), median [IQR] | Stage (I, II), n (%) | Stage (III, IV), n (%) | Unknown, n (%) |
|-----------------|-------------|----------|------------------------|------------------|----------------------------------------|----------------------|------------------------|----------------|
| Non-cancer      | —           | 429      | 57 [49–66]             | 120:309 (28/72)  | 25 [23–27]                             | —                    | —                      | —              |
| PRC             | Prostate    | 100      | 72 [67–82]             | 0:100 (0/100)    | 24 [22–26]                             | 26 (26)              | 49 (49)                | 25 (25)        |
| OVC             | Ovarian     | 70       | 56 [47–63]             | 70:0 (100/0)     | 24 [21–27]                             | 8 (11)               | 4 (6)                  | 58 (83)        |
| LC              | Lung        | 300      | 66 [60–72]             | 138:162 (46/54)  | 24 [22–25]                             | 228 (76)             | 27 (9)                 | 45 (15)        |
| PAC             | Pancreatic  | 100      | 68 [61–73]             | 50:50 (50/50)    | 22 [19–24]                             | 37 (37)              | 54 (54)                | 9 (9)          |
| CRC             | Colorectal  | 300      | 67 [60–76]             | 107:193 (36/64)  | 24 [22–26]                             | 132 (44)             | 143 (48)               | 25 (8)         |
| BLC             | Bladder     | 299      | 72 [64–80]             | 45:254 (15/85)   | 24 [22–27]                             | 291 (97)             | 8 (3)                  | 0              |
| BRE             | Breast      | 30       | 62 [51–72]             | 30:0 (100/0)     | 25 [22–29]                             | 23 (77)              | 0                      | 7 (23)         |
| <i>p</i> -value |             |          | <0.001                 | <0.001           | <0.001                                 |                      |                        |                |

Continuous variables are presented as median [interquartile range]. Categorical variables are presented as n (%). Stage is classified as early (I, II) vs. advanced (III, IV). Non-cancer includes NOR (n = 100), DIA (n = 100), HBP (n = 100), H.D. (n = 100), and YNOR (n = 29). PAC includes CPAN (n = 70) and YPAN (n = 30). *p*-values were calculated using the Kruskal–Wallis test for continuous variables and Chi-square test for categorical variables.