Table 1 Clinicopathological features of study participants.

Characteristic	Non-cancer				Cancer					p-Value
	NOR (n=100)	DIA (n=100)	HBP (n=100)	H.D. (n=100)	PRC (n=100)	OVC (n=70)	LC (n=300)	PAC (n=70)	CRC (n=300)	
Age (yr), median [IQR]	44.5 [36.8, 55.2]	60.0 [52.8, 66.0]	59.5 [54.0, 67.0]	62.5 [54.0, 70.2]	72.5 [67.0, 82.0]	56.5 [47.2, 62.8]	66.0 [60.0, 72.0]	68.0 [63.2, 73.8]	67.0 [59.8, 76.0]	**<0.001**
Sex, n (%)										**<0.001**
Female	23 (23.0)	28 (28.0)	27 (27.0)	24 (24.0)	0 (0.0)	70 (100.0)	138 (46.0)	36 (51.4)	107 (35.7)	
Male	77 (77.0)	72 (72.0)	73 (73.0)	76 (76.0)	100 (100.0)	0 (0.0)	162 (54.0)	34 (48.6)	193 (64.3)	
BMI (kg/m²), median [IQR]	24.2 [22.4, 25.6]	24.2 [22.2, 26.4]	25.4 [22.9, 27.5]	25.1 [23.1, 27.0]	24.4 [22.0, 26.1]	23.5 [21.1, 26.7]	23.6 [21.7, 25.4]	21.4 [18.7, 24.3]	24.1 [21.5, 26.4]	**<0.001**
Smoking status, n (%)										
Never	42 (42.0)	37 (39.4)	41 (42.3)	38 (40.0)	49 (51.0)	29 (87.9)	0 (0.0)	—	—	
Former	31 (31.0)	36 (38.3)	40 (41.2)	34 (35.8)	34 (35.4)	0 (0.0)	10 (29.4)	—	—	
Current	27 (27.0)	21 (22.3)	16 (16.5)	23 (24.2)	13 (13.5)	4 (12.1)	24 (70.6)	—	—	
Drinking status, n (%)										
Never	32 (32.0)	42 (45.2)	36 (37.5)	33 (34.4)	74 (77.9)	27 (90.0)	0 (0.0)	—	—	
Former	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.1)	0 (0.0)	0 (0.0)	—	—	
Current	68 (68.0)	51 (54.8)	60 (62.5)	63 (65.6)	20 (21.1)	3 (10.0)	33 (100.0)	—	—	
Stage, n (%)										
Early (I–II)	—	—	—	—	24 (24.0)	8 (11.4)	228 (76.0)	39 (55.7)	253 (84.3)	
Advanced (III–IV)	—	—	—	—	51 (51.0)	5 (7.1)	27 (9.0)	29 (41.4)	47 (15.7)	
Unknown	—	—	—	—	25 (25.0)	57 (81.4)	45 (15.0)	2 (2.9)	0	

Abbreviations: BMI, body mass index; DIA, diabetes; H.D., hypertension + diabetes; HBP, hypertension; IQR, interquartile range; yr, years.
Values are presented as median [IQR] or n (%). p-Values from Kruskal–Wallis test (continuous) or χ^2 test (categorical).
Stage: Early (I–II), Advanced (III–IV). Based on AJCC Cancer Staging Manual, 8th Edition.