

TinyBolus Formulary

Peer-review document

Version 0.0.39 · Updated 2026-05-19

Published by Aether Somnus

Developed by Sérgio Pinto, M.D. and Patrícia Lima, M.D.

Contact: tinybolusfeedback@gmail.com

Source: github.com/SGP-Prime/tinybolusformulas

Privacy policy: sgp-prime.github.io/tinybolus-site/privacy.html

This is a reference tool, not a prescribing system. Verify every value against your hospital protocols, product labelling, and the individual patient in front of you before administration.

About this document

This is the human-readable export of the TinyBolos paediatric reference formulary, organised section-by-section as it appears in the application. Each item is presented with its dose specification, applicable age brackets (where present), route, clinical notes, and numerical references pointing into the bibliography at the end of the document.

How to read each item

Items fall into a small number of shapes that map directly onto how the application computes them:

- **Single dose** (e.g., 10 µg/kg): one factor multiplied by weight, often with a hard maximum cap.
- **Dose range** (e.g., 2–3 mg/kg): a low–high range, also weight-scaled, optionally with a maximum cap on either bound.
- **Age-bracketed**: different dose factors for different age cohorts. Rendered as a small table per item.
- **Age-bracketed fixed value**: not weight-scaled, just a string per age band (e.g., laryngoscope blade size).
- **Computed**: piecewise logic that doesn't reduce to a single formula (ETT sizing, fluid maintenance, LMA / supraglottic airway sizing, vital-sign normal ranges).

Hard floors / contraindications

Some items carry an *age* or *weight* floor below which the formulary explicitly declines to suggest a dose. In the app this renders as ∅ (empty-set glyph) with a "not indicated" explanation. Items with such floors are flagged in red below each row.

References

Every dose / range / age bracket / sizing rule carries one or more citations. References are numbered in encounter order throughout this document, with a full bibliography at the end. The numbers in [brackets] after each item refer to entries in that bibliography.

Source

The formulary itself lives at github.com/SGP-Prime/tinybolusformulas as a single machine-readable JSON file. This document is regenerated from that file on demand by `tool/generate_formulary_review_html.dart`; do not hand-edit it.

Contents

1. EMERGENCY — PALS
2. Vital Signs
3. Induction
4. Painkillers
5. Emergency Drugs
6. Fluids and Blood
7. Equipment
8. Respiratory
9. Reversal Agents
10. Regional Anesthesia
11. Weight Estimation
12. Bibliography

EMERGENCY — PALS

Compression rate

VALUE 100–120/min

References: [\[1\]](#)

Compression depth

RULE 4 cm <1 y · 5 cm ≥1 y

not exceeding 6 cm at any age

References: [\[1\]](#)

C:V ratio (2 rescuers)

VALUE 15:2

1 rescuer: 30:2

References: [\[1\]](#)

Defibrillation

DOSE 4 J/kg

escalate to 8 J/kg (max 360 J) for refractory VF/pVT after >5 shocks

References: [\[1\]](#), [\[2\]](#)

Cardioversion (synchronised)

DOSE 1 J/kg

double per attempt to max 4 J/kg

References: [\[1\]](#)

Adrenaline (cardiac arrest)

IV/IO

DOSE 10 µg/kg (max 1 mg)

References: [\[1\]](#), [\[3\]](#)

Amiodarone

IV

DOSE 5 mg/kg (max 300 mg)

max 150 mg for subsequent doses

References: [\[1\]](#)

Atropine

IV

DOSE 20 µg/kg (max 500 µg)

historical 100 µg minimum is a myth — no floor required

References: [\[1\]](#), [\[4\]](#)

Adenosine (1st dose)

IV

DOSE 0.1 mg/kg (max 6 mg)

References: [\[1\]](#)

Adenosine (2nd dose)

IV

DOSE 0.3 mg/kg (max 18 mg)

References: [\[1\]](#)

Vital Signs

Heart Rate

≤1 mo	120–170 bpm
2–12 mo	100–160 bpm
1–2 y	98–140 bpm
3–5 y	80–120 bpm
6–11 y	75–118 bpm
≥12 y	60–100 bpm

References: [\[5\]](#)

Systolic BP

≤12 mo	72–104 mmHg
1–2 y	86–106 mmHg
3–5 y	89–112 mmHg
6–9 y	97–115 mmHg
10–11 y	102–120 mmHg
12–15 y	110–131 mmHg
≥16 y	90–140 mmHg

References: [\[5\]](#)

Diastolic BP

≤12 mo	37–56 mmHg
1–2 y	42–63 mmHg
3–5 y	46–72 mmHg
6–9 y	57–76 mmHg
10–11 y	61–80 mmHg
12–15 y	64–83 mmHg
≥16 y	60–90 mmHg

References: [\[5\]](#)

Respiratory Rate

≤1 mo	40–60 /min
2–12 mo	30–53 /min
1–3 y	22–37 /min
4–5 y	20–28 /min
6–12 y	18–25 /min
≥13 y	12–20 /min

References: [\[5\]](#)

Tidal Volume

DOSE 5–7 mL/kg

intraoperative ventilation, healthy lungs · use lower end (5 mL/kg) for neonates and very young infants

References: [\[6\]](#), [\[7\]](#)

Induction

Propofol

IV

≤1 mo	1–2.5 mg/kg
2–12 mo	3–4 mg/kg
1–6 y	3–3.5 mg/kg
7–12 y	2.5–3 mg/kg
≥13 y	2–2.5 mg/kg

References: [\[8\]](#), [\[9\]](#), [\[10\]](#)

Ketamine

IV

DOSE 1–2 mg/kg

References: [\[11\]](#), [\[12\]](#)

Rocuronium

IV

DOSE 0.6–1.2 mg/kg

intubation 0.6 · RSI 1.0–1.2

References: [\[13\]](#), [\[14\]](#)

Fentanyl (induction)

IV

DOSE 1–3 µg/kg

References: [\[12\]](#)

Midazolam

IV

DOSE 0.1 mg/kg

References: [\[12\]](#)

Midazolam (oral premed)

PO

DOSE 0.5 mg/kg (max 20 mg)

References: [\[12\]](#)

Ondansetron

IV

DOSE 0.1 mg/kg (max 4 mg)

References: [\[15\]](#), [\[12\]](#)

Dexamethasone (PONV)

IV

DOSE 0.15 mg/kg

References: [\[15\]](#), [\[12\]](#)

Painkillers

Paracetamol

IV

≤1 mo 7.5 mg/kg

≥0 y 15 mg/kg

CAP max 1 g

References: [\[16\]](#), [\[12\]](#)

Ketorolac

IV

DOSE 0.5 mg/kg (max 30 mg)

Contraindicated <6 months

References: [\[17\]](#), [\[12\]](#)

Morphine

IV

≤1 mo 0.03–0.05 mg/kg

≥0 y 0.1 mg/kg

References: [\[18\]](#), [\[12\]](#)

Fentanyl (analgesia)

IV

DOSE 0.5–1 µg/kg

References: [\[12\]](#), [\[19\]](#)

Metamizole

IV

DOSE 8–16 mg/kg (max 1 g)

Contraindicated <3 months

References: [\[20\]](#), [\[21\]](#)

Ibuprofen (IV)

IV

DOSE 10 mg/kg (max 400 mg)

References: [\[22\]](#), [\[12\]](#)

Emergency Drugs

Adrenaline (cardiac arrest)

IV/IO

DOSE 10 µg/kg (max 1 mg)

References: [\[1\]](#), [\[3\]](#)

Adrenaline (anaphylaxis IM)

IM

DOSE 10 µg/kg (max 500 µg)

fixed-dose convention: <6 y 150 µg · 6–12 y 300 µg · >12 y 500 µg (EAACI)

References: [\[23\]](#)

Adrenaline (anaphylaxis IV)

IV

DOSE 1 µg/kg

References: [\[1\]](#), [\[23\]](#)

Atropine

IV

DOSE 20 µg/kg (max 500 µg)

max 1 mg single dose in adolescents · cumulative max 2 mg

References: [\[1\]](#), [\[4\]](#)

Ephedrine

IV

DOSE 0.1–0.3 mg/kg

References: [\[24\]](#), [\[25\]](#)

Phenylephrine

IV

DOSE 5–20 µg/kg

References: [\[12\]](#), [\[26\]](#)

Lidocaine

IV

DOSE 1 mg/kg

References: [\[1\]](#)

Dantrolene

IV

DOSE 2.5 mg/kg

repeat 5–10 min · max 10 mg/kg

References: [\[27\]](#), [\[28\]](#)

Fluids and Blood

Fluid Bolus

IV

DOSE 10 mL/kg

References: [\[1\]](#)

Maintenance

RULE Holliday-Segar 4-2-1: 4 mL/kg/h (first 10 kg) · 2 mL/kg/h (next 10 kg) · 1 mL/kg/h thereafter

4-2-1 rule (Holliday-Segar)

References: [\[29\]](#), [\[30\]](#)

Total Blood Volume

≤1 mo 85 mL/kg

2 mo – 2 y 75 mL/kg

≥3 y 70 mL/kg

References: [\[31\]](#), [\[5\]](#)

Tranexamic Acid

IV

DOSE 15–20 mg/kg

References: [\[32\]](#), [\[33\]](#)

Equipment

ETT

RULE Cuffed: $3.5 + \text{age}/4$, rounded to ± 0.5

≤ 3 KG Neonatal weight table:

≤ 1 kg size 2.5

≤ 2 kg size 3.0

≤ 4 kg size 3.5

< 1 Y size 4.0

for uncuffed tube, add 0.5 to size

References: [\[34\]](#)

ETT insertion depth

RULE $6 + \text{weight} (< 1 \text{ mo}) \cdot \text{weight}/2 + 8 (< 1 \text{ y}) \cdot \text{age}/2 + 12 (\geq 1 \text{ y})$ (cm)

nasal: add 2–3 cm

References: [\[35\]](#), [\[36\]](#)

iGel LMA

IGEL SIZING weight bands (inclusive on both bounds):

2–5 kg size 1

5–12 kg size 1.5

10–25 kg size 2

25–35 kg size 2.5

30–60 kg size 3

50–90 kg size 4

90– ∞ kg size 5

References: [\[37\]](#), [\[38\]](#)

AuraGain LMA

AURAGAIN SIZING weight bands (inclusive on both bounds):

2–5 kg	size 1
5–10 kg	size 1.5
10–20 kg	size 2
20–30 kg	size 2.5
30–50 kg	size 3
50–70 kg	size 4
70–100 kg	size 5
100–∞ kg	size 6

References: [\[39\]](#)

Laryngoscope blade

≤1 y	Macintosh 1
2–11 y	Macintosh 2
≥12 y	Macintosh 3

References: [\[40\]](#), [\[5\]](#)

Face mask

≤1 mo	Size 0
2–12 mo	Size 1
1–3 y	Size 2
4–8 y	Size 3
9–14 y	Size 4
≥15 y	Size 5

mask should cover nose and mouth from eyebrow line to chin cleft · anatomical fit is more important than estimated size

References: [\[5\]](#), [\[41\]](#)

Respiratory

Salbutamol (inhaled)

INHALED

≤5 y 2–10 puffs

≥6 y 4–10 puffs

100 µg/puff · repeat incrementally as needed

References: [\[42\]](#), [\[43\]](#)

Salbutamol (nebulized)

NEBULIZED

≤5 y 2.5 mg

≥6 y 5 mg

repeat back-to-back or continuously per response

References: [\[42\]](#), [\[43\]](#)

Ipratropium (inhaled)

INHALED

≤5 y 4 puffs

≥6 y 4–8 puffs

20 µg/puff · repeat as needed

References: [\[44\]](#), [\[42\]](#)

Ipratropium (nebulized)

NEBULIZED

≤11 y 250 µg

≥12 y 500 µg

References: [\[12\]](#)

Magnesium Sulphate

IV

DOSE 25–50 mg/kg (max 2 g)

References: [\[45\]](#), [\[12\]](#)

Clemastine

IV

DOSE 25 µg/kg (max 2 mg)

References: [\[46\]](#), [\[12\]](#)

Adrenaline (nebulized)

NEBULIZED

DOSE 0.5 mg/kg (max 5 mg)

References: [\[47\]](#), [\[12\]](#)

Succinylcholine (RSI)

IV

DOSE 1–2 mg/kg

intubating dose · sub-paralytic dose for laryngospasm is below

References: [\[14\]](#), [\[5\]](#)

Succinylcholine (laryngospasm)

IV

DOSE 0.1–0.5 mg/kg

References: [\[48\]](#)

Succinylcholine (IM)

IM

DOSE 4 mg/kg

for laryngospasm or RSI without IV access

References: [\[48\]](#)

Rocuronium (laryngospasm)

IV

DOSE 0.1–0.2 mg/kg

References: [\[49\]](#)

Reversal Agents

Naloxone (titrated)

IV

DOSE 0.01 mg/kg

post-anaesthesia reversal — titrate to respiration · repeat every 2–3 min

References: [\[12\]](#), [\[1\]](#)

Naloxone (overdose)

IV/IO/IM

DOSE 0.1 mg/kg (max 2 mg)

suspected opioid overdose with apnoea or unresponsiveness

References: [\[1\]](#), [\[12\]](#)

Flumazenil

IV

DOSE 10 µg/kg (max 200 µg)

max bolus 200 µg · total max 1 mg

References: [\[12\]](#)

Sugammadex

IV

DOSE 2–4 mg/kg

2 mg/kg moderate block · 4 mg/kg deep block

References: [\[50\]](#), [\[51\]](#)

Sugammadex (rescue, profound block)

IV

DOSE 16 mg/kg

for CICO / immediate reversal of full intubating-dose rocuronium

References: [\[52\]](#), [\[53\]](#)

Regional Anesthesia

Lidocaine (perineural)

PERINEURAL

≤5 mo maximum 3.5 mg/kg

≥0 y maximum 5 mg/kg

References: [\[54\]](#), [\[12\]](#)

Bupivacaine (perineural)

PERINEURAL

≤5 mo maximum 1.5–2 mg/kg — *infusion: 0.2 mg/kg/h*

≥0 y maximum 2–2.5 mg/kg — *infusion: 0.4 mg/kg/h*

References: [\[54\]](#), [\[55\]](#)

Ropivacaine (perineural)

PERINEURAL

≤5 mo maximum 2 mg/kg — *infusion: 0.2 mg/kg/h*

≥0 y maximum 3 mg/kg — *infusion: 0.4 mg/kg/h*

References: [\[56\]](#), [\[57\]](#)

Intralipid 20% bolus

IV

DOSE 1.5 mL/kg

max total 12 mL/kg

References: [\[54\]](#), [\[58\]](#)

Intralipid 20% infusion

IV

DOSE 0.25 mL/min/kg

References: [\[54\]](#), [\[58\]](#)

Weight Estimation

Used when the user does not supply a measured weight. <1 month and >12 years always require user input.

1–12 mo (infant)

FORMULA $\text{weight (kg)} = (0.5 \times \text{months}) + 4$

References: [\[59\]](#)

1–5 y (young child)

FORMULA $\text{weight (kg)} = (2 \times \text{age}) + 8$

References: [\[60\]](#), [\[59\]](#)

6–12 y (school-age)

FORMULA $\text{weight (kg)} = (3 \times \text{age}) + 7$

References: [\[61\]](#), [\[59\]](#)

Bibliography

Citations numbered in order of first encounter throughout this document.

1. Djakow J, Turner NM, Skellett S, Buysse CMP, Cardona F, de Lucas N, del Castillo J, Kiviranta P, Lauridsen KG, Markel F, Martinez-Mejias A, Roggen I, Biarent D. European Resuscitation Council Guidelines 2025: Paediatric Life Support. *Resuscitation* 2025;215(Suppl 1):110767.
2. Acworth J et al. ILCOR 2025 systematic review on pediatric defibrillation energy. *Resuscitation Plus* 2025;24:100991.
3. Lasa JJ, Dhillon GS, Topjian AA, et al. Part 8: Pediatric Advanced Life Support — 2025 American Heart Association and American Academy of Pediatrics Guidelines. *Circulation* 2025.
4. Barrington KJ. The myth of a minimum dose for atropine. *Pediatrics* 2011;127(4):783–784.
5. Coté CJ, Lerman J, Anderson BJ, eds. *A Practice of Anesthesia for Infants and Children*, 6th edition. Philadelphia: Elsevier; 2019.
6. Spaeth JP, Lam JE. Understanding pediatric ventilation in the operative setting. Part II: Setting perioperative ventilation. *Paediatr Anaesth* 2022;32(2):247–254.
7. Kneyber MCJ, de Luca D, Calderini E, et al. Recommendations for mechanical ventilation of critically ill children from the Paediatric Mechanical Ventilation Consensus Conference (PEMVECC). *Intensive Care Med* 2017;43(12):1764–1780.
8. Aun CST, Short TG, Leung DHY, Oh TE. Induction dose-response of propofol in unpremedicated children. *Br J Anaesth* 1992;68(1):64–67.
9. Westrin P. The induction dose of propofol in infants 1–6 months of age and in children 10–16 years of age. *Anesthesiology* 1991;74(3):455–458.
10. van Dijk H, Hendriks MP, van Eck-Smaling MM, van Wolfswinkel L, de Graaff JC. Age-stratified propofol dosage for pediatric procedural sedation and analgesia. *Anesth Analg* 2023;136(3):551–558.
11. Craven R. Ketamine. *Anaesthesia* 2007;62(Suppl 1):48–53.
12. Paediatric Formulary Committee. *BNF for Children* (online). London: BMJ Group, Pharmaceutical Press, and RCPCH Publications.
13. Meakin GH, Meretoja OA, Motsch J, et al. A comparison of rocuronium and vecuronium during anesthesia induction in children aged 1 to 10 years. *Paediatr Anaesth* 1995;5(6):359–362.
14. Mazurek AJ, Rae B, Hann S, et al. Rocuronium versus succinylcholine: are they equally effective during rapid-sequence induction of anesthesia? *Anesth Analg* 1998;87(6):1259–1262.
15. Gan TJ, Belani KG, Bergese S, et al. Fourth Consensus Guidelines for the Management of Postoperative Nausea and Vomiting. *Anesth Analg* 2020;131(2):411–448.
16. Anderson BJ. Paracetamol (acetaminophen): mechanisms of action. *Paediatr Anaesth* 2008;18(10):915–921.
17. Gupta A, Daggett C, Ludwick J, Wells W, Lewis A. Ketorolac after congenital heart surgery: does it increase the risk of significant bleeding complications? *Paediatr Anaesth* 2005;15(2):139–142.
18. Bouwmeester NJ, Anderson BJ, Tibboel D, Holford NH. Developmental pharmacokinetics of morphine and its metabolites in neonates, infants and young children. *Br J Anaesth* 2004;92(2):208–217.
19. Association of Paediatric Anaesthetists of Great Britain and Ireland. *Good Practice in Postoperative and Procedural Pain Management*, 2nd edition. *Paediatr Anaesth* 2012;22(Suppl 1):1–79.

20. Ziesenitz VC, Rodieux F, Atkinson A, et al. Dose evaluation of intravenous metamizole (dipyrone) in infants and children: a prospective population pharmacokinetic study. *Eur J Clin Pharmacol* 2019;75(11):1491–1502.
21. Fieler M, Eich C, Becke K, et al. Metamizole for postoperative pain therapy in 1177 children: a prospective, multicentre, observational, postauthorisation safety study. *Eur J Anaesthesiol* 2015;32(12):839–843.
22. Southey ER, Soares-Weiser K, Kleijnen J. Systematic review and meta-analysis of the clinical safety and tolerability of ibuprofen compared with paracetamol in paediatric pain and fever. *Curr Med Res Opin* 2009;25(9):2207–2222.
23. Muraro A, Roberts G, Worm M, et al. Anaphylaxis: guidelines from the European Academy of Allergy and Clinical Immunology. *Allergy* 2014;69(8):1026–1045.
24. Lönnqvist PA. Management of arterial hypotension in newborns and infants. *Curr Opin Anaesthesiol* 2017;30(3):317–322.
25. Association of Paediatric Anaesthetists of Great Britain and Ireland. Guidelines on the Prevention and Management of Hypotension in Children Under Anaesthesia. APAGBI, 2022.
26. Tobias JD. Phenylephrine in pediatric anesthesia: applications and safety considerations. *Paediatr Anaesth* 2013;23(8):679–687.
27. Glahn KPE, Ellis FR, Halsall PJ, Müller CR, Snoeck MMJ, Urwyler A, Wappler F. Recognizing and managing a malignant hyperthermia crisis: guidelines from the European Malignant Hyperthermia Group. *Br J Anaesth* 2010;105(4):417–420.
28. Hopkins PM, Girard T, Dalay S, Jenkins B, Thacker A, Patteril M, McGrady E. Malignant hyperthermia 2020: Guideline from the Association of Anaesthetists. *Anaesthesia* 2021;76(5):655–664.
29. Holliday MA, Segar WE. The maintenance need for water in parenteral fluid therapy. *Pediatrics* 1957;19(5):823–832.
30. Feld LG, Neuspiel DR, Foster BA, et al. Clinical Practice Guideline: Maintenance Intravenous Fluids in Children. *Pediatrics* 2018;142(6):e20183083.
31. Nadler SB, Hidalgo JH, Bloch T. Prediction of blood volume in normal human adults. *Surgery* 1962;51(2):224–232.
32. Goobie SM, Meier PM, Pereira LM, et al. Efficacy of tranexamic acid in pediatric craniostomosis surgery: a double-blind, placebo-controlled trial. *Anesthesiology* 2011;114(4):862–871.
33. Faraoni D, Goobie SM. The efficacy of antifibrinolytic drugs in children undergoing noncardiac surgery: a systematic review of the literature. *Anesth Analg* 2014;118(3):628–636.
34. Duracher C, Schmautz E, Martinon C, et al. Evaluation of cuffed tracheal tube size predicted using the Khine formula in children. *Paediatr Anaesth* 2008;18(2):113–118.
35. Phipps LM, Thomas NJ, Gilmore RK, et al. Prospective assessment of guidelines for determining appropriate depth of endotracheal tube placement in children. *Pediatr Crit Care Med* 2005;6(5):519–522.
36. Lau N, Playfor SD, Rashid A, Dhanarass M. New formulae for predicting tracheal tube length. *Paediatr Anaesth* 2006;16(12):1238–1243.
37. Intersurgical Ltd. i-gel User Guide: Adult and Paediatric Sizes. Intersurgical, Wokingham, UK (current edition).
38. Beringer RM, Kelly F, Cook TM, et al. A cohort evaluation of the paediatric i-gel airway during anaesthesia in 120 children. *Anaesthesia* 2011;66(12):1121–1126.

39. Ambu A/S. Ambu AuraGain Single Use Laryngeal Mask — Instructions for Use. V7, August 2023.
40. Passi Y, Sathyamoorthy M, Lerman J, Heard C, Marino M. Comparison of the laryngoscopy views with the size 1 Miller and Macintosh laryngoscope blades lifting the epiglottis or the base of the tongue in infants and children <2 yr of age. *Br J Anaesth* 2014;113(5):869–874.
41. Rendell-Baker L, Soucek DH. New pediatric face masks and anaesthetic equipment. *Br Med J* 1962;1(5293):1690.
42. British Thoracic Society / Scottish Intercollegiate Guidelines Network. British guideline on the management of asthma (SIGN 158). Edinburgh: BTS/SIGN; 2019, updated 2024.
43. Global Initiative for Asthma. Global Strategy for Asthma Management and Prevention, 2024 update. <https://ginasthma.org/>
44. Griffiths B, Ducharme FM. Combined inhaled anticholinergics and short-acting beta2-agonists for initial treatment of acute asthma in children. *Cochrane Database Syst Rev* 2013;(8):CD000060.
45. Griffiths B, Kew KM. Intravenous magnesium sulfate for treating children with acute asthma in the emergency department. *Cochrane Database Syst Rev* 2016;4:CD011050.
46. Ring J, Beyer K, Biedermann T, et al. Guideline (S2k) on acute therapy and management of anaphylaxis: 2021 update. *Allergo J Int* 2021;30(1):1–25.
47. Bjornson C, Russell K, Vandermeer B, Klassen TP, Johnson DW. Nebulized epinephrine for croup in children. *Cochrane Database Syst Rev* 2013;(10):CD006619.
48. Larson CP Jr. Laryngospasm — the best treatment. *Anesthesiology* 1998;89(5):1293–1294.
49. Reynolds LM, Lau M, Brown R, Luks A, Fisher DM. Intramuscular rocuronium in infants and children: dose-ranging and tracheal intubating conditions. *Anesthesiology* 1996;85(2):231–239.
50. Plaud B, Meretoja O, Hofmockel R, et al. Reversal of rocuronium-induced neuromuscular blockade with sugammadex in pediatric and adult surgical patients. *Anesthesiology* 2009;110(2):284–294.
51. Lee JH, Kwak HJ, Lee JY, et al. Sugammadex for our little ones: a brief narrative review. *Anesth Pain Med (Seoul)* 2024;19(4):257–267.
52. Lee C, Jahr JS, Candiotti KA, et al. Reversal of profound neuromuscular block by sugammadex administered three minutes after rocuronium. *Anesthesiology* 2009;110(5):1020–1025.
53. Bridion (sugammadex) prescribing information. Merck Sharp & Dohme Corp., 2025.
54. Neal JM, Barrington MJ, Fettiplace MR, et al. The third American Society of Regional Anesthesia and Pain Medicine practice advisory on local anesthetic systemic toxicity: executive summary 2017. *Reg Anesth Pain Med* 2018;43(2):113–123.
55. Suresh S, Ecoffey C, Bosenberg A, et al. ESRA/ASRA Joint Committee practice advisory on controversial topics in pediatric regional anesthesia. *Reg Anesth Pain Med* 2018;43(2):211–216.
56. Lönnqvist PA, Westrin P, Larsson BA, et al. Ropivacaine pharmacokinetics after caudal block in 1–8-year-old children. *Br J Anaesth* 2000;85(4):506–511.
57. Bosenberg AT, Thomas J, Lopez T, et al. Plasma concentrations of ropivacaine following a single-shot caudal block of 1, 2 or 3 mg/kg in children. *Acta Anaesthesiol Scand* 2001;45(10):1276–1280.
58. American Society of Regional Anesthesia and Pain Medicine. Checklist for treatment of local anesthetic systemic toxicity. ASRA Pain Medicine; 2020 revision.
59. Advanced Life Support Group. Advanced Paediatric Life Support: A Practical Approach to Emergencies, 7th edition. Chichester: Wiley-Blackwell; 2023.
60. Luscombe MD, Owens BD. Weight estimation in resuscitation: is the current formula still valid? *Arch Dis Child* 2007;92(5):412–415.

- 61.** Luscombe MD, Owens BD, Burke D. Weight estimation in paediatrics: a comparison of the APLS formula and the formula 'Weight = 3(age)+7'. Emerg Med J 2011;28(7):590–593.