

Machine Learning and Data Analytics

ME 5013- Fall 2019

Lecture 04

- Multiple Linear Regression



The University of Texas at San Antonio™

Adel Alaeddini, PhD

Associate Professor of Mechanical Engineering

Advanced Data Engineering Lab

adel.alaeddini@utsa.edu

- Here our model is

$$Y = \beta_0 + \beta_1 X_1 + \beta_2 X_2 + \cdots + \beta_p X_p + \epsilon,$$

- We interpret β_j as the *average* effect on Y of a one unit increase in X_j , *holding all other predictors fixed*. In the advertising example, the model becomes

$$\textit{sales} = \beta_0 + \beta_1 \times \textit{TV} + \beta_2 \times \textit{radio} + \beta_3 \times \textit{newspaper} + \epsilon.$$

- The ideal scenario is when the predictors are uncorrelated
 - a *balanced design*:
 - Each coefficient can be estimated and tested separately.
 - Interpretations such as “*a unit change in X_j is associated with a β_j change in Y , while all the other variables stay fixed*”, are possible.
- Correlations amongst predictors cause problems:
 - The variance of all coefficients tends to increase, sometimes dramatically
 - Interpretations become hazardous — when X_j changes, everything else changes.
- *Claims of causality* should be avoided for observational data.

“Data Analysis and Regression” Mosteller and Tukey 1977

- A regression coefficient β_j estimates the expected change in Y per unit change in X_j , *with all other predictors held fixed*. But predictors usually change together!
- Example: Y total amount of change in your pocket;
 X_1 = # of coins; X_2 = # of pennies, nickels and dimes. By itself, regression coefficient of Y on X_2 will be > 0 . But how about with X_1 in model?
- Y = number of tackles by a football player in a season; W and H are his weight and height. Fitted regression model is $\hat{Y} = b_0 + 0.5W - 0.1H$. How do we interpret $\hat{\beta}_2 < 0$?

“Essentially, all models are wrong, but some are useful”

George Box

“The only way to find out what will happen when a complex system is disturbed is to disturb the system, not merely to observe it passively”

Fred Mosteller and John Tukey, paraphrasing George Box


- Given estimates $\hat{\beta}_0, \hat{\beta}_1, \dots, \hat{\beta}_p$, we can make predictions using the formula

$$\hat{y} = \hat{\beta}_0 + \hat{\beta}_1 x_1 + \hat{\beta}_2 x_2 + \cdots + \hat{\beta}_p x_p$$

- We estimate $\beta_0, \beta_1, \dots, \beta_p$ as the values that minimize the sum of squared residuals

$$\begin{aligned} \text{RSS} &= \sum_{i=1}^n (y_i - \hat{y}_i)^2 \\ &= \sum_{i=1}^n (y_i - \hat{\beta}_0 - \hat{\beta}_1 x_{i1} - \hat{\beta}_2 x_{i2} - \cdots - \hat{\beta}_p x_{ip})^2. \end{aligned}$$

This is done using standard statistical software. The values $\hat{\beta}_0, \hat{\beta}_1, \dots, \hat{\beta}_p$ that minimize RSS are the multiple least squares regression coefficient estimates.



	Coefficient	Std. Error	t-statistic	p-value
Intercept	2.939	0.3119	9.42	< 0.0001
TV	0.046	0.0014	32.81	< 0.0001
radio	0.189	0.0086	21.89	< 0.0001
newspaper	-0.001	0.0059	-0.18	0.8599

Correlation

	TV	radio	newspaper	sales
TV	1.0000	0.0548	0.0567	0.7822
radio		1.0000	0.3541	0.5762
newspaper			1.0000	0.2283
sales				1.0000

1. Is at least one of the predictors X_1, X_2, \dots, X_p useful in predicting the response?
2. Do all the predictors help to explain Y , or is only a subset of the predictors useful?
3. How well does the model fit the data?
4. Given a set of predictor values, what response value should we predict, and how accurate is our prediction?

$$F = \frac{(\text{TSS} - \text{RSS})/p}{\text{RSS}/(n - p - 1)} \sim F_{p, n-p-1}$$

Quantity	Value
Residual Standard Error	1.69
R^2	0.897
F-statistic	570

- The most direct approach is called **all subsets** or **best subsets** regression: we compute the least squares fit for all possible subsets and then choose between them based on some criterion that balances training error with model size.
- However we often can't examine all possible models, since they are 2^p of them; for example when $p = 40$ there are over a billion models!
- Instead we need an automated approach that searches through a subset of them. We discuss two commonly use approaches next.

- Begin with the *null model* — a model that contains an intercept but no predictors.
- Fit p simple linear regressions and add to the null model the variable that results in the lowest RSS.
- Add to that model the variable that results in the lowest RSS amongst all two-variable models.
- Continue until some stopping rule is satisfied, for example when all remaining variables have a p-value above some threshold.


- Start with all variables in the model.
- Remove the variable with the largest p-value — that is, the variable that is the least statistically significant.
- The new $(p - 1)$ -variable model is fit, and the variable with the largest p-value is removed.
- Continue until a stopping rule is reached. For instance, we may stop when all remaining variables have a significant p-value defined by some significance threshold.

- Later we discuss more systematic criteria for choosing an “optimal” member in the path of models produced by forward or backward stepwise selection.
- These include *Mallow's C_p* , *Akaike information criterion (AIC)*, *Bayesian information criterion (BIC)*, *adjusted R^2* and *Cross-validation (CV)*.

Qualitative Predictors

- Some predictors are not *quantitative* but are *qualitative*, taking a discrete set of values.
- These are also called *categorical* predictors or *factor variables*.
- See for example the scatterplot matrix of the credit card data in the next slide.

In addition to the 7 quantitative variables shown, there are four qualitative variables: *gender*, *student* (student status), *status* (marital status), and *ethnicity* (Caucasian, African American (AA) or Asian).



Example: investigate differences in credit card balance between males and females, ignoring the other variables. We create a new variable

$$x_i = \begin{cases} 1 & \text{if } i\text{th person is female} \\ 0 & \text{if } i\text{th person is male.} \end{cases}$$

Resulting model:

$$y_i = \beta_0 + \beta_1 x_i + \epsilon_i = \begin{cases} \beta_0 + \beta_1 + \epsilon_i & \text{if } i\text{th person is female} \\ \beta_0 + \epsilon_i & \text{if } i\text{th person is male.} \end{cases}$$

Intpretatio?

Results for gender model:

	Coefficient	Std. Error	t-statistic	p-value
Intercept	509.80	33.13	15.389	< 0.0001
gender[Female]	19.73	46.05	0.429	0.6690

- With more than two levels, we create additional dummy variables. For example, for the **ethnicity** variable we create two dummy variables. The first could be

$$x_{i1} = \begin{cases} 1 & \text{if } i\text{th person is Asian} \\ 0 & \text{if } i\text{th person is not Asian,} \end{cases}$$

and the second could be

$$x_{i2} = \begin{cases} 1 & \text{if } i\text{th person is Caucasian} \\ 0 & \text{if } i\text{th person is not Caucasian.} \end{cases}$$

- Then both of these variables can be used in the regression equation, in order to obtain the model

$$y_i = \beta_0 + \beta_1 x_{i1} + \beta_2 x_{i2} + \epsilon_i = \begin{cases} \beta_0 + \beta_1 + \epsilon_i & \text{if } i\text{th person is Asian} \\ \beta_0 + \beta_2 + \epsilon_i & \text{if } i\text{th person is Caucasian} \\ \beta_0 + \epsilon_i & \text{if } i\text{th person is AA.} \end{cases}$$

- There will always be one fewer dummy variable than the number of levels. The level with no dummy variable — African American in this example — is known as the *baseline*.

	Coefficient	Std. Error	t-statistic	p-value
Intercept	531.00	46.32	11.464	< 0.0001
ethnicity[Asian]	-18.69	65.02	-0.287	0.7740
ethnicity[Caucasian]	-12.50	56.68	-0.221	0.8260

Removing the additive assumption: *interactions* and *nonlinearity*


Interactions:

- In our previous analysis of the *Advertising* data, we assumed that the effect on *sales* of increasing one advertising medium is independent of the amount spent on the other media.
- For example, the linear model

$$\widehat{\text{sales}} = \beta_0 + \beta_1 \times \text{TV} + \beta_2 \times \text{radio} + \beta_3 \times \text{newspaper}$$

states that the average effect on *sales* of a one-unit increase in *TV* is always β_1 , regardless of the amount spent on *radio*.

- But suppose that spending money on radio advertising actually increases the effectiveness of TV advertising, so that the slope term for **TV** should increase as **radio** increases.
- In this situation, given a fixed budget of \$100, 000, spending half on **radio** and half on **TV** may increase **sales** more than allocating the entire amount to either **TV** or to **radio**.
- In marketing, this is known as a **synergy** effect, and in statistics it is referred to as an **interaction** effect.



When levels of either **TV** or **radio** are low, then the true **sales** are lower than predicted by the linear model.
But when advertising is split between the two media, then the model tends to underestimate **sales**.

Model takes the form

$$\begin{aligned} \text{sales} &= \beta_0 + \beta_1 \times TV + \beta_2 \times \text{radio} + \beta_3 \times (\text{radio} \times TV) + \epsilon \\ &= \beta_0 + (\beta_1 + \beta_3 \times \text{radio}) \times TV + \beta_2 \times \text{radio} + \epsilon \end{aligned}$$

Results:

	Coefficient	Std. Error	t-statistic	p-value
Intercept	6.7502	0.248	27.23	< 0.0001
TV	0.0191	0.002	12.70	< 0.0001
radio	0.0289	0.009	3.24	0.0014
TV×radio	0.0011	0.000	20.73	< 0.0001

- The results in this table suggests that interactions are important.
- The p-value for the interaction term $TV \times radio$ is extremely low, indicating that there is strong evidence for $H_A : \beta_3 \neq 0$.
- The R^2 for the interaction model is 96.8%, compared to only 89.7% for the model that predicts sales using TV and radio without an interaction term.

- This means that $(96.8 - 89.7)/(100 - 89.7) = 69\%$ of the variability in **sales** that remains after fitting the additive model has been explained by the interaction term.
- The coefficient estimates in the table suggest that an increase in TV advertising of \$1, 000 is associated with increased sales of $(\hat{\beta}_1 + \hat{\beta}_3 \times \text{radio}) \times 1000 = 19 + 1.1 \times \text{radio}$ units.
- An increase in radio advertising of \$1, 000 will be associated with an increase in sales of $(\hat{\beta}_2 + \hat{\beta}_3 \times \text{TV}) \times 1000 = 29 + 1.1 \times \text{TV}$ units.

- Sometimes it is the case that an interaction term has a very small p-value, but the associated main effects (in this case, TV and radio) do not.
- The *hierarchy principle*:

If we include an interaction in a model, we should also include the main effects, even if the p-values associated with their coefficients are not significant.



- The rationale for this principle is that interactions are hard to interpret in a model without main effects — their meaning is changed.
- Specifically, the interaction terms also contain main effects, if the model has no main effect terms.

Consider the **Credit** data set, and suppose that we wish to predict **balance** using **income** (quantitative) and **student** (qualitative).

Without an interaction term, the model takes the form


$$\begin{aligned} \text{balance}_i &\approx \beta_0 + \beta_1 \times \text{income}_i + \begin{cases} \beta_2 & \text{if } i\text{th person is a student} \\ 0 & \text{if } i\text{th person is not a student} \end{cases} \\ &= \beta_1 \times \text{income}_i + \begin{cases} \beta_0 + \beta_2 & \text{if } i\text{th person is a student} \\ \beta_0 & \text{if } i\text{th person is not a student.} \end{cases} \end{aligned}$$

$$\begin{aligned} \text{balance}_i &\approx \beta_0 + \beta_1 \times \text{income}_i + \begin{cases} \beta_2 + \beta_3 \times \text{income}_i & \text{if student} \\ 0 & \text{if not student} \end{cases} \\ &= \begin{cases} (\beta_0 + \beta_2) + (\beta_1 + \beta_3) \times \text{income}_i & \text{if student} \\ \beta_0 + \beta_1 \times \text{income}_i & \text{if not student} \end{cases} \end{aligned}$$



Credit data; Left: no interaction between **income** and **student**.
 Right: with an interaction term between **income** and **student**.

polynomial regression on Auto data



The figure suggests that

$$\text{mpg} = \beta_0 + \beta_1 \times \text{horsepower} + \beta_2 \times \text{horsepower}^2 + E$$

may provide a better fit.

	Coefficient	Std. Error	t-statistic	p-value
Intercept	56.9001	1.8004	31.6	< 0.0001
horsepower	-0.4662	0.0311	-15.0	< 0.0001
horsepower ²	0.0012	0.0001	10.1	< 0.0001