European Grapevine Moth (*Lobesia botrana*): A Comprehensive Report

I. Executive Summary

The European Grapevine Moth (EGVM), *Lobesia botrana* (Denis & Schiffermüller, 1776) is one of the most destructive lepidopteran pests of grapevines worldwide. Native to Southern Italy, this moth has spread across Europe, North Africa, the Middle East, and most recently North and South America. It poses major economic and ecological threats to viticulture (the cultivation and production of grapes, particularly for wine), due to its multivoltine (capable of producing more than two generations in a single year) life cycle, broad host range, and close association with secondary fungal pathogens such as *Botrytis cinerea*.

First detected in the United States in 2009 (Napa County, California), the EGVM triggered federal quarantine and eradication programs led by USDA–APHIS and the California Department of Food and Agriculture (CDFA). Understanding its biology, ecology, and management is essential to preventing reinfestation and to developing sustainable integrated-pest-management (IPM) programs.

II. Introduction

Viticulture is a cornerstone of Mediterranean and Californian agriculture, generating billions of dollars annually in table-grape, raisin, and wine industries. Among the numerous insect pests that affect vines, the European Grapevine Moth stands out for its unique ability to damage grapes directly while simultaneously inducing severe fungal infections such as the causative agent of bunch rot (Botrytis cinerea).

This report consolidates scientific findings from European, South American, and U.S. research programs to present an integrated review of the EGVM. It covers taxonomy, morphology, host range, developmental biology, damage patterns, and the current state of monitoring and management. The purpose is to provide an evidence-based overview that supports both academic study and practical vineyard decision-making.

III. Taxonomy and Distribution

Taxonomic position.

The EGVM is a member of the family Tortricidae, also known as leaf-roller moths, two were originally described from specimens collected near Vienna, Austria, in 1776, Lobesia has been historically known by synonyms including *Eudemis botrana*, *Polychrosis botrana*, and *Olindia rosmarinana*. It belongs to the subfamily Olethreutinae and is closely related to other vineyard

tortricids such as *Sparganothis pilleriana* and *Endopiza viteana* (the native grape berry moth of North America).

Global distribution.
Today, the EGVM occurs across the entire Mediterranean basin, Central Europe, the Middle East, and parts of North and West Africa. It has also been recorded in eastern Russia, Chile (2008), Argentina (2010), and Japan (likely misidentifications).
In the United States, infestations were confined to northern California (Napa, Sonoma, Solano, Mendocino, Monterey, and Fresno Counties) between 2009 and 2016 before eradication was declared in 2016 (CDFA, 2016).

IV. Host Range

The EGVM is primarily a pest of cultivated grapes (*Vitis vinifera*), though larvae can also develop on a variety of wild and cultivated plants. Host choice is influenced by fruit chemistry, phenology, and microclimate. While grape remains the preferred host, alternative plants sustain populations during early spring or in marginal habitats.

Table 1. Major and Minor Host Plants of *Lobesia botrana*

Category	Scientific Name	Common Name	Notes
Primary Hosts	*Vitis vinifera*	Cultivated grapevine	Main host throughout Europe and the Americas
	Daphne gnidium	Spurge-flax daphne	Key wild reservoir supporting early generations
Secondary Hosts	*Olea europaea*	Olive	Occasional infestation of flower clusters
	Prunus avium, *P. domestica*, *P. persica*	Cherry, plum, peach	Larvae may feed on blossoms or green fruit
	Punica granatum	Pomegranate	Damage reported in Spain and Israel
	Actinidia chinensis	Kiwi	Minor host in mixed orchards
Wild Hosts	*Rubus fruticosus*, *Ribes nigrum*	Blackberry, currant	Support off-season populations

Host range breadth helps explain the moth's persistence even where grapes are sparse. Early-season development on *Daphne gnidium* or *Ligustrum vulgare* (privet) enables migration into vineyards once grapes flower, ensuring continuous population growth.

V. Morphology and Identification

Adult moths are small (6–8 mm long; wingspan 11–13 mm). The Forewings exhibit a cream to tan ground color overlaid with bluish-gray and brown mottling, forming an irregular mosaic pattern. The hindwings are gray with fringed margins. At rest, adults characteristically hold their wings in a tent-like posture over the abdomen.

Eggs are lentil-shaped, 0.6–0.8 mm diameter, and laid singly on flower pedicels or berries. Newly laid eggs are iridescent white, turning yellow and then black near hatching.
Larvae pass through five instars, growing from 1 mm to 15 mm. Early in stars are creamy white with black heads; later instars become greenish or pinkish as gut contents show through the transparent cuticle. The prothoracic shield has a dark posterior margin and distinct white tubercles at the base of body setae. Fully grown larvae spin grayish-white silken cocoons under bark or in cluster debris before pupating.
Pupae measure 4–9 mm (longer in females) and are initially cream colored, darkening to brown. Diagnostic characters include a five-to-six-toothed anal comb and horizontal L-pinaculum on T1 (Gilligan & Epstein, 2012).

In the field, *L. botrana* may be confused with *Eupoecilia ambiguella* or *Endopiza viteana*; however, its wing pattern and genitalic features are distinctive. So identification is best confirmed by dissecting male genitalia or DNA barcoding of COI sequences.

VI. Biology and Seasonal Life Cycle

The EGVM is polyvoltine, producing two to four generations per year depending on climate, and temperature and photoperiod regulate development and diapause. In northern Europe (two generations), southern Europe (three), and Mediterranean basins (four partial generations), the life cycle adapts closely to vine phenology (Brière & Pracros, 1998; Milonas et al., 2001).

Adults emerge from overwintering pupae once temperatures exceed 10 °C (50 °F) for 10–12 days. Males emerge first and locate females through sex pheromones. Mating occurs shortly after dusk when temperatures are above 12 °C (54 °F). Females typically mate once and can lay 80–140 eggs, averaging 35 per day. Adult longevity ranges from 7 to 20 days depending on weather.

Table 2. Seasonal Life-Cycle Summary of *Lobesia botrana*

Generation	Time Period (Europe)	Host Stage Attacked	Key Damage	Notes
1st Generation	May – June	Flower clusters	Webbing and aborted flowers	Larvae form “glomerules” of silk-bound blossoms

Generation	Time Period (Europe)	Host Stage Attacked	Key Damage	Notes
2nd Generation	July – August	Green berries	Berry hollowing and frass deposition	Moderate crop loss and secondary fungi
3rd Generation	August – September	Ripening berries	Deep feeding and fruit rot	Highest economic impact
Overwintering Stage	October – April	—	Pupal diapause beneath bark or soil	High cold tolerance; resumes in spring

Egg incubation takes 3–10 days depending on temperature; larval development 20–30 days; and pupal stage 6–14 days. Developmental thresholds range from 10 °C (lower) to 30 °C (upper), with optimum between 26–29 °C and 40–70 % relative humidity (Del Tío et al., 2001).
Larvae in the first-generation feed on flower pedicels and can cause cluster drying. Second-generation larvae enter green berries, creating entry holes that invite fungal infection. Third-generation larvae feed within tight clusters near harvest, causing severe losses and encouraging *Botrytis* bunch rot.
Overwintering pupae survive temperatures as low as –10 °C and can endure for months in diapause. Adults resume flight in spring once degree-day accumulation surpasses ~833 DDF for the first generation and ~904 DDF for the second (Brière & Pracros, 1998).

VII. Damage and Economic Importance

The EGVM causes direct and indirect damage that can result in severe yield and quality losses in viticulture. Larval feeding disrupts the grapevine's reproductive structures across successive generations.

First-generation damage typically occurs in May and June when larvae feed on flower clusters, webbing blossoms together in "glomerules." Although yield loss from this stage may be modest, affected clusters dry up or drop prematurely.

Second-generation larvae (July–August) bore into developing berries, hollowing them out while leaving the skins intact. These damaged berries exude juice that attracts *Drosophila* fruit flies and ants.

Third-generation larvae (August–September) cause the most devastating injury as they penetrate ripening berries. Feeding creates entry points for opportunistic fungi such as *Botrytis cinerea*, *Aspergillus niger*, *Alternaria*, *Rhizopus*, *Cladosporium*, and *Penicillium*. This leads to "bunch rot," characterized by shriveled berries covered in fungal growth and frass.

Economic losses vary depending on infestation level, climate, and cultivar. In Mediterranean wine regions, yield reductions of **30–80%** have been recorded in untreated vineyards (Coscollá,

1998). In Chile, severe infestations following the 2008 introduction led to total crop loss in some areas (USDA–APHIS, 2010). In California, the potential threat was estimated at over $2 billion annually in grape-related economic activity (CDFA, 2016).

Beyond yield, *L. botrana* affects grape and wine quality through:

- Contaminated clusters increase volatile acidity and reduce sugar concentration.
- Fungal colonization alters flavor compounds and aromatic balance.
- Wineries face additional costs in sorting, cleaning, and fungicide applications.

These combined impacts make *L. botrana* not only a pest of economic significance but also a phytosanitary concern for global trade in grapes and grape products.

VIII. Monitoring and Detection

Early detection of *Lobesia botrana* populations is essential for successful management. The use of pheromone traps, field inspections, and phenological models allow precise timing of control measures.

Table 3. Monitoring and Management Summary for *Lobesia botrana*

Aspect	Recommended Practice	Scientific Basis / Source
Monitoring traps	Deploy red or orange delta traps baited with synthetic pheromone (1 trap per 30 acres, ≥5 ft above ground).	Del Tío et al., 2001
Egg and larval inspection	Weekly inspection of 100 clusters for eggs, webbing, or hollow berries.	UC Davis IPM, 2010
Degree-day models	Predict emergence using 833 DDF (1st gen.) and 904 DDF (2nd gen.) thresholds.	Brière & Pracros, 1998
Biological control	Release or conserve parasitoids such as *Dibrachys affinis* and *Campoplex capitator*.	Thiéry, 2008; Coscollá, 1998
Chemical control	Apply reduced-risk insecticides (spinosyns, Bt, methoxyfenozide) targeting newly hatched larvae.	Sáenz-de-Cabezón et al., 2005
Mating disruption	Use Isomate-EGVM pheromone dispensers (≥10 acres for best effect).	Louis et al., 2002
Quarantine procedures	Fumigation with methyl bromide under 7 CFR 305 T101 if exporting from infested areas.	USDA–APHIS, 2010

Pheromone monitoring.
Male moths are highly responsive to the female-produced sex pheromone, a blend dominated by (E, Z)-7,9-dodecadienyl acetate. Traps capture males to determine flight peaks and guide insecticide timing. In California, pheromone trapping was the primary detection method during the 2009–2016 eradication campaign (Gilligan & Epstein, 2012).

Field surveys.
Visual inspection complements trapping. Eggs are best found on pedicels or berry skins; their color transition (white → yellow → black) indicates developmental stage. Larvae can be detected by silk webbing, frass, and hollow berries. Inspecting one cluster per vine across representative vineyard blocks provides reliable population estimates.

Data integration.
Modern IPM programs use temperature data to calculate degree-days and forecast larval emergence. These predictive tools improve timing for control applications and reduce pesticide use.

IX. Management and Control Strategies

Management of *L. botrana* requires integration of biological, chemical, and cultural methods within a broader regulatory framework. The pest's multivoltine nature and cryptic habits make single method control ineffective.

1. Cultural Control

- **Sanitation:** Removal of infested clusters and pruning residues eliminates overwintering pupae.
- **Harvest hygiene:** Timely harvest reduces late-season infestation and fungal spread.
- **Canopy management:** Proper pruning improves air circulation and light penetration, creating unfavorable conditions for larval development and *Botrytis* infection.

2. Biological Control

Numerous natural enemies attack *L. botrana*, especially parasitoid wasps in the families Ichneumonidae, Braconidae, and Pteromalidae.
Key species include:

- *Campoplex capitator* (Ichneumonidae) — parasitizes larvae and pupae in Italy.
- *Dibrachys affinis* and *D. cavus* (Pteromalidae) — major pupal parasitoids causing up to 70% mortality in Spain (Thiéry, 2008).
- Tachinid flies (Diptera) — contribute additional control under natural vineyard ecosystems.

3. Mating Disruption

Pheromone-mediated mating disruption confuses males, reducing mating success and suppress population growth. Commercial formulations such as **Isomate-EGVM** or **Isonet-L Plus** release synthetic pheromone blends that mimic female signals.
Studies in Germany and Italy demonstrated up to **95% reduction in male captures** and **>80% reduction in larval density** when applied over continuous vineyard blocks exceeding 10 hectares (Louis et al., 2002).
Effectiveness depends on low initial population density, proper dispenser spacing, and minimal interference from adjacent untreated vineyards.

4. Chemical Control

Where populations exceed threshold levels, selective insecticides are used to protect susceptible stages. Recommended options include:

- **Spinosad (Entrust®)** and **spinetoram (Delegate®):** target young larvae with minimal non-target effects.
- **Methoxyfenozide (Intrepid®):** insect growth regulator disrupting molting.
- **Bacillus thuringiensis var. kurstaki (Bt):** microbial biopesticide used in organic vineyards.
  Spray timing is synchronized with egg hatch, typically shortly after peak male flight.
  Broad-spectrum pyrethroids (e.g., beta-cyfluthrin) are discouraged due to their negative effects on beneficial fauna.

5. Quarantine and Eradication

Following the 2009 detection in California, USDA–APHIS imposed strict quarantine measures under the **Plant Protection Act (§412a)**. Movement of grapes and nursery stock from infested areas required inspection, hot-water treatment, or fumigation. Methyl bromide fumigation (T101 protocol) was limited to export shipments where infestation was confirmed.

Extensive trapping, host removal, and coordinated pesticide applications eradicated *L. botrana* from California by 2016. The program demonstrated that area-wide IPM, combined with public cooperation, can achieve elimination of an invasive vineyard pest.

X. Quarantine and Regulatory Considerations

International trade regulations classify *L.botrana* as a quarantine pest of high concern under EPPO A2 List and the U.S. Federal Plant Pest Act.
Regulatory strategies emphasize:

- **Early detection:** mandatory pheromone trapping in risk zones.
- **Containment:** restriction on movement of grape materials from infested zones.

- **Documentation:** inspection and certification of nursery stock and harvested grapes.

The **Environmental Assessment for EGVM (USDA–APHIS, 2010)** identified minimal environmental risk from quarantine treatments such as methyl bromide fumigation when used under controlled conditions. The global exemption for quarantine and pre-shipment (QPS) applications ensures compliance with the **Montreal Protocol** while balancing pest control and environmental protection.

XI. Ongoing Research and Future Prospects

Contemporary research focuses on refining population modeling, improving biological-control implementation, and reducing pesticide reliance.
Key areas of investigation include:

- **Molecular identification**: COI barcoding distinguishes *L. botrana* from native tortricids, improving surveillance accuracy.
- **Semiochemical innovation:** improved pheromone blends and aerosol emitters for cost-efficient disruption.
- **Climate adaptation:** modeling range expansion under changing temperatures and precipitation regimes.
- **Microbial biocontrol:** entomopathogenic fungi and bacteria as environmentally friendly alternatives.
- **Behavioral ecology:** understanding female oviposition cues on host volatiles such as methyl salicylate and β-ocimene (Maher & Thiéry, 2006).

Such research enhances long-term management and supports sustainable viticulture practices globally.

XII. Conclusions

The European Grapevine Moth remains one of the most studied vineyard pests worldwide due to its adaptability, economic impact, and quarantine relevance. Effective management depends on precise monitoring, understanding of life cycles, and integration of cultural, biological, and pheromone-based tactics.
The California eradication program stands as a model of how science-based, cooperative management can successfully eliminate an invasive species.
Continued vigilance is required to prevent reintroduction and to protect the global grape industry from renewed losses

References

1. **Armendáriz, I., Campillo, G., Pérez-Sanz, A., & Marcos, M. A. (2007).**
   The grapevine moth Lobesia botrana in the Arribes D.O. (2004–2006 seasons).
   Boletín de Sanidad Vegetal – Plagas, 33, 477–489.

2. **Brière, J.-F., & Pracros, P. (1998).**
   Comparison of temperature-dependent models for the development of the grapevine moth *Lobesia botrana.*
   Environmental Entomology, 27(1), 94–101. https://doi.org/10.1093/ee/27.1.94

3. **California Department of Food and Agriculture (CDFA). (2016).**
   European Grapevine Moth (EGVM) Eradication Program Report.
   https://www.cdfa.ca.gov/plant/pdep/EGVM/index.html

4. **Coscollá, R. (1998).**
   Grapevine Cluster Moths (Lobesia botrana Den. & Schiff.).
   Madrid: Ministry of Agriculture, Spain.

5. **Del Tío, R., Martínez, J. L., Ocete, R., & Ocete, M. E. (2001).**
   Relationship between pheromone trap catches and degree-day accumulation in Sherry vineyards for *Lobesia botrana.*
   Journal of Applied Entomology, 125(1–2), 9–14. https://doi.org/10.1046/j.1439-0418.2001.00504.x

6. **Gilligan, T. M., & Epstein, M. E. (2012).**
   Identification and Biology of the European Grapevine Moth (Lobesia botrana).
   USDA–APHIS Technical Bulletin No. 2022.
   https://www.aphis.usda.gov/plant_health/plant_pest_info/egvm/downloads/egvm-id-biology.pdf

7. **Louis, F., Schmidt-Tiedemann, A., & Schirra, K. (2002).**
   Control of *Lobesia botrana* in German vineyards using pheromone-mediated mating disruption.
   IOBC/WPRS Bulletin, 25(7), 1–9.

8. **Maher, N., & Thiéry, D. (2006).**
   Influence of wild host *Daphne gnidium* on oviposition and larval performance of *Lobesia botrana.*
   Chemoecology, 16(3), 135–144. https://doi.org/10.1007/s00049-006-0335-2

9. **Milonas, P. G., Savopoulou-Soultani, M., & Stavridis, D. G. (2001).**
   Day-degree models for predicting *Lobesia botrana* flight activity in Greece.
   Journal of Applied Entomology, 125(9–10), 515–518. https://doi.org/10.1046/j.1439-0418.2001.00585.x

10. **Moreau, J., Benrey, B., & Thiéry, D. (2006).**
Grape cultivar influences larval and adult performance of *Lobesia botrana.*
Bulletin of Entomological Research, 96(3), 205–212.
https://doi.org/10.1079/BER2005417

11. **Sáenz-de-Cabezón, F., Maron, V., Zalom, F. G., & Pérez-Moreno, I. (2005).**
Efficacy of insect growth regulators on *Lobesia botrana* (Lepidoptera: Tortricidae).*
Pest Management Science, 61(8), 817–823. https://doi.org/10.1002/ps.1042

12. **Thiéry, D. (2008).**
Biology and ecology of the European Grapevine Moth (Lobesia botrana) in Mediterranean vineyards.
INRA Publications, Bordeaux, France.

13. **United States Department of Agriculture (USDA–APHIS). (2010).**
Environmental Assessment for the Control of the European Grapevine Moth in California.
https://www.aphis.usda.gov/plant_health/ea/downloads/egvm-ea.pdf