Quasi-Monte Carlo-Based Conditional Malliavin Method for Continuous-Time Asian Option Greeks

Chao Yu Xiaoqun Wang

September 30, 2021

Abstract

Although many methods for computing the Greeks of discrete-time Asian options are proposed, few methods to calculate the Greeks of continuous-time Asian options are known. In this paper, we develop an integration by parts formula in the multi-dimensional Malliavin calculus, and apply it to obtain the Greeks formulae for continuous-time Asian options in the multi-asset situation. We combine the Malliavin method with the quasi-Monte Carlo method to calculate the Greeks in simulation. We discuss the asymptotic convergence of simulation estimates for the continuous-time Asian option Greeks obtained by Malliavin derivatives. We propose to use the conditional quasi-Monte Carlo method to smooth Malliavin Greeks, and show that the calculation of conditional expectations analytically is viable for many types of Asian options. We prove that the new estimates for Greeks have good smoothness. For binary Asian options, Asian call options and up-and-out Asian call options, for instance, our estimates are infinitely times differentiable. We take the gradient principal component analysis method as a dimension reduction technique in simulation. Numerical experiments demonstrate the large efficiency improvement of the proposed method, especially for Asian options with discontinuous payoff functions.

1 Introduction

In mathematical finance, the growing emphasis on risk management issues has hastened efficient techniques for the computations of price sensitivities with respect to model parameters, namely, the Greeks, which can be expressed as

$$Greek := \frac{\partial}{\partial \alpha} \mathbb{E}_{\mathbb{Q}} \left[e^{-rT} H \right],$$

where $\mathbb{E}_{\mathbb{Q}}\left[e^{-rT}H\right]$ is the price of some option with the payoff H, and α is some model parameter. The Greeks play an important role in hedging financial derivatives.

There are numerous ways to calculate the Greeks of discrete path-dependent options, like the finite difference (FD) method [10], the pathwise (PW) method [15, 3] and the likelihood ratio (LR) method [11, 3]. However, when handling the Greeks of continuous path-dependent options, some of these methods have limitations in many cases due to the complexity of the continuous-time performance of the risky asset S_t . For example, the LR method could not be used directly to calculate the Greeks of continuous-time simple Asian options with payoff $H = f(\bar{S}_T)$, where $\bar{S}_T = \frac{1}{T} \int_0^T S_t dt$ is the continuous-time average of S_t and f is the payoff function, due to the unknown of the density of \bar{S}_T (hereafter we omit the 'continuous-time'). Moreover, the FD method produces biased estimates with large variance, and the PW method is invalid when the payoff function f is discontinuous (see [10]).

Fournié *et al.* [7] developed a new method to calculate the Greeks of options based on Malliavin calculus, and Benhamou [2] generalized the method for the Asian options. The main idea is to use the integration by parts formula in Malliavin calculus to eliminate the need of taking the derivative of the payoff function. By this mean, the Greeks of options have the following expression

$$Greek = \mathbb{E}_{\mathbb{Q}}\left[e^{-rT}H \cdot weight\right],$$

where *weight* is some random variable.

Fournié *et al.* [7] gave a particular solution for the weighting function in the case of the *delta* of simple Asian options with payoff $H = f(\bar{S}_T)$ in the one-asset case. However, there are very few results in the literature with regard to complex Asian options with payoff $H = f(S_T, \bar{S}_T)$ and those in the multi-asset case. Notice that neither the extension

from simple Asian options to complex Asian options nor that from the one-asset case to the multi-asset case is trivial since the original integration by parts formula cannot be applied directly (see Remarks 2.4 and 2.5 in Section 2).

To our knowledge, only Benhamou [2] considered the *delta* of complex Asian options. However, the calculation of the weight function is very complicated. For the multi-asset case, although Xu *et al.* [32] gave the *delta* and *gamma* of simple Asian options, the calculation is complex. Some idea can be extended to a more general integration by parts formula. For complex Asian options in the multi-asset case, no general integration by parts formula is known.

On the other hand, the Malliavin method gives us a chance to derive unbiased estimates of the Asian option Greeks. Nevertheless, we still need to use simulation to calculate Greeks since these estimates have no closed forms. The Monte Carlo (MC) and the quasi-Monte Carlo (QMC) methods are the usual simulation methods to estimate the integrals. Differently from the MC method, the QMC method uses deterministic low discrepancy points instead of random points to estimate the integrals. The QMC method provides a faster asymptotic convergence than MC for many financial problems (see [10]). But the high dimensionality and the lack of smoothness limit the advantage of QMC over MC. The gradient principal component analysis (GPCA) method [31] and the conditional quasi-Monte Carlo (CQMC) method [30, 1] are the two main powerful methods to overcome the high dimensionality and the lack of smoothness, respectively.

When applying the MC and the QMC methods to the Asian option Greeks, we need to discretize the path, which is different from the discrete path-dependent options. It leads to two important issues, namely, how to deal with the bias of the discrete simulation estimates and how to reduce the variance.

For the first issue, the discrete simulation estimates are directly used to approximate the original estimates of Malliavin Greeks in most relevant literature (see [2, 32]). There is few literature discussing the bias of the simulation estimates. For the second issue, Fournié *et al.* [7] were the first to develop a variance reduction technique, which localizes the integration by part formula around the singularity, to estimate the Malliavin Greeks in the MC setting. Xu *et al.* [32] showed the superiority of the QMC method over the MC method. However, as they mentioned, no variance reduction technique is taken. We believe that a more powerful method combining the QMC method with a more general variance reduction technique could be developed.

Based on the multi-dimensional Malliavin calculus, we develop a general integration by parts formula, and apply it to obtain the Greeks formulae for both simple and complex Asian options in the multi-asset Black-Scholes model. In simulation, we prove the asymptotic unbiasedness of the simulation estimates when the payoff function f is continuous with linear growth. Then based on the CQMC method, we develop a new method to calculate the Asian option Greeks. We smooth the involved payoff functions by taking conditional expectations, which is proved to be continuous under some uniform convergence conditions, and derive new estimates of the Greeks. The main contributions of this paper are as follows:

- In *m*-dimensional Malliavin calculus, we develop the integration by parts formula. Then we apply it to calculate the Greeks (*delta*, *gamma* and *vega*) for both simple and complex Asian options in the *m*-asset Black-Scholes model. In particular, we obtain the Greeks formulae for European options.
- We convert the estimates of the Malliavin Greeks to the discrete simulation estimates, and apply the CQMC method to calculate the option Greeks in continuous situation. For payoff functions with linear growth, we prove that the simulation estimates of Asian option Greeks are asymptotic unbiased.
- We propose to smooth the integrands for Greek estimates by taking conditional expectations. The Malliavin
 estimates for Greeks are usually discontinuous. However, after taking conditional expectations, the estimates
 for many Asian options have good smoothness. After then some potent dimension reduction techniques (such
 as the GPCA method) can be combined with the QMC method to obtain good effectiveness.
- We show how the conditional expectations can be calculated analytically. Since the simulation estimates for many Asian option Greeks (obtained by Malliavin calculus) satisfy the separation of variables condition, our method has wide practicability.

This paper is organized as follows. In Section 2, we introduce the basic theory of the Malliavin calculus and generalize the integration by parts formula. In Section 3, we use our formula to obtain the Greeks formulae for Asian options in the Black-Scholes model. We introduce the QMC method in Section 4. The main idea of our method is introduced in Section 5. The implementation is demonstrated on three examples in Section 6. Numerical experiments are performed in Section 7. Finally, we summarize our conclusions in Section 8.

2 Malliavin calculus: basic theory and results

In this section, we briefly introduce the basic theory of Malliavin Calculus in the multi-dimensional case. In addition, we develop an integration by parts formula in order to calculate the Greeks for the complex Asian options in Section 3. For more details of this subject, we refer to [22, 23].

Consider the classical filtered probability space $(\Omega, \mathscr{F}, \{\mathscr{F}_t\}_{0 \leq t \leq T}, \mathbb{P})$ and the m-dimensional Brownian motion $\mathbf{W} = \{\mathbf{W}_t = (W_t^{(1)}, \cdots, W_t^{(m)})'; 0 \leq t \leq T\}$, where Ω is the space of all continuous \mathbb{R}^m -valued functions ω on [0, T] such that $\omega(0) = \mathbf{0}$, \mathscr{F} is the completion of the Borel σ -algebra $\mathscr{B}(\Omega)$ with respect to the Wiener measure \mathbb{P} , $\mathbf{W}_t(\omega) = \omega(t)$ is the coordinate process and $\{\mathscr{F}_t\}_{0 \leq t \leq T}$ is the \mathbb{P} -augmentation of the filtration generated by \mathbf{W} , which satisfies the usual condition (see [17]).

We denote by $C_p^{\infty}(\mathbb{R}^n)$ the set of all infinitely continuously differentiable functions ϕ such that ϕ and all of its partial derivatives have polynomial growth. Let $\mathscr S$ denote the class of smooth random variables such that a random variable $X \in \mathscr S$ has the form

$$X = \varphi \left(\int_0^T \boldsymbol{h}_1(s) \cdot d\boldsymbol{W}_s, \cdots, \int_0^T \boldsymbol{h}_n(s) \cdot d\boldsymbol{W}_s \right),$$

where $n \in \mathbb{N}_+$, $\varphi \in C_p^{\infty}(\mathbb{R}^{m \times n})$ and $h_j \in L^2([0,T];\mathbb{R}^m)$ for $j=1,\cdots,n$. Note that \mathscr{S} is dense in $L^2(\Omega)$. The Malliavin gradient $D_t X$ of the smooth random variable X is defined as the $L^2([0,T];\mathbb{R}^m)$ -valued random variable $D_t X = (D_t^{(1)}X,\cdots,D_t^{(m)}X)'$ with components

$$D_t^{(i)}X = \sum_{i=1}^n \frac{\partial \varphi}{\partial x_{ij}} \left(\int_0^T \boldsymbol{h}_1(s) \cdot d\boldsymbol{W}_s, \cdots, \int_0^T \boldsymbol{h}_n(s) \cdot d\boldsymbol{W}_s \right) h_j^{(i)}(t), \quad i = 1, \cdots, m.$$

We can check that D_t is a closable operator from $\mathscr{S} \subset L^2(\Omega)$ to $L^2(\Omega; L^2([0,T];\mathbb{R}^m))$. Denote by $D^{1,2}(\Omega)$ the closure of the class of smooth random variables \mathscr{S} with respect to the graph norm

$$||X||_{D^{1,2}(\Omega)} := \left[||X||_{L^{2}(\Omega)}^{2} + ||D_{t}X||_{L^{2}(\Omega;L^{2}([0,T];\mathbb{R}^{m}))}^{2} \right]^{\frac{1}{2}}$$

$$= \left[\mathbb{E}\left[|X|^{2} \right] + \mathbb{E}\left[\int_{0}^{T} \sum_{i=1}^{m} |D_{t}^{(i)}X|^{2} dt \right] \right]^{\frac{1}{2}}.$$

Then D_t is a closed dense operator from $L^2(\Omega)$ to $L^2(\Omega; L^2([0,T];\mathbb{R}^m))$ with dense domain Dom $D_t = D^{1,2}(\Omega)$, which is a Banach space under the norm $\|\cdot\|_{D^{1,2}(\Omega)}$.

Similar to the ordinary gradient operator ∇_n in \mathbb{R}^n , there is also a chain rule for the Malliavin gradient D_t , which is important in the proof of the following integration by parts formula.

Proposition 2.1. (Proposition 1.2.4, [22]) Let $\varphi : \mathbb{R}^n \to \mathbb{R}$ be a Lipschitz function. Suppose that $X = (X_1, \dots, X_n)$ is a stochastic vector whose components belong to $D^{1,2}(\Omega)$, and the law of X is absolutely continuous with respect to the Borel-Lebesgue measure on \mathbb{R}^n . Then $\varphi(X) \in D^{1,2}(\Omega)$, and

(1)
$$D_t \varphi(X) = \sum_{j=1}^n \frac{\partial \varphi}{\partial x_j}(X) D_t X_j.$$

Now we define by $\tilde{\delta}: L^2(\Omega; L^2([0,T];\mathbb{R}^m)) \to L^2(\Omega)$ with domain Dom $\tilde{\delta}$ the adjoint of the closed dense operator $D_t: L^2(\Omega) \to L^2(\Omega; L^2([0,T];\mathbb{R}^m))$ that is called the Malliavin divergence operator. Although $\tilde{\delta}$ is abstract to understand, it coincides with the Itô integral with respect to W (see [17]) when $v \in L^2(\Omega; L^2([0,T];\mathbb{R}^m)) \cong L^2(\Omega \times [0,T];\mathbb{R}^m)$ is an \mathscr{F}_t -adapted \mathbb{R}^m -valued process, namely, $\tilde{\delta}(v) = \int_0^T v_t \cdot \mathrm{d}W_t$. We will keep use of the notation $\int_0^T v_t \cdot \mathrm{d}W_t := \tilde{\delta}(v)$ when $v \in \mathrm{Dom}\ \tilde{\delta}$. Thus we also call $\tilde{\delta}$ the Skorohod integral.

The following lemma is the multiplication formula of $\tilde{\delta}$, which can be very useful to the calculation in Section 3. The proof can be found in [22].

Lemma 2.2. Let $X \in D^{1,2}(\Omega)$ and $v \in Dom \ \tilde{\delta}$. Suppose that $Xv \in L^2(\Omega; L^2([0,T];\mathbb{R}^m))$. Then $Xv \in Dom \ \tilde{\delta}$, and we have

(2)
$$\tilde{\delta}(Xv) = X\tilde{\delta}(v) - \int_0^T \sum_{i=1}^m D_t^{(i)} X v_t^{(i)} dt.$$

¹In fact, this space is induced by the classical Wiener space $(\Omega, \mathcal{H}, \mathbb{P})$, where $\mathcal{H} \cong L^2([0,T]; \mathbb{R}^m)$ (see [22]).

In Proposition 6.2.1 of [22], Nualart gave an integration by parts formula, which plays an important role in calculating the Greeks of options in mathematical finance. Here we extend it to the three-dimensional case in the following theorem, which will be strongly used in Section 3 especially for the case of the complex Asian options in the multi-asset case.

Theorem 2.3. Let X_1, \dots, X_n be m-dimensional random row vectors whose components belong to $D^{1,2}(\Omega)$, and Y be an n-dimensional random vector. Consider an $L^2([0,T])$ -valued random vector $\mathbf{v}_t = (v_t^{(1)}, \dots, v_t^{(n)})$. Assume that the following conditions are satisfied

- (a) There exists an invertible $m \times m$ real matrix $\mathbf{c} = (\mathbf{c}_1, \dots, \mathbf{c}_m)$ with inverse $\mathbf{b} = (\mathbf{b}_1, \dots, \mathbf{b}_m)'$, such that for all $j = 1, \dots, n$, $\mathbf{D}_t \mathbf{X}_j = (\mathbf{c}_1 \mathbf{X}_t^{(j1)}, \dots, \mathbf{c}_m \mathbf{X}_t^{(jm)})$ for some $\mathbf{X}_t^{(jk)} \in L^2(\Omega; L^2([0,T]))$, $k = 1, \dots, m$;
- (b) The law of (X_1, \dots, X_n) is absolutely continuous with respect to the Borel-Lebesgue measure on \mathbb{R}^{mn} ;
- (c) For all $k = 1, \dots, m$, the $n \times n$ matrix $\zeta^{(k)} = (\zeta_{ij}^{(k)})_{n \times n} := ((X_t^{(jk)}, v_t^{(i)})_{L^2([0,T])})_{n \times n}$ is invertible a.s. with inverse $\tilde{\zeta}^{(k)}$, and $\sum_{k=1}^n \sum_{i=1}^n Y_i \tilde{\zeta}_{ik}^{(l)} v_t^{(k)} b_t' \in Dom \, \tilde{\delta}$, where $(\cdot, \cdot)_{L^2([0,T])}$ is the inner product in $L^2([0,T])$.

Then for any Lipschitz function $\varphi : \mathbb{R}^{mn} \to \mathbb{R}$, we have for all $l = 1, \dots, m$,

(3)
$$\mathbb{E}\left[\sum_{j=1}^{n} \frac{\partial \varphi}{\partial x_{jl}}(\boldsymbol{X}_{1}, \dots, \boldsymbol{X}_{n})Y_{j}\right] = \mathbb{E}\left[\varphi(\boldsymbol{X}_{1}, \dots, \boldsymbol{X}_{n})\tilde{\delta}\left(\sum_{k=1}^{n} \sum_{j=1}^{n} Y_{j}\tilde{\zeta}_{jk}^{(l)}v_{t}^{(k)}b_{l}^{\prime}\right)\right].$$

Proof. By Proposition 2.1, we have

$$\begin{split} & \boldsymbol{D}_{t}\boldsymbol{\varphi}(\boldsymbol{X}_{1},\cdots,\boldsymbol{X}_{n}) \\ &= \left(\begin{array}{cccc} \boldsymbol{D}_{t}\boldsymbol{X}_{1}, & \cdots, & \boldsymbol{D}_{t}\boldsymbol{X}_{n} \end{array}\right) \left(\begin{array}{c} \frac{\partial \boldsymbol{\varphi}}{\partial \boldsymbol{x}_{1}}(\boldsymbol{X}_{1},\cdots,\boldsymbol{X}_{n})' \\ & \vdots \\ \frac{\partial \boldsymbol{\varphi}}{\partial \boldsymbol{x}_{n}}(\boldsymbol{X}_{1},\cdots,\boldsymbol{X}_{n})' \end{array}\right) \\ &= \left(\begin{array}{cccc} (\boldsymbol{c}_{1}\boldsymbol{X}_{t}^{(11)},\cdots,\boldsymbol{c}_{m}\boldsymbol{X}_{t}^{(1m)}), & \cdots, & (\boldsymbol{c}_{1}\boldsymbol{X}_{t}^{(n1)},\cdots,\boldsymbol{c}_{m}\boldsymbol{X}_{t}^{(nm)}) \end{array}\right) \left(\begin{array}{c} \frac{\partial \boldsymbol{\varphi}}{\partial \boldsymbol{x}_{1}}(\boldsymbol{X}_{1},\cdots,\boldsymbol{X}_{n})' \\ & \vdots \\ \frac{\partial \boldsymbol{\varphi}}{\partial \boldsymbol{x}_{n}}(\boldsymbol{X}_{1},\cdots,\boldsymbol{X}_{n})' \end{array}\right). \end{split}$$

Multiplying both sides of the above equation by b_l yields

(4)
$$\sum_{i=1}^{m} b_{li} D_t^{(i)} \varphi(\boldsymbol{X}_1, \dots, \boldsymbol{X}_n) = \sum_{j=1}^{n} \frac{\partial \varphi}{\partial x_{jl}} (\boldsymbol{X}_1, \dots, \boldsymbol{X}_n) X_t^{(jl)}.$$

Take inner product of (4) and $v_t^{(k)}$ in $L^2([0,T])$ for $k=1,\cdots,n$. Then we have

$$(5) \qquad \begin{pmatrix} \int_{0}^{T} \sum_{i=1}^{m} D_{t}^{(i)} \varphi(\boldsymbol{X}_{1}, \cdots, \boldsymbol{X}_{n}) b_{li} v_{t}^{(1)} dt \\ \vdots \\ \int_{0}^{T} \sum_{i=1}^{m} D_{t}^{(i)} \varphi(\boldsymbol{X}_{1}, \cdots, \boldsymbol{X}_{n}) b_{li} v_{t}^{(n)} dt \end{pmatrix} = \begin{pmatrix} \zeta_{11}^{(l)} & \cdots & \zeta_{1n}^{(l)} \\ \vdots & \ddots & \vdots \\ \zeta_{n1}^{(l)} & \cdots & \zeta_{nn}^{(l)} \end{pmatrix} \begin{pmatrix} \frac{\partial \varphi}{\partial x_{ll}} (\boldsymbol{X}_{1}, \cdots, \boldsymbol{X}_{n}) \\ \vdots \\ \frac{\partial \varphi}{\partial x_{nl}} (\boldsymbol{X}_{1}, \cdots, \boldsymbol{X}_{n}) \end{pmatrix}.$$

Since $\zeta^{(l)}$ is invertible a.s. with inverse $\tilde{\zeta}^{(l)}$, we left multiply both sides of (5) by $Y\tilde{\zeta}^{(l)}$. Then formula (3) follows from the fact that $\tilde{\delta}$ is the adjoint of D_t .

Remark 2.4. The Malliavin gradient of X_j can be decomposed into a coefficient matrix c and a series of processes $(X_t^{(j1)}, \dots, X_t^{jm})$ for $j = 1, \dots, n$ in Theorem 2.3, which is the main extension from the original integration by parts formula in [22]. That will be more convenient to calculate the Greeks in the multi-asset case in Section 3.

Remark 2.5. It might be unrealistic to obtain a useful integration by parts formula in the three-dimensional case without the condition that $D_t X_j = (c_1 X_t^{(j1)}, \dots, c_m X_t^{(jm)})$ for $j = 1, \dots, n$, which means that $D_t X_1, \dots, D_t X_n$ share the same coefficient matrix c. This condition allows Theorem 2.3 to be used to calculate the complex Asian option Greeks in Section 3.2.

If we take m = n = 1, c = b = 1 and $X_t = D_t X$ in Theorem 2.3, we can obtain the classical integration by parts formula in the following corollary.

Corollary 2.6. (Proposition 6.2.1, [22]) Let $X \in D^{1,2}(\Omega)$, and Y be a random variable. Consider an $L^2([0,T])$ -valued random variable v_t . Assume that $(DX,v)_{L^2([0,T])} \neq 0$, $Yv(DX,v)_{L^2([0,T])} \in Dom\ \tilde{\delta}$, and the law of X is absolutely continuous with respect to the Borel-Lebesgue measure on \mathbb{R} . Then for any Lipschitz function $\varphi : \mathbb{R} \to \mathbb{R}$, we have

(6)
$$\mathbb{E}\left[\varphi'(X)Y\right] = \mathbb{E}\left[\varphi(X)\tilde{\delta}\left(Yv(DX,v)_{L^{2}([0,T])}^{-1}\right)\right].$$

3 Malliavin Greeks for Asian options

We assume that the financial market consists of one risk-free asset (bond) B and m risky assets (stocks) $S^{(1)}, \dots, S^{(m)}$. We recall the Black-Scholes model in the multi-asset case (see [10, 17]). Let $\mathbf{W} = \{\mathbf{W}_t = (W_t^{(1)}, \dots, W_t^{(m)})'; 0 \le t \le T\}$ be an m-dimensional Brownian motion on a filtered probability space $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{0 \le t \le T}, \mathbb{P})$, where $\{\mathcal{F}_t\}$ is the \mathbb{P} -augmentation of the filtration generated by \mathbf{W} , which satisfies the usual condition. The risky assets are the (unique) strong solutions of the following stochastic differential equation for $i = 1, \dots, m$

$$dS_t^{(i)} = \mu_i S_t^{(i)} dt + S_t^{(i)} \sum_{j=1}^m \sigma_{ij} dW_t^{(j)}, \quad 0 \le t \le T,$$

where $\mu = (\mu_1, \dots, \mu_m)'$ is a real vector and $\sigma = (\sigma_{ij})_{m \times m}$ is a real $m \times m$ matrix. By Itô's formula (see [17]), we can obtain the specific expression of $S_t^{(i)}$ as follows

$$S_t^{(i)} = S_0^{(i)} \exp\left\{ \left(\mu_i - \frac{1}{2} \sum_{j=1}^m \sigma_{ij}^2 \right) t + \sum_{j=1}^m \sigma_{ij} W_t^{(j)} \right\}, \quad 0 \le t \le T,$$

for $i = 1, \dots, m$. The risk-free asset B has the form $B_t = e^{rt}$, where r > 0 is the risk-free interest rate. We set $a := \sigma \sigma'$ and assume that for some $\lambda > 0$,

(7)
$$x'ax \ge \lambda ||x||^2, \quad \forall x \in \mathbb{R}^m.$$

Under condition (7), the process $M_t := \exp\{-\gamma \cdot W_t - \frac{1}{2} \|\gamma\|^2 t\}$ is a continuous martingale, where $\gamma = \sigma^{-1}(\mu - r\underline{1})$ and $\underline{1}$ is the m-dimensional column vector with every component equal to 1. By the Girsanov Theorem (see [17]), $\tilde{W}_t := W_t + \gamma t$ is an m-dimensional Brownian motion under the new probability measure (the risk-neutral probability measure) $\mathbb{Q} := \int M_T d\mathbb{P}$. Thus for $i = 1, \dots, m$, $S_t^{(i)}$ can be rewritten as

(8)
$$S_t^{(i)} = S_0^{(i)} \exp\left\{\omega_i t + \sum_{j=1}^m \sigma_{ij} \tilde{W}_t^{(j)}\right\}, \quad 0 \le t \le T,$$

where $\omega_i = r - \frac{1}{2} \sum_{j=1}^m \sigma_{ij}^2$.

3.1 Greeks for simple Asian options

We assume that the payoff of the simple Asian option is $f(\bar{S}_T)$, where the payoff function $f: \mathbb{R}^m \to \mathbb{R}$ is a nonnegative measurable function such that $f(\bar{S}_T) \in L^1(\Omega, \mathcal{F}, \mathbb{Q})$, and

$$\bar{S}_T = \frac{1}{T} \int_0^T S_t \mathrm{d}t$$

is the continuous-time average of S_t over the time interval [0,T]. Then the value of the option at time 0 is given by (see [17])

(9)
$$V = \mathbb{E}_{\mathbb{Q}} \left[e^{-rT} f(\bar{S}_T) \right].$$

The Greek of the option is the derivative (or higher derivative) of V with respect to a model parameter α given by

(10)
$$Greek := \frac{\partial V}{\partial \alpha} = \frac{\partial}{\partial \alpha} \mathbb{E}_{\mathbb{Q}} \left[e^{-rT} f(\bar{S}_T) \right].$$

if the above derivative exists. We only consider $\alpha = S_0^{(l)}$ and $\alpha = \sigma_{lp}$ for any fixed $l \in \{1, \cdots, m\}$ and $p \in \{1, \cdots, m\}$ in this paper. The correspondingly

If f is Lipschitz continuous, then (10) makes sense, and we can interchange the expectation and differentiation as follows (see [10])

(11)
$$\frac{\partial}{\partial \alpha} \mathbb{E}_{\mathbb{Q}} \left[e^{-rT} f(\bar{S}_T) \right] = e^{-rT} \mathbb{E}_{\mathbb{Q}} \left[\frac{\partial f}{\partial x_l} (\bar{S}_T) \frac{\partial \bar{S}_T^{(l)}}{\partial \alpha} \right].$$

If we take the partial derivative of f directly in (11) and obtain the relevant Greek formula, the well-known method is called the PW method (see [10]). In this case, the estimate of the Greek is dependent on the partial derivative of the payoff function f, which implies that this method might be failed when f is not Lipschitz continuous.

In view of the above, we turn to another way, i.e., the Malliavin method, which turns (11) to another form which excludes the partial derivative of f.

Since we only consider the expectation here, we identify the filtered probability space $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{0 \le t \le T}, \mathbb{Q})$ with the classical filtered probability space in Section 2.

As for the Malliavin derivatives we are going to use, we have for all $u \in [0, T]$ fixed (see [32, 22]),

$$D_{t}^{(i)}\tilde{W}_{u}^{(j)} = \delta_{ij}\mathbf{1}\{t \leq u\}, \qquad i, j = 1, \dots, m; \\ D_{t}S_{u}^{(j)} = (\sigma_{j1}\cdots,\sigma_{jm})'S_{u}^{(j)}\mathbf{1}\{t \leq u\}, \quad j = 1, \dots, m; \\ D_{t}\bar{S}_{u}^{(j)} = (\sigma_{j1}\cdots,\sigma_{jm})'\frac{1}{u}\int_{t}^{u}S_{v}^{(j)}dv, \quad j = 1, \dots, m,$$

where $\bar{S}_u := \frac{1}{u} \int_0^u S_v dv$, $\mathbf{1}\{\cdot\}$ is the indicator function, and δ denotes the Kronecker delta function.

Applying Theorem 2.3 to the right hand of (11) with n=1, $X=\bar{S}_T^{\prime}$, $Y=\frac{\partial \bar{S}_T^{(l)}}{\partial \alpha}$, $v_l=S_l^{(l)}$, $c=\sigma'$ and $X_l^{(k)}=S_l^{(k)}$ $\frac{1}{T} \int_t^T S_u^{(k)} du$ for $k = 1, \dots, m$, we can obtain the Greek formula by Malliavin calculus as follows

(12)
$$Greek = \mathbb{E}_{\mathbb{Q}} \left[e^{-rT} f(\bar{S}_T) \tilde{\delta} \left(\frac{\partial \bar{S}_T^{(l)}}{\partial \alpha} \frac{2S_t^{(l)}}{T \left(\bar{S}_T^{(l)} \right)^2} \sigma_l^{-1} \right) \right] =: \mathbb{E}_{\mathbb{Q}} [e^{-rT} f(\bar{S}_T) \cdot weight],$$

where σ_l^{-1} denotes the *l*th column of the matrix σ^{-1} , and *weight* is some random variable independent of the payoff function f. Then Lemma 2.2 can be used to obtain the specific expression of (12).

Since (12) excludes the partial derivative of f, we can extend it to the case that f is not necessarily Lipschitz continuous by the following lemma.

Lemma 3.1. (Lemma 12.28, [5]) Let $\alpha \mapsto w_{\alpha}$ be a process with the parameter α (i.e., $\alpha = S_0^{(l)}$ or $\alpha = \sigma_{lp}$) such that $\alpha \mapsto \mathbb{E}_{\mathbb{Q}}[w_{\alpha}^2] < \infty$ is locally bounded. Further assume that

(13)
$$\frac{\partial}{\partial \alpha} \mathbb{E}_{\mathbb{Q}}[e^{-rT} f(\bar{\mathbf{S}}_T)] = e^{-rT} \mathbb{E}_{\mathbb{Q}}[f(\bar{\mathbf{S}}_T) w_{\alpha}]$$

is valid for all $f \in C_c^{\infty}(\mathbb{R}^m)$ (i.e., f is an infinitely differentiable function with compact support). Then (13) also holds for all locally integrable function $f : \mathbb{R}^m \to \mathbb{R}$ such that $f(\bar{S}_T) \in L^2(\Omega, \mathcal{F}, \mathbb{Q})$.

Remark 3.2. We can also replace $f(\bar{S}_T)$ by $f(S_T, \bar{S}_T)$ in Lemma 3.1 in order to give a similar approximation of the Greeks for complex Asian options in Section 3.2.

We conclude the above analysis by the following theorem of the formulae for three common Greeks (delta, gamma and *vega*) of simple Asian options.

Theorem 3.3. Let $f(\bar{S}_T)$ be the payoff of the simple Asian option. If f is locally integrable and $f(\bar{S}_T) \in L^2(\Omega, \mathscr{F}, \mathbb{Q})$, then the formulae for delta, gamma and vega are given by

$$\begin{split} \textit{delta} : & \frac{\partial V}{\partial S_0^{(l)}} = \mathbb{E}_{\mathbb{Q}} \left[f(\bar{S}_T) \frac{2e^{-rT}}{S_0^{(l)}} \tau \right]; \\ \textit{gamma} : & \frac{\partial^2 V}{\partial S_0^{(p)} \partial S_0^{(l)}} = \mathbb{E}_{\mathbb{Q}} \left\{ f(\bar{S}_T) \frac{2e^{-rT}}{S_0^{(l)}} \left[\frac{2}{TS_0^{(p)} \bar{S}_T^{(p)}} \left(\tau \sum_{i=1}^m \int_0^T \sigma_{ip}^{-1} S_t^{(p)} \mathrm{d} \tilde{W}_t^{(i)} - \frac{\delta_{lp}}{\left(T \bar{S}_T^{(l)} \right)^2} \sum_{i=1}^m \sigma_{il}^{-1} H_{ilp} \right. \right. \\ & \left. + \frac{\tau T \bar{S}_T^{(p)}}{2} - \frac{\sigma_{lp}^{-1}}{T \bar{S}_T^{(l)}} \sum_{i=1}^m \sigma_{il}^{-1} \int_0^T S_t^{(l)} S_t^{(p)} \mathrm{d} t \right) - \frac{\delta_{lp}}{S_0^{(l)}} \tau \right] \right\}; \\ \textit{vega} : & \frac{\partial V}{\partial \sigma_{lp}} = \mathbb{E}_{\mathbb{Q}} \left[f(\bar{S}_T) \frac{2e^{-rT}}{\left(T \bar{S}_T^{(l)} \right)^2} \left(\int_0^T S_t^{(l)} g_t \mathrm{d} t \left(\tau + \frac{1}{2} \right) T \bar{S}_T^{(l)} - \int_0^T S_u^{(l)} \int_u^T \left(S_t^{(l)} g_t + S_t^{(l)} \sigma_{pl}^{-1} \right) \mathrm{d} t \mathrm{d} u \right) \right], \\ \text{where } g_t := \tilde{W}_t^{(p)} - \sigma_{lp} t, \ \tau := \frac{\sum_{i=1}^m \int_0^T \sigma_{il}^{-1} S_t^{(l)} \mathrm{d} \tilde{W}_t^{(i)}}{T \bar{S}_T^{(l)}} + \frac{1}{2}, \text{ and} \\ H_{ilp} := T \bar{S}_T^{(l)} \int_0^T S_u^{(p)} \int_u^T S_t^{(l)} \mathrm{d} \tilde{W}_t^{(i)} \mathrm{d} u - \int_0^T S_t^{(l)} \mathrm{d} \tilde{W}_t^{(i)} \int_0^T S_u^{(p)} \int_u^T S_t^{(l)} \mathrm{d} t \mathrm{d} u. \end{split}$$

When m = 1, from Theorem 3.3, we obtain formulae for *delta*, *gamma* and *vega* in one dimensional case in the following corollary.

Corollary 3.4. Let m = 1 and $f(\bar{S}_T)$ be the payoff of the simple Asian option. If f is locally integrable and $f(\bar{S}_T) \in L^2(\Omega, \mathcal{F}, \mathbb{Q})$, then the formulae for delta, gamma and vega are given by

$$\begin{aligned} & \textit{delta}: \frac{\partial V}{\partial S_0} = \mathbb{E}_{\mathbb{Q}}\left[f(\bar{S}_T)\frac{2e^{-rT}}{S_0\sigma^2}\left(\frac{S_T - S_0}{T\bar{S}_T} - \omega\right)\right]; \\ & \textit{gamma}: \frac{\partial^2 V}{\partial S_0^2} = \mathbb{E}_{\mathbb{Q}}\left[f(\bar{S}_T)\frac{4e^{-rT}}{\sigma^4S_0^2T^2\bar{S}_T^2}\left(S_T^2 - 2S_TS_0 + S_0^2 + \omega rT^2\bar{S}_T^2 - 2rTS_T\bar{S}_T + 2\omega TS_0\bar{S}_T\right)\right]; \\ & \textit{vega}: \frac{\partial V}{\partial \sigma} = \mathbb{E}_{\mathbb{Q}}\left[f(\bar{S}_T)\frac{2e^{-rT}}{\sigma^2T^2\bar{S}_T^2}\left(\left(S_T - S_0 - \left(r - \sigma^2\right)T\bar{S}_T\right)\int_0^T S_tg_t\mathrm{d}t - \sigma^2\int_0^T S_u\int_u^T S_tg_t\mathrm{d}t\mathrm{d}u - \frac{\sigma T^2}{2\sigma^2T^2\bar{S}_T^2}\right)\right]. \end{aligned}$$

The formulae for *delta* and *gamma* in Theorem 3.3 can also be found in [32]. However, our method is more concise and general since all the results are based on a general integration by parts formula in Theorem 2.3 rather than the complicated calculation for each specific example in [32]. Moreover, Theorem 2.3 could also be allowed to calculate the Greeks formulae for complex Asian options in the next subsection.

3.2 Greeks for complex Asian options

Now we consider the payoff of the complex Asian option $f(S_T, \bar{S}_T)$, where the payoff function $f: \mathbb{R}^{2m} \to \mathbb{R}$ is a nonnegative measurable function such that $f(S_T, \bar{S}_T) \in L^1(\Omega, \mathcal{F}, \mathbb{Q})$.

Similar to the case of the simple Asian option, if f is Lipschitz continuous, we can obtain the Greek of the complex Asian option as follows

(14)
$$Greek = \frac{\partial}{\partial \alpha} \mathbb{E}_{\mathbb{Q}} [e^{-rT} f(S_T, \bar{S}_T)]$$

$$= e^{-rT} \mathbb{E}_{\mathbb{Q}} \left[\frac{\partial f}{\partial x_{1l}} (S_T, \bar{S}_T) \frac{\partial S_T^{(l)}}{\partial \alpha} + \frac{\partial f}{\partial x_{2l}} (S_T, \bar{S}_T) \frac{\partial \bar{S}_T^{(l)}}{\partial \alpha} \right].$$

However, when dealing with the complex Asian option, the choice of v_t in Theorem 2.3 is important since a good choice can simplify our formulae greatly. We introduce the set Λ defined by

$$\Lambda := \{ a \in L^2([0,T]) : \int_0^T a_t dt = 0, \int_0^T a_t \int_0^t S_u^{(k)} du dt \neq 0, \forall k = 1, \dots, m \}.$$

Let $a \in \Lambda$. Then applying Theorem 2.3 to the right hand of (14) with n = 2, $X_1 = S_T'$, $X_2 = \bar{S}_T'$, $Y = (\frac{\partial S_T^{(l)}}{\partial \alpha}, \frac{\partial \bar{S}_T^{(l)}}{\partial \alpha})$, $v_t = (a_t, S_t^{(l)})$, $c = \sigma'$, $X_t^{(1k)} = S_T^{(k)}$ and $X_t^{(2k)} = \frac{1}{T} \int_t^T S_u^{(k)} du$ for $k = 1, \dots, m$, we can obtain the matrix

(15)
$$\zeta = \begin{pmatrix} 0 & -\frac{1}{T} \int_0^T a_l \int_0^t S_u^{(l)} du dt \\ T S_T^{(l)} \bar{S}_T^{(l)} & \frac{T}{2} \left(\bar{S}_T^{(l)} \right)^2 \end{pmatrix},$$

and the Greek formula

Greek

$$(16) \qquad = \mathbb{E}_{\mathbb{Q}} \left[e^{-rT} f(S_T, \bar{S}_T) \tilde{\delta} \left(\left(\frac{\frac{T}{2} \bar{S}_T^{(l)} \frac{\partial S_T^{(l)}}{\partial \alpha} a_t - T S_T^{(l)} \frac{\partial \bar{S}_T^{(l)}}{\partial \alpha} a_t}{S_T^{(l)} \int_0^T a_v \int_0^v S_u^{(l)} du dv} + \frac{\partial S_T^{(l)}}{\partial \alpha} \frac{S_t^{(l)}}{T S_T^{(l)} \bar{S}_T^{(l)}} \right) \sigma_l^{-1} \right) \right]$$

$$= : \mathbb{E}_{\mathbb{Q}} [e^{-rT} f(S_T, \bar{S}_T) \cdot weight].$$

By Lemma 3.1 and Remark 3.2, the formula (16) still holds when f is locally integrable and $f(S_T, \bar{S}_T) \in L^2(\Omega, \mathcal{F}, \mathbb{Q})$. We conclude the above analysis by the following theorem of the formulae for the Greeks for complex Asian options.

Theorem 3.5. Let $f(S_T, \bar{S}_T)$ be the payoff of the complex Asian option, and $a \in \Lambda$. If f is locally integrable and $f(S_T, \bar{S}_T) \in L^2(\Omega, \mathcal{F}, \mathbb{Q})$, then the formulae for delta, gamma and vega are given by

$$\begin{split} delta : \frac{\partial V}{\partial S_0^{(I)}} &= \mathbb{E}_{\mathbb{Q}} \left[f(S_T, \bar{S}_T) \frac{e^{-rT}}{S_0^{(I)}} \left(\frac{1}{T \bar{S}_T^{(I)}} \sum_{i=1}^m \int_0^T \sigma_{il}^{-1} S_t^{(I)} \mathrm{d} \tilde{W}_t^{(i)} + \frac{\int_0^T a_t \int_t^T S_u^{(I)} \mathrm{d} u \mathrm{d} t}{2h_T} \right. \\ &\quad + \frac{1}{2} - \frac{T \bar{S}_T^{(I)} \sum_{i=1}^m \int_0^T \sigma_{il}^{-1} a_t \mathrm{d} \tilde{W}_t^{(i)}}{2h_T} - \frac{T \bar{S}_T^{(I)} \int_0^T a_t \int_t^T a_v \int_t^v S_u^{(I)} \mathrm{d} u \mathrm{d} v \mathrm{d} t}{2h_T^2} \right) \right]; \\ gamma : \frac{\partial^2 V}{\partial S_0^{(P)} \partial S_0^{(I)}} &= \mathbb{E}_{\mathbb{Q}} \left[f(S_T, \bar{S}_T) \frac{e^{-rT}}{S_0^{(P)} S_0^{(I)}} \tilde{\delta} \left(\sigma_l^{-1} \left(\frac{T}{2} \bar{S}_T^{(I)} w_1 a_t - T S_T^{(I)} w_2 a_t + \frac{T}{2} \bar{S}_T^{(I)} w_1 a_t - T S_T^{(I)} w_2 a_t \right) \right) \right]; \\ vega : \frac{\partial V}{\partial \sigma_{lp}} &= \mathbb{E}_{\mathbb{Q}} \left[f(S_T, \bar{S}_T) e^{-rT} \left(\frac{T \bar{S}_T^{(I)} g_T}{2h_T} \sum_{i=1}^m \int_0^T \sigma_{il}^{-1} a_t \mathrm{d} \tilde{W}_t^{(i)} + \frac{g_T}{2} - \sigma_{pl}^{-1} + \frac{T \bar{S}_T^{(I)} g_T}{2h_T^2} \int_0^T a_t \int_t^T a_v \int_t^V S_u^{(I)} \mathrm{d} u \mathrm{d} v \mathrm{d} t \right. \\ &\quad - \frac{1}{2h_T} \int_0^T \left(g_T a_t \int_t^T S_u^{(I)} \mathrm{d} u + T \bar{S}_T^{(I)} \sigma_{pl}^{-1} a_t \right) \mathrm{d} t - \frac{S_T^{(I)} \int_0^T S_t^{(I)} g_t \mathrm{d} t}{S_T^{(I)} h_T} \sum_{i=1}^m \int_0^T \sigma_{il}^{-1} a_t \mathrm{d} \tilde{W}_t^{(i)} \\ &\quad - \frac{\int_0^T S_t^{(I)} g_t \mathrm{d} t}{h_T^2} \int_0^T a_t \int_t^T a_v \int_t^V S_u^{(I)} \mathrm{d} u \mathrm{d} v \mathrm{d} t + \frac{1}{h_T} \int_0^T a_t \int_t^T \left(S_u^{(I)} g_u + \sigma_{pl}^{-1} S_u^{(I)} \right) \mathrm{d} u \mathrm{d} t \right) \right], \end{aligned}$$

where $h_t := \int_0^t a_V \int_0^V S_u^{(l)} du dv$, $\tilde{h}_t := \int_0^t a_V \int_0^V S_u^{(p)} du dv$, $g_t := \tilde{W}_t^{(p)} - \sigma_{lp}t$, and

$$\begin{split} w_1 &= -\frac{T\bar{S}_T^{(p)}S_T^{(l)}}{2\tilde{h}_T} \sum_{i=1}^m \int_0^T \sigma_{ip}^{-1} a_t \mathrm{d} \tilde{W}_t^{(i)} - \delta_{lp} S_T^{(l)} + \frac{1}{2} + \frac{S_T^{(l)}}{2\tilde{h}_T} \int_0^T a_t \int_t^T S_u^{(p)} \mathrm{d} u \mathrm{d} t - \frac{T\bar{S}_T^{(p)}S_T^{(l)}}{2\tilde{h}_T^2} \int_0^T a_t \int_t^T a_v \int_t^V S_u^{(p)} \mathrm{d} u \mathrm{d} v \mathrm{d} t \\ &+ \frac{S_T^{(l)}}{T\bar{S}_T^{(p)}} \sum_{i=1}^m \int_0^T \sigma_{ip}^{-1} S_t^{(p)} \mathrm{d} \tilde{W}_t^{(i)}; \\ w_2 &= -\frac{T\bar{S}_T^{(p)}\bar{S}_T^{(l)}}{2\tilde{h}_T} \sum_{i=1}^m \int_0^T \sigma_{ip}^{-1} a_t \mathrm{d} \tilde{W}_t^{(i)} + \frac{\bar{S}_T^{(l)}}{2\tilde{h}_T} \int_0^T a_t \int_t^T S_u^{(p)} \mathrm{d} u \mathrm{d} t - \frac{\delta_{lp}\bar{S}_T^{(p)}}{2} + \frac{\bar{S}_T^{(l)}}{2} + \delta_{lp} \frac{\bar{S}_T^{(p)}}{2\tilde{h}_T} \int_0^T a_t \int_t^T S_u^{(l)} \mathrm{d} u \mathrm{d} t \\ &- \frac{T\bar{S}_T^{(p)}\bar{S}_T^{(l)}}{2\tilde{h}_T^2} \int_0^T a_t \int_t^T a_v \int_t^V S_u^{(p)} \mathrm{d} u \mathrm{d} v \mathrm{d} t + \frac{\bar{S}_T^{(l)}}{T\bar{S}_T^{(p)}} \sum_{i=1}^m \int_0^T \sigma_{ip}^{-1} S_t^{(p)} \mathrm{d} \tilde{W}_t^{(i)}. \end{split}$$

Remark 3.6. If $f(x_1, x_2)$ is only dependent on x_1 in Theorem 3.5, namely, $f(S_T, \bar{S}_T) = f(\bar{S}_T)$, we can obtain the Greeks formulae for European options.

When m = 1, we can choose $a_t = \frac{T}{2} - t$ in Theorem 3.5 to obtain the specific expression of the formulae in one dimensional case. For example, the formula for *delta* is given in the following corollary.

Corollary 3.7. Let m = 1 and $f(S_T, \bar{S}_T)$ be the payoff of the complex Asian option. If f is locally integrable and $f(S_T, \bar{S}_T) \in L^2(\Omega, \mathcal{F}, \mathbb{Q})$, then the formula for delta is given by

$$delta: \frac{\partial V}{\partial S_0} = \mathbb{E}_{\mathbb{Q}} \left[f(S_T, \bar{S}_T) \frac{e^{-rT}}{S_0} \left(\frac{1}{\sigma^2 T \bar{S}_T} \left(S_T - S_0 - rT \bar{S}_T \right) - \frac{T \bar{S}_T}{2\sigma h_T} \left(\int_0^T \tilde{W}_t dt - \frac{T}{2} \tilde{W}_T \right) + \frac{\int_0^T \int_t^T S_u \left(\frac{T}{2} - t \right) du dt}{2h_T} \right) \right] + \frac{1}{2} - \frac{T \bar{S}_T \int_0^T \int_t^T \int_t^V S_u \left(\frac{T}{2} - v \right) \left(\frac{T}{2} - t \right) du dv dt}{2h_T^2} \right].$$

Lacking closed form solutions for the expectations (integrals) in Theorems 3.3 and 3.5, the Greeks above could only be calculated by numerical methods. The rest of this paper is focused on developing a QMC-based method to calculate the expectations in Theorems 3.3 and 3.5 more efficiently.

4 Quasi-Monte Carlo method

MC and QMC are two important methods in numerical simulation to calculate high-dimensional integral over the cube $[0,1)^d$,

$$I(F) = \int_{(0,1]^d} F(\boldsymbol{u}) d\boldsymbol{u} = \mathbb{E}[F(\boldsymbol{U})],$$

where U is a d-dimensional uniform random variable in $(0,1]^d$. Compared with the MC method, QMC uses deterministic points, which are more uniformly distributed over $(0,1]^d$ instead of random points. The QMC estimate of I(F) can be written as

$$Q_n(F) = \frac{1}{n} \sum_{i=1}^n F(\mathbf{u}_i),$$

where u_1, \dots, u_n are deterministic low discrepancy points. The QMC error can be bounded by the Koksma-Hlawka inequality

$$|I(F) - Q_n(F)| < V_{HK}(F)D^*(u_1, \dots, u_n),$$

where $V_{\rm HK}(F)$ is the variation of F in the sense of the paper of Hardy and Krause (see [21]), and $D^*(u_1, \dots, u_n)$ is the star discrepancy. This inequality asserts that the convergence rate of QMC for functions of finite variation is $O(n^{-1}(\ln n)^d)$, which is asymptotically much better than the MC rate $O(n^{-1/2})$. We can choose many low discrepancy sequences to achieve this rate, such as the Sobol' sequence (see [25]), the Faure sequence (see [6]) and the Halton sequence (see [12]).

Despite the advantage of QMC over MC in asymptotic convergence rate, there are two crucial factors limiting the efficiency of QMC in practice, namely, the smoothness of the integrand and the high dimensionality (see [20]).

Although high dimensionality is a big challenge for QMC, it may still provide better performance if the function has low effective dimension (see [4, 27]). In order to enhance the efficiency of QMC, some path generation methods (PGMs) are developed to reduce the effective dimension. To introduce PGM, we consider the following expectation in mathematical finance

$$I(g) = \mathbb{E}[g(X)]$$

for some function g, where $X \sim N(\mathbf{0}_d, \Sigma)$ and Σ is the covariance matrix. The essence of a PGM is to generate the samples of X as X = AY, where $Y \sim N(\mathbf{0}_d, I_d)$ and the matrix A satisfies $AA' = \Sigma$. This is equivalent to transform the integral (18) as

(19)
$$I(g) = \mathbb{E}[g(X)] = \mathbb{E}[g(AY)].$$

The choices of A make no difference in MC setting from the point of view of variance (because the variance is unchanged), whereas different choices of A may lead to different efficiency in QMC setting due to the possible change of the effective dimension and the smoothness properties. Classical methods of PGM include standard (STD) construction, Brownian bridge (BB) construction and principal components analysis (PCA) construction (see [10]). Many new methods are proposed to reduce the effective dimension, such as the linear transform (LT) construction (see [16]), the clustering QR (CQR) construction (see [29]) and the GPCA construction (see [31]). The GPCA construction

performs best in many cases, especially when the integrand has complicated structure, since it uses the structure and information of the integrand (see [30]).

Bad smoothness is another factor that could seriously decrease the efficiency of the QMC method. Strenuous efforts are devoted to conquering it, such as the orthogonal transformation (OT) method (see [28]), the QR method (see [14]), and the CQMC method (see [30]).

In practice, the Koksma-Hlawka inequality is not very useful although it asserts that QMC provides a better asymptotic convergence rate. The first reason is that it is hard to calculate the variation $V_{\rm HK}(f)$. The second reason is that many functions in finance have unbounded variation. Moreover, since the QMC points are deterministic, we need to randomize QMC points to obtain error estimation. Randomized quasi-Monte Carlo (RQMC) methods are proposed to solve those problems, such as the scrambling, the digital shift, and the random shift (see [24, 19, 18]).

5 QMC-based conditional Malliavin method for Asian option Greeks

We focus on calculating the following expectation

$$(20) I(g) = \mathbb{E}[g(X)],$$

where $X = (X_1, \dots, X_d)'$ is a d-dimensional random vector with independent components, and g(X) is integrable. Let $\rho_k(x_k)$ be the density function of X_k , and $\rho(x_u) = \prod_{k \in u} \rho_k(x_k)$ for $u \subseteq \{1, \dots, d\}$.

5.1 Conditional QMC method

CQMC method is a smoothing method similar to the CMC method (see [9]). We describe the main idea of CQMC method below. We refer to [30] for details.

Suppose that the function g satisfies the so-called variable separation condition

(21)
$$g(\boldsymbol{x}) = h(\boldsymbol{x})\mathbf{1}\{\psi_d(\boldsymbol{z}) < x_j < \psi_u(\boldsymbol{z})\},$$

for some x_j , where h is a continuous function and ψ_d and ψ_u are functions of $z := (x_1, \dots, x_{j-1}, x_{j+1}, \dots, x_d)'$. Then the expectation (20) can be written as

$$I(g) = \mathbb{E}\left[g(\boldsymbol{X})\right] = \int_{\mathbb{R}^{d-1}} \left(\int_{\psi_d}^{\psi_d} h(\boldsymbol{x}) \rho_j(x_j) \mathrm{d}x_j\right) \mathbf{1}\{\psi_d(\boldsymbol{z}) < \psi_u(\boldsymbol{z})\} \rho(\boldsymbol{z}) \mathrm{d}\boldsymbol{z}.$$

Suppose that the conditional expectation

(22)
$$\mathbb{E}\left[g(\boldsymbol{X})|\boldsymbol{Z}\right] = \int_{\psi_d}^{\psi_u} h(\boldsymbol{x})\rho_j(x_j) \mathrm{d}x_j \mathbf{1}\{\psi_d(\boldsymbol{z}) < \psi_u(\boldsymbol{z})\} =: G(\boldsymbol{Z})$$

is analytically available, then we have

(23)
$$I(g) = \mathbb{E}[g(X)] = \mathbb{E}[\mathbb{E}[g(X)|Z]] = \mathbb{E}[G(Z)].$$

Therefore, G(Z) is a new estimate of I(g), which is called the conditional estimate.

Many functions in financial engineering satisfy the separation of variables condition (see the expression (27) for the function g of Asian option Greeks in the next subsection). The method of conditional expectation above is a form of CMC to reduce the variance. The idea can also be used in QMC setting if we use the QMC points instead of MC points in calculating $\mathbb{E}[G(Z)]$, and we call it CQMC method.

It is well-known that the performance of QMC integration strongly depends on the smoothness of the integrand. If the integrand possesses excellent smoothness, the calculation efficiency may be increased significantly. Furthermore, the effective dimension reduction method (such as the GPCA method) can be used to achieve better results.

The next theorem can be proved similarly as the Theorem A.1 in [33]. Therefore, we omit the proof of Theorem 5.2. It shows the good smoothness of the new function G. Before that we need to introduce a definition of uniform convergence.

Definition 5.1. We say the integral $\int \xi(y,x) dx$ converges uniformly on a set \mathscr{O} , if for any $\varepsilon > 0$ and $y \in \mathscr{O}$, there exists a constant A independent of y such that $\left| \int_{\|x\| > A} \xi(y,x) dx \right| < \varepsilon$.

Note that here y and x can be scalars or vectors, and ||x|| represents the Euclidean norm of x.

Theorem 5.2. Suppose that function g has the form in (21) and G is defined by (22). Assume that the following conditions are satisfied

- (a) ψ_d and ψ_u are continuous;
- (b) h is nonnegative.

Then the following two statements are equivalent

- (c) If one of ψ_u and ψ_d is infinite, then the integral $\int_{\mathbb{R}} h(x) \rho_j(x_j) dx_j$ converges uniformly on a compact neighborhood of z, where $z := (x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_d)'$ for some j;
- (d) G is continuous.

Theorem 5.2 gives conditions for which G is a continuous function. For most common Asian options, condition (a) is hold. Note that condition (c) is still sufficient for (d) even without the condition (b). Thus if both ψ_d and ψ_u are finite, then G is continuous, and if one of them is infinite, we need an extra condition of uniform convergence. The next lemma, which is similar to the Lemma 4.2 in [33], shows that the assumption of uniform convergence is also satisfied for most common Asian options.

Lemma 5.3. For fixed j, assume that h(x) has the form $h(x) = \sum_{k=0}^{n} A_k x_j^k e^{Bx_j} - C$ for some positive integer n, where A_k , B and C are functions of z (given in Theorem 5.2), bounded on a compact set Λ but independent of x_j . Then $\int h(x)\phi(x_j) dx_j$ converges uniformly on Λ , where ϕ is the density function of the standard normal variable x_j .

By Lemma 5.3 we can easily verify the uniform convergence condition (c) in Theorem 5.2. Fortunately, in common cases the function h has the form given in Lemma 5.3 (see Sections 6 for the following concrete examples: binary Asian options, Asian call options and up-and-out Asian call options).

5.2 QMC-based conditional Malliavin method for calculating Greeks

For convenience, from now on we only consider the Greeks for simple Asian options in the one-dimensional case (hereafter we omit the 'simple'). However, all results in this subsection can be extended to the general situation.

We presented the formulae for Asian option Greeks by Malliavin calculus in Corollary 3.4. For *delta*, *gamma* and *vega*, the Geeks have the following expression

(24)
$$= \mathbb{E}_{\mathbb{Q}} \left[e^{-rT} f(\bar{S}_T) \cdot weight \left(\bar{S}_T, S_T, \int_0^T S_t \left(\tilde{W}_t - \sigma t \right) dt, \int_0^T S_u \int_u^T S_t \left(\tilde{W}_t - \sigma t \right) dt du \right) \right]$$

$$= : \mathbb{E}_{\mathbb{Q}} [\hat{\theta}]$$

$$= : \theta.$$

In the numerical simulation, we usually take MC methods to estimate θ , and this requires us to discretize the continuous path first. We may choose equidistant points $0 = t_0 < t_1 < \cdots < t_d = T$ between time 0 and the maturity date T, at which we get the price of underlying asset S_{t_j} in discrete times. For simplicity, we denote S_{t_j} by S_j . Based on the expression (8) of S under the risk-neutral probability measure \mathbb{Q} , we have

(25)
$$S_j = S_0 \exp\left(\omega t_j + \sigma \tilde{W}_{t_j}\right), \quad 0 \le t \le T.$$

To simulate the continuous-time average $\bar{S}_T = \frac{1}{T} \int_0^T S_t dt$, let

$$S_A := \frac{1}{d} \sum_{i=1}^d S_i$$

be the discrete average of the asset prices. Then by the theory of Riemann integral, we have the pointwise convergence as follows

$$\lim_{d\to+\infty} S_A = \bar{S}_T.$$

It is natural to take the following simulation estimate $\hat{\theta}_d$ to approximate θ in (24)

(26)
$$\hat{\theta}_d = e^{-rT} f(S_A) \cdot weight\left(S_A, S_d, \frac{T}{d} \sum_{j=1}^d S_j(\tilde{W}_{t_j} - \sigma t_j), \frac{T^2}{d^2} \sum_{i=1}^d \sum_{j=i}^d S_i S_j(\tilde{W}_{t_j} - \sigma t_j)\right).$$

Although in most literature $\hat{\theta}_d$ is treated as identical to $\hat{\theta}$ directly, we can not ignore the fact that $\hat{\theta}_d$ is a biased estimate of θ . Fortunately, we can prove that $\hat{\theta}_d$ is an asymptotic unbiased estimate with respect to d when f is a continuous function with linear growth.

Theorem 5.4. Let $f(\bar{S}_T)$ be the payoff of the Asian option. If f is continuous with linear growth, then $\hat{\theta}_d$ is an asymptotic unbiased estimate of θ with respect to d, i.e.,

$$\lim_{d \to +\infty} \mathbb{E}_{\mathbb{Q}}[\hat{ heta}_d] = \mathbb{E}_{\mathbb{Q}}[\hat{ heta}] = heta.$$

Proof. From the formulae of Greeks in Corollary 3.4, we only need to consider the convergence of the following expectations

$$\mathbb{E}_{\mathbb{Q}}\left[f(S_A)\frac{S_d^{m_1}}{S_A^{m_2}}\right],$$

$$\mathbb{E}_{\mathbb{Q}}\left[f(S_A)\frac{S_d^{m_1}}{S_A^{m_2}}\frac{1}{d}\sum_{j=1}^d S_j(\tilde{W}_{t_j}-\sigma t_j)\right],$$

$$\mathbb{E}_{\mathbb{Q}}\left[\frac{f(S_A)}{S_A^2}\frac{1}{d^2}\sum_{i=1}^d \sum_{j=i}^d S_i S_j(\tilde{W}_{t_j}-\sigma t_j)\right],$$

where m_1 and m_2 are nonnegative integers.

Since f is continuous, we have $\lim_{d\to +\infty} f(S_A) = f(\bar{S}_T)$. Thus each integrand in the above expectations are convergent pointwisely. By dominated convergence theorem, we only need to consider the upper and lower bounds of each integrand.

For the first integrand, since f is nonnegative with linear growth, we have

$$0 \leq f(S_A) \frac{S_d^{m_1}}{S_A^{m_2}} \leq C(1+S_A) \frac{S_d^{m_1}}{S_A^{m_2}} \leq C\left(1 + \max_{0 \leq t \leq T} S_t\right) \frac{S_d^{m_1}}{\left(\min_{0 \leq t \leq T} S_t\right)^{m_2}},$$

where *C* is a positive constant.

For the second integrand, we have

$$\begin{split} f(S_A) \frac{S_d^{m_1}}{S_A^{m_2}} \frac{1}{d} \sum_{j=1}^d S_j(\tilde{W}_{t_j} - \sigma t_j) &\geq f(S_A) \frac{S_d^{m_1}}{S_A^{m_2}} \min_{0 \leq t \leq T} S_t(\tilde{W}_t - \sigma t) \\ &\geq -C \left(1 + \max_{0 \leq t \leq T} S_t \right) \frac{S_d^{m_1}}{(\min_{0 \leq t \leq T} S_t)^{m_2}} \left[\min_{0 \leq t \leq T} S_t(\tilde{W}_t - \sigma t) \right]^-, \end{split}$$

and

$$f(S_A) \frac{S_d^{m_1}}{S_A^{m_2}} \frac{1}{d} \sum_{j=1}^d S_j(\tilde{W}_{t_j} - \sigma t_j) \le f(S_A) \frac{S_d^{m_1}}{S_A^{m_2}} \max_{0 \le t \le T} S_t(\tilde{W}_t - \sigma t) \\ \le C \left(1 + \max_{0 \le t \le T} S_t \right) \frac{S_d^{m_1}}{(\min_{0 \le t \le T} S_t)^{m_2}} \left[\max_{0 \le t \le T} S_t(\tilde{W}_t - \sigma t) \right]^+,$$

where $[\cdot]^+$ and $[\cdot]^-$ denotes the positive part and the negative part of a function, respectively. For the third integrand, we have

$$\begin{split} \frac{f(S_A)}{S_A^2} \frac{1}{d^2} \sum_{i=1}^{d} \sum_{j=i}^{d} S_i S_j (\tilde{W}_{t_j} - \sigma t_j) &\geq \frac{f(S_A)}{S_A^2} \frac{d^2 + d}{2d^2} \min_{\substack{0 \leq u \leq T \\ u \leq t \leq T}} S_u S_t (\tilde{W}_t - \sigma t) \\ &\geq -\frac{C \left(1 + \max_{0 \leq t \leq T} S_t\right)}{\left(\min_{0 \leq t < T} S_t\right)^2} \left[\min_{\substack{0 \leq u \leq T \\ u \leq t \leq T}} S_u S_t (\tilde{W}_t - \sigma t) \right]^{-}, \end{split}$$

and

$$\begin{split} \frac{f(S_A)}{S_A^2} \frac{1}{d^2} \sum_{i=1}^{d} \sum_{j=i}^{d} S_i S_j (\tilde{W}_{t_j} - \sigma t_j) &\leq \frac{f(S_A)}{S_A^2} \frac{d^2 + d}{2d^2} \max_{\substack{0 \leq u \leq T \\ u \leq t \leq T}} S_u S_t (\tilde{W}_t - \sigma t) \\ &\leq \frac{C \left(1 + \max_{0 \leq t \leq T} S_t\right)}{\left(\min_{0 \leq t \leq T} S_t\right)^2} \left[\max_{\substack{0 \leq u \leq T \\ u \leq t < T}} S_u S_t (\tilde{W}_t - \sigma t) \right]^+. \end{split}$$

Thus the conclusion follows.

From the above proof, we can also get the following result in the Asian options pricing.

Corollary 5.5. Let $f(\bar{S}_T)$ be the payoff of the Asian option. If f is continuous with linear growth, then we have

$$\lim_{d\to+\infty}\mathbb{E}_{\mathbb{Q}}[f(S_A)]=\mathbb{E}_{\mathbb{Q}}[f(\bar{S}_T)],$$

where $S_A = \frac{1}{d} \sum_{j=1}^{d} S_j$ is the discrete average of the asset prices.

It is open about the bias or the asymptotic unbiasedness of $\hat{\theta}_d$ when f is not continuous. Nevertheless, we still use $\hat{\theta}_d$ to estimate θ in practice.

Note that $\mathbb{E}_{\mathbb{Q}}[\hat{\theta}_d]$ can be expressed as

(27)
$$\mathbb{E}_{\mathbb{Q}}[\hat{\theta}_d] = \mathbb{E}_{\mathbb{Q}}[g(X)],$$

where $X = (X_1, \dots, X_d)'$ is a d-dimensional standard normal random vector under the risk-neutral probability measure \mathbb{Q} (see the next section). Thus the CQMC method can be applied to calculate the Greeks.

We defined $G(Z) = \mathbb{E}_{\mathbb{Q}}[g(X)|Z]$ the conditional Malliavin (CMV) estimate of the Greeks in Section 5.1. If the CMV estimate G(Z) is analytically available, we can use QMC method to estimate the Greeks as follows

(28)
$$\mathbb{E}_{\mathbb{Q}}[g(\boldsymbol{X})] = \mathbb{E}_{\mathbb{Q}}[G(\boldsymbol{Z})] \approx \frac{1}{n} \sum_{i=1}^{n} G(\Phi^{-1}(\boldsymbol{u}_i)),$$

where u_1, \dots, u_n are (d-1)-dimensional RQMC points, and Φ is the distribution function of Z. This is QMC-based CMV method. We call it QMC-CMV method for short.

Note that we only consider the continuous-time Asian option Greeks in this paper. For the discrete-time Asian option Greeks, the Malliavin estimate degenerates into the likelihood ratio (LR) estimate, and there are many powerful methods based on the LR method and the pathwise (PW) method combined with CQMC method to improve the smoothness. We refer to [30, 33] for more details.

6 QMC-CMV method: implementation and examples

In this section, we show the implementation of the QMC-CMV method, and give the CMV estimates for the Greeks of three common Asian option Greeks (i.e., binary Asian options, Asian call options and up-and-out Asian call options) as illustrations.

By discretization, we get the price of underlying asset S_i at time t_i . To separate variables, let

(29)
$$\tilde{S}_j = S_0 \exp\left(\omega(t_j - t_1) + \sigma(\tilde{W}_{t_j} - \tilde{W}_{t_1})\right) = S_0 \exp\left(\omega(t_j - t_1) + \sigma\bar{W}_{t_j - t_1}\right),$$

where

$$\bar{W}_t := \tilde{W}_{t+t_1} - \tilde{W}_{t_1}.$$

Obviously, $\bar{W} = \{\bar{W}_t, \mathscr{F}_{t+t_1}\}_{0 \le t \le T-t_1}$ is also a Brownian motion under \mathbb{Q} . Comparing (25) with (29), we have

$$(30) S_j = \tilde{S}_j \exp\left(\omega t_1 + \sigma \tilde{W}_{t_1}\right).$$

Let $\bar{\boldsymbol{W}} = (\bar{W}_{t_2-t_1}, \cdots, \bar{W}_{t_d-t_1})'$. Because \tilde{W}_{t_1} and $\bar{\boldsymbol{W}}$ are independent and normally distributed, we can generate them by

(31)
$$\tilde{W}_{t_1} = \sqrt{t_1} X_1, \quad X_1 \sim N(0, 1),$$

and

(32)
$$\bar{\boldsymbol{W}} = \boldsymbol{A}\boldsymbol{Z}, \quad \boldsymbol{Z} \sim N(\boldsymbol{0}_{d-1}, \boldsymbol{I}_{d-1}),$$

where $Z = (X_2, \dots, X_d)'$, $\mathbf{0}_{d-1}$ is a (d-1)-dimensional zero column vector, I_{d-1} is a $(d-1) \times (d-1)$ identify matrix, and A is a $(d-1) \times (d-1)$ matrix satisfying $AA' = \Sigma$ with

$$\Sigma = \begin{pmatrix} t_2 - t_1 & t_2 - t_1 & \cdots & t_2 - t_1 \\ t_2 - t_1 & t_3 - t_1 & \cdots & t_3 - t_1 \\ \vdots & \vdots & \ddots & \vdots \\ t_2 - t_1 & t_3 - t_1 & \cdots & t_d - t_1 \end{pmatrix}.$$

Different choice of A may lead to different efficiency of QMC-based method. Since the Greek estimates can be viewed as a function of X, namely, $\mathbb{E}_{\mathbb{Q}}[\hat{\theta}_d] = \mathbb{E}_{\mathbb{Q}}[g(X)]$, and we have shown that the CMV estimate $G(Z) = \mathbb{E}_{\mathbb{Q}}[g(X)|Z]$ is continuous for common options, some PGMs (such as GPCA) can be used to reduce the effective dimension of G(Z).

Let $X = (X_1, Z')'$. Its components are independently and normally distributed. Let $S = (S_1, \dots, S_d)$. We have

(33)
$$\tilde{S}_{i} = S_{0} \exp\left(\omega(t_{i} - t_{1}) + \sigma A_{i-1} Z\right),$$

and

$$(34) S_i = S_0 \exp\left(\omega t_i + \sigma \sqrt{t_1} X_1 + \sigma A_{i-1} Z\right) = \exp\left(\omega t_1 + \sigma \sqrt{t_1} X_1\right) \tilde{S}_i,$$

where $A_0 = \mathbf{0}'_{d-1}$ is a zero row vector and A_j is the *j*th row of A. The variable S_j is a function of X, with the component X_1 and Z being separated (\tilde{S}_j is a function of Z).

By the above method, we achieve the goal of variables separation. In the expression of the Greek estimates $\mathbb{E}_{\mathbb{Q}}[g(X)]$, the variable separation condition is usually satisfied for g. In some cases, we can calculate the analytical expression of G(Z).

Example 6.1. As a warm-up example, we consider the binary Asian option whose payoff function is given by

$$f(\bar{S}_T) = \mathbf{1}\{\bar{S}_T > K\},\,$$

where K is the strike price. By Corolarry 3.4 and (26), the simulation estimate of delta is given by

$$\mathbb{E}_{\mathbb{Q}}[\hat{\theta}_d] = \mathbb{E}_{\mathbb{Q}}\left[\frac{2e^{-rT}}{S_0\sigma^2}\left(\frac{S_d - S_0}{TS_A} - \omega\right)\mathbf{1}\{S_A > K\}\right].$$

Since an indicator function is involved in the integrand, a direct use of QMC method could lead to large loss of efficiency. Based on (34), we have

$$S_A = \exp\left(\omega t_1 + \sigma \sqrt{t_1} X_1\right) \frac{1}{d} \sum_{i=1}^d \tilde{S}_j = \exp\left(\omega t_1 + \sigma \sqrt{t_1} X_1\right) \tilde{S}_A,$$

where

$$\tilde{S}_A := \frac{1}{d} \sum_{j=1}^d \tilde{S}_j = \frac{1}{d} \sum_{j=1}^d S_0 \exp\left(\boldsymbol{\omega}(t_j - t_1) + \boldsymbol{\sigma} \boldsymbol{A}_{j-1} \boldsymbol{Z}\right).$$

Thus we have

$${S_A > K} = {X_1 > \psi_d},$$

where

$$\psi_d := \frac{\ln K - \ln \tilde{S}_A - \omega t_1}{\sigma_{\sqrt{t_1}}}$$

is a function of Z. So we achieve variable separation, and the delta can be written by

$$\mathbb{E}_{\mathbb{Q}}[\hat{\theta}_d] = \mathbb{E}_{\mathbb{Q}}\left[\frac{2e^{-rT}}{S_0\sigma^2}\left(\frac{S_d - S_0}{TS_A} - \omega\right)\mathbf{1}\{X_1 > \psi_d\}\right] =: \mathbb{E}_{\mathbb{Q}}\left[g(\boldsymbol{X})\right],$$

where g(x) has the form (21).

We denote the density and distribution function of one-dimensional normal distribution by ϕ and Φ , respectively. To use CMV method (22), we need to calculate the analytical expression of $\mathbb{E}_{\mathbb{Q}}[g(X)|Z]$. This can be done as follows

$$\begin{split} &\mathbb{E}_{\mathbb{Q}}\left[g(\boldsymbol{X})|\boldsymbol{Z}\right] \\ &= \int_{\psi_d}^{\infty} \frac{2e^{-rT}}{S_0\sigma^2} \left(\frac{S_d - S_0}{TS_A} - \boldsymbol{\omega}\right) \phi(x_1) \mathrm{d}x_1 \\ &= \frac{2e^{-rT}\tilde{S}_d}{TS_0\tilde{S}_A\sigma^2} \Phi(-\psi_d) - \frac{2e^{-rT - rt_1 + \sigma^2 t_1}}{T\tilde{S}_A\sigma^2} \Phi(-\sigma\sqrt{t_1} - \psi_d) - \frac{2e^{-rT}\boldsymbol{\omega}}{S_0\sigma^2} \Phi(-\psi_d) \\ &= : G(\boldsymbol{Z}). \end{split}$$

Then G(Z) is the CMV estimate of *delta* for the binary Asian option. Clearly, G(Z) has good smoothness.

Similarly, we can calculate CMV estimates of *gamma* and *vega* for the binary Asian option. To sum up, we give the following theorem.

Theorem 6.2. For the binary Asian option, the CMV estimates of delta, gamma and vega are given by

$$\begin{split} delta: G_{\Delta,1}(K; \mathbf{Z}) &= \left(\frac{2e^{-rT}\tilde{S}_d}{TS_0\tilde{S}_A\sigma^2} - \frac{2e^{-rT}\omega}{S_0\sigma^2}\right) \Phi_1 - \frac{2e^{-rT-rt_1+\sigma^2t_1}}{T\tilde{S}_A\sigma^2} \Phi_2; \\ gamma: G_{\Gamma,1}(K; \mathbf{Z}) &= \frac{4e^{-rT}}{\sigma^4S_0^2T^2} \left[\left(\frac{\tilde{S}_d^2}{\tilde{S}_A^2} + \omega r T^2 - \frac{2rT\tilde{S}_d}{\tilde{S}_A}\right) \Phi_1 + \frac{S_0^2}{\tilde{S}_A^2} e^{-2rt_1+3\sigma^2t_1} \Phi_3 + \left(\frac{2\omega TS_0}{\tilde{S}_A} - \frac{2S_0\tilde{S}_d}{\tilde{S}_A^2}\right) e^{-rt_1+\sigma^2t_1} \Phi_2 \right]; \\ vega: G_{\vartheta,1}(K; \mathbf{Z}) &= \Phi_1 \left(\frac{2e^{-rT}(\tilde{S}_d - (r-\sigma^2)T\tilde{S}_A)}{\sigma^2T\tilde{S}_A^2d} \sum_{j=1}^d \tilde{S}_j(A_{j-1}\mathbf{Z} - \sigma t_j) - \frac{2e^{-rT}}{d^2\tilde{S}_A^2} \sum_{i=1}^d \sum_{j=i}^d \tilde{S}_i\tilde{S}_j(A_{j-1}\mathbf{Z} - \sigma t_j) - \frac{e^{-rT}}{\sigma} \right) \\ &+ \Phi_2 \left(\frac{2e^{-rT}S_0\sqrt{t_1}}{\sigma^2T\tilde{S}_A}\sigma\sqrt{t_1}e^{\sigma^2t_1/2-\omega t_1} - \frac{2S_0e^{-rT}}{\sigma^2Td\tilde{S}_A^2}e^{-rt_1+\sigma^2t_1} \sum_{j=1}^d \tilde{S}_j(A_{j-1}\mathbf{Z} - \sigma t_j) \right) \\ &+ \frac{1}{\sqrt{2\pi}}e^{-\psi_d^2/2} \left(\frac{2e^{-rT}\tilde{S}_d\sqrt{t_1}}{\sigma^2T\tilde{S}_A} - \frac{2e^{-rT}}{\sigma^2}\sqrt{t_1}(r-\sigma^2) - \frac{2e^{-rT}\sqrt{t_1}}{\tilde{S}_A^2d^2} \sum_{i=1}^d \sum_{j=i}^d \tilde{S}_i\tilde{S}_j \right) \\ &- \frac{2e^{-rT}S_0\sqrt{t_1}}{\sigma^2T\tilde{S}_A} \frac{1}{\sqrt{2\pi}}e^{\sigma^2t_1/2-\omega t_1}e^{-(\psi_d+\sigma\sqrt{t_1})^2/2}, \end{split}$$

where
$$\Phi_1 := \Phi(-\psi_d)$$
, $\Phi_2 := \Phi(-\sigma\sqrt{t_1} - \psi_d)$ and $\Phi_3 := \Phi(-2\sigma\sqrt{t_1} - \psi_d)$.

We can easily see that all CMV estimates given above are infinitely times differentiable, which are QMC-friendly. On the other hand, based on Corollary 3.4, the original Malliavin estimates of the Greeks involve discontinuities. Thus we may expect that CMV method could provide better results than the original Malliavin method in QMC.

As an example, for delta, the QMC-CMV estimate is given by

$$delta \approx \frac{1}{n} \sum_{i=1}^{n} \left[\frac{2e^{-rT} \tilde{S}_{d}(\Phi^{-1}(\boldsymbol{u}_{i}))}{T S_{0} \tilde{S}_{A}(\Phi^{-1}(\boldsymbol{u}_{i})) \sigma^{2}} \Phi(-\psi_{d}(\Phi^{-1}(\boldsymbol{u}_{i}))) - \frac{2e^{-rT-rt_{1}+\sigma^{2}t_{1}}}{T \tilde{S}_{A}(\Phi^{-1}(\boldsymbol{u}_{i})) \sigma^{2}} \Phi(-\sigma \sqrt{t_{1}} - \psi_{d}(\Phi^{-1}(\boldsymbol{u}_{i}))) - \frac{2e^{-rT-rt_{1}+\sigma^{2}t_{1}}}{T \tilde{S}_{A}(\Phi^{-1}(\boldsymbol$$

where u_1, \dots, u_n are QMC points.

Example 6.3. Consider the Asian call option with the following payoff function

$$f(\bar{S}_T) = (\bar{S}_T - K)^+$$
.

Using the notations and imitating the steps in Example 6.1, we can obtain the CMV estimates for the Asian call option Greeks in the following theorem.

Theorem 6.4. For the Asian call option, the CMV estimates for delta, gamma and vega are given by

$$\begin{split} delta: & G_{\Delta,2}(K; Z) = -\frac{2e^{-rT}}{T\sigma^2} \Phi_1 + \left(\frac{2e^{-rT}\tilde{S}_d}{TS_0\sigma^2} e^{rt_1} - \frac{2e^{-rT}\omega\tilde{S}_A}{S_0\sigma^2} e^{rt_1}\right) \tilde{\Phi}_2 - KG_{\Delta,1}(K; Z); \\ gamma: & G_{\Gamma,2}(K; Z) = \left(\frac{8\omega e^{-rT}}{\sigma^4S_0T} - \frac{8e^{-rT}\tilde{S}_d}{\sigma^4S_0T^2\tilde{S}_A}\right) \Phi_1 + \frac{4e^{-rT}}{\sigma^4T^2\tilde{S}_A} e^{-rt_1 + \sigma^2t_1} \Phi_2 \\ & + \left(\frac{4e^{-rT}\tilde{S}_d^2}{\sigma^4S_0^2T^2\tilde{S}_A} e^{rt_1} + \frac{4e^{-rT}\omega\tilde{S}_A^2}{\sigma^4S_0^2} e^{rt_1} - \frac{8r\tilde{S}_de^{-rT}}{\sigma^4S_0^2T} e^{rt_1}\right) \tilde{\Phi}_2 - KG_{\Gamma,1}(K; Z); \\ vega: & G_{\vartheta,2}(K; Z) = -\frac{2e^{-rT}S_0}{\sigma^2T\tilde{S}_Ad} \sum_{j=1}^d \tilde{S}_j (A_{j-1}Z - \sigma t_j) \Phi_1 - KG_{\vartheta,1}(K; Z) - \frac{2e^{-rT}S_0\sqrt{t_1}}{\sigma^2T} \frac{1}{\sqrt{2\pi}} e^{-\psi_d^2/2} \\ & + \left[\frac{2e^{-rT}\tilde{S}_d}{\sigma^2T\tilde{S}_Ad} \sum_{j=1}^d \tilde{S}_j (A_{j-1}Z - \sigma t_j) e^{rt_1} + \left(\frac{2e^{-rT}\tilde{S}_d\sqrt{t_1}}{\sigma^2T} - \frac{2e^{-rT}\tilde{S}_A}{\sigma^2} \sqrt{t_1} (r - \sigma^2) \right) \right. \\ & - \frac{2e^{-rT}\sqrt{t_1}}{\tilde{S}_Ad^2} \sum_{i=1}^d \sum_{j=i}^d \tilde{S}_i \tilde{S}_j \right) \sigma \sqrt{t_1} e^{\sigma^2t/2 + \omega t} - \frac{2e^{-rT}(r - \sigma^2)}{\sigma^2d} \sum_{j=1}^d \tilde{S}_j e^{rt_1} (A_{j-1}Z - \sigma t_j) \\ & - \frac{2e^{-rT}\sqrt{t_1}}{\tilde{S}_Ad^2} \sum_{i=1}^d \sum_{j=i}^d \tilde{S}_i \tilde{S}_j (A_{j-1}Z - \sigma t_j) e^{rt_1} - \frac{\tilde{S}_Ae^{-rT}}{\sigma} e^{rt_1} \right] \tilde{\Phi}_2 + \left(\frac{2e^{-rT}\tilde{S}_d\sqrt{t_1}}{\sigma^2T} - \frac{2e^{-rT}\tilde{S}_A}{\sigma^2} \sqrt{t_1} (r - \sigma^2) \right. \\ & - \frac{2e^{-rT}\sqrt{t_1}}{\tilde{S}_Ad^2} \sum_{i=1}^d \sum_{j=i}^d \tilde{S}_i \tilde{S}_j \right) \frac{1}{\sqrt{2\pi}} e^{\sigma^2t/2 + \omega t - (\psi_d - \sigma\sqrt{t_1})^2/2}, \end{split}$$

where $\Phi_1 := \Phi(-\psi_d)$, $\Phi_2 := \Phi(-\sigma\sqrt{t_1} - \psi_d)$, and $\tilde{\Phi}_2 := \Phi(\sigma\sqrt{t_1} - \psi_d)$; and $G_{\Delta,1}$, $G_{\Gamma,1}$ and $G_{\vartheta,1}$ are the CMV estimates for the binary Asian option Greeks in Theorem 6.2.

Note that in this case, the CMV estimates are the asymptotic unbiased estimates of the Greeks with respect to *d* according to Theorem 5.4. We can easily see that all estimates given above are infinitely times differentiable, which are QMC-friendly as well.

Example 6.5. Consider the up-and-out Asian call option with the following payoff function

$$f(\bar{S}_T) = (\bar{S}_T - K)^+ \mathbf{1} \{\bar{S}_T \le H\}.$$

Using the notations and imitating the steps in Example 6.1, we can obtain the CMV estimates for the up-and-out Asian call option Greeks in the following theorem.

Theorem 6.6. For the up-and-out Asian call option, the CMV estimates for delta, gamma and vega are given by

$$\begin{aligned} & \textit{delta}: G_{\Delta,3}(K,H; \boldsymbol{Z}) = G_{\Delta,2}(K; \boldsymbol{Z}) - (G_{\Delta,2}(H; \boldsymbol{Z}) + (H-K)G_{\Delta,1}(H; \boldsymbol{Z})); \\ & \textit{gamma}: G_{\Gamma,3}(K,H; \boldsymbol{Z}) = G_{\Gamma,2}(K; \boldsymbol{Z}) - (G_{\Gamma,2}(H; \boldsymbol{Z}) + (H-K)G_{\Gamma,1}(H; \boldsymbol{Z})); \\ & \textit{vega}: G_{\vartheta,3}(K,H; \boldsymbol{Z}) = G_{\vartheta,2}(K; \boldsymbol{Z}) - (G_{\vartheta,2}(H; \boldsymbol{Z}) + (H-K)G_{\vartheta,1}(H; \boldsymbol{Z})), \end{aligned}$$

where $G_{\Delta,1}$, $G_{\Gamma,1}$ and $G_{\vartheta,1}$ are the CMV estimates for the binary Asian option Greeks in Theorem 6.2, and $G_{\Delta,2}$, $G_{\Gamma,2}$ and $G_{\vartheta,2}$ are the CMV estimates for the Greeks in Theorem 6.4.

Again, we can easily see that all estimates given above are infinitely times differentiable.

7 Numerical experiments

In this section, we present numerical experiments for Examples 6.1-6.5 in Section 6 by comparing the variance reduction factors (VRFs) of different methods to illustrate the effectiveness of QMC-CMV method. We choose RQMC points instead of QMC points since the RQMC points are random which allow the variance to be estimated.

Let $\mathscr{P} = \{u_1, \cdots, u_N\}$ be the (d-1)-dimensional QMC points. We generate M independent random versions $\mathscr{P}_j = \{u_1^{(j)}, \cdots, u_N^{(j)}\}, j = 1, \cdots, M$. For each batch \mathscr{P}_j , we compute the estimate

$$Q_N^{(j)} = \frac{1}{N} \sum_{i=1}^N G(u_i^{(j)}),$$

Greeks	K	d	MC-MV	QMC-MV	MC-CMV	QMC-CMV
delta	90	64	1	11	1	55,545
		128	1	12	1	41,135
	100	64	1	11	1	35,126
		128	1	8	1	21,778
	110	64	1	8	1	38,228
		128	1	7	1	28,820
gamma	90	64	1	5	1	5,647
		128	1	3	1	4,113
	100	64	1	6	1	7,154
		128	1	3	1	7,508
	110	64	1	4	1	4,678
		128	1	3	1	3,565
vega	90	64	1	5	1	8,139
		128	1	4	1	5,742
	100	64	1	4	1	6,582
		128	1	4	1	6,738
	110	64	1	4	1	6,154
		128	1	3	1	6,415

where G(Z) is the payoff function. The ultimate estimate is given by

$$\tilde{Q} = \frac{1}{M} \sum_{j=1}^{M} Q_N^{(j)},$$

and the variance of the estimate \tilde{Q} is estimated by

$$\sigma_{\text{RQMC}}^2 = \frac{1}{M(M-1)} \sum_{j=1}^{M} (Q_N^{(j)} - \tilde{Q})^2.$$

In our experiments, the number of samples N and the number of batches M are the same for each method. We take H=120 in the up-and-out Asian call options. In each numerical experiment, there are M=500 independent batches of simulation and each batch contains $N=2^{15}$ samples. The Malliavin (MV) estimate in MC is chosen as the benchmark, and we denote it by MC-MV. Let the variance of the MC-MV be V_0^2 . If the variance of an alternative estimate is V^2 , then the VRF of this estimate with respect to MC-MV is given by

$$VRF := \frac{V_0^2}{V^2}.$$

In QMC setting, GPCA method is used as the PGM to reduce the effective dimension (due to its superiority in Section 4) and the scrambled Sobol' points proposed by Matoušek [19] are used as the RQMC points. We set parameters $S_0 = 100$, $\sigma = 0.2$, r = 0.1, T = 1 for all examples and set d = 64, 256, and K = 90, 100, 110. The option is in the money if K = 90, at the money if K = 100, and out of money if K = 110.

For each Greek, we compare the MV method and the CMV method. The MV method includes MC-MV and QMC-MV, which combine MV method with MC and QMC, respectively. The CMV method includes MC-CMV and QMC-CMV, which combine CMV method with MC and QMC, respectively.

In this paper, all the programs are run under macOS using MATLAB (version 8.4.0). The computer is a MacBook Pro laptop with Intel Core i5 Duo CPU @ 2.4 GHz, 8 GB of RAM.

The results for binary Asian options, Asian call options and up-and-out Asian call options are presented in Tables 1-9, respectively. We observe the following from the experiments.

Table 2: The Greek values (×10⁻³) for binary Asian options with payoff function $f = \mathbf{1}\{\bar{S}_T > K\}$.

Greeks	K	d	MC-MV	QMC-MV	MC-CMV	QMC-CMV
delta	90	64	14.083	14.053	14.071	14.056
		128	14.024	14.016	13.999	14.015
	100	64	29.204	29.188	29.187	29.185
		128	29.179	29.199	29.193	29.198
	110	64	27.802	27.802	27.812	27.805
		128	27.747	27.728	27.747	27.730
gamma	90	64	-1.7400	-1.7407	-1.7414	-1.7404
		128	-1.7216	-1.7195	-1.7184	-1.7185
	100	64	-1.1893	-1.1895	-1.1890	-1.1896
		128	-1.1690	-1.1693	-1.1688	-1.1713
	110	64	0.8000	0.8018	0.8035	0.8010
		128	0.8229	0.8208	0.8203	0.8196
vega	90	64	-1092.7	-1095.0	-1096.1	-1095.0
		128	-1107.2	-1109.5	-1111.2	-1110.3
	100	64	-831.14	-831.92	-833.07	-831.66
		128	-839.68	-840.38	-839.58	-840.47
	110	64	512.06	511.82	512.05	512.27
		128	510.34	509.42	511.49	510.53

Table 3: The simulation times (unit: s) for binary Asian options with payoff function $f = \mathbf{1}\{\bar{S}_T > K\}$.

Greeks	K	d	MC-MV	QMC-MV	MC-CMV	QMC-CMV
delta	90	64	157	172	123	152
		128	301	356	241	302
	100	64	147	179	117	150
		128	309	369	247	312
	110	64	156	187	124	153
		128	305	370	252	329
gamma	90	64	153	191	132	156
		128	304	387	267	348
	100	64	176	202	132	173
		128	337	415	287	360
	110	64	176	200	139	175
		128	344	410	290	354
vega	90	64	968	949	1601	1794
		128	3170	3884	7197	7091
	100	64	930	919	1517	1549
		128	3074	3625	7190	6990
	110	64	994	1022	1615	1644
		128	3319	3426	6897	6951

Table 4: The VRFs for Asian call options with payoff function $f = (\bar{S}_T - K)^+$.

Greeks	K	d	MC-MV	QMC-MV	MC-CMV	QMC-CMV
delta	90	64	1	50	1	12,804
		128	1	22	1	13,396
	100	64	1	17	1	5,809
		128	1	12	1	6,695
	110	64	1	9	1	5,007
		128	1	7	1	4,987
gamma	90	64	1	5	1	919
		128	1	3	1	1,132
	100	64	1	3	1	727
		128	1	3	1	739
	110	64	1	3	1	481
		128	1	2	1	494
vega	90	64	1	4	1	754
		128	1	4	1	724
	100	64	1	3	1	537
		128	1	3	1	643
	110	64	1	3	1	437
		128	1	2	1	349

Table 5: The Greek values (×10⁻³) for Asian call options with payoff function $f = (\bar{S}_T - K)^+$.

Greeks	K	d	MC-MV	QMC-MV	MC-CMV	QMC-CMV
delta	90	64	884.51	883.76	884.44	883.83
		128	879.06	879.46	879.72	879.28
	100	64	659.73	660.31	659.90	660.19
		128	655.69	655.75	655.53	655.78
	110	64	359.25	359.40	359.35	359.42
		128	355.36	355.11	355.13	355.18
gamma	90	64	12.113	12.196	12.101	12.157
		128	12.500	12.351	12.349	12.365
	100	64	29.165	29.220	29.258	29.160
		128	29.218	29.117	29.130	29.187
	110	64	30.802	30.821	30.906	30.773
		128	30.589	30.590	30.579	30.596
vega	90	64	9119.8	9090.1	9005.7	9069.4
		128	8552.1	8530.9	8638.5	8563.6
	100	64	20364	20397	20346	20379
		128	19939	19921	19874	19922
	110	64	21823	21773	21820	21796
		128	21497	21480	21470	21458

Table 6: The simulation times (unit: s) for Asian call options with payoff function $f = (\bar{S}_T - K)^+$.

Greeks	K	d	MC-MV	QMC-MV	MC-CMV	QMC-CMV
delta	90	64	143	168	119	150
		128	286	376	243	316
	100	64	152	173	119	150
		128	291	379	245	324
	110	64	152	182	125	164
		128	301	386	259	347
gamma	90	64	159	198	142	177
		128	313	392	266	334
	100	64	153	187	137	172
		128	299	375	254	321
	110	64	147	183	131	163
		128	292	354	249	326
vega	90	64	990	1101	1772	1762
		128	3412	3415	6808	6862
	100	64	987	998	1704	1734
		128	3315	3384	6813	7045
	110	64	1106	1182	2081	1739
		128	3340	3358	6185	6827

Table 7: The VRFs for up-and-out Asian call options with payoff function $f = (\bar{S}_T - K)^+ \mathbf{1} \{\bar{S}_T \le H\}$, where H = 120.

Greeks	K	d	MC-MV	QMC-MV	MC-CMV	QMC-CMV
delta	90	64	1	2	1	28,707
		128	1	2	1	20,327
	100	64	1	2	2	51,475
		128	1	2	1	38,146
	110	64	1	1	2	67,723
		128	1	1	1	37,032
gamma	90	64	1	2	1	1,355
		128	1	1	1	1,052
	100	64	1	2	1	1,838
		128	1	2	1	2,235
	110	64	1	1	2	3,032
		128	1	1	2	2,369
vega	90	64	1	1	1	18,247
		128	1	2	1	11,787
	100	64	1	1	2	18,495
		128	1	1	1	11,064
	110	64	1	1	2	12,718
		128	1	1	1	8,812

Table 8: The Greek values (×10⁻³) for up-and-out Asian call options with payoff function $f = (\bar{S}_T - K)^+ \mathbf{1} \{\bar{S}_T \le H\}$, where H = 120.

Greeks	K	d	MC-MV	QMC-MV	MC-CMV	QMC-CMV
delta	90	64	283.52	283.79	283.73	283.64
		128	287.00	286.92	287.00	287.03
	100	64	212.65	212.83	212.77	212.73
		128	214.62	214.60	214.70	214.58
	110	64	64.706	64.644	64.684	64.689
		128	64.987	65.069	65.005	65.043
gamma	90	64	-47.024	-47.043	-47.057	-47.035
		128	-46.568	-46.552	-46.511	-46.533
	100	64	-16.453	-16.452	-16.468	-16.454
		128	-16.129	-16.147	-16.165	-16.144
	110	64	-1.2664	-1.2597	-1.2706	-1.2707
		128	-1.1690	-1.1668	-1.1666	-1.1675
vega	90	64	-32905	-32884	-32910	-32895
		128	-32828	-32857	-32844	-32850
	100	64	-12151	-12151	-12154	-12156
		128	-12068	-12062	-12080	-12071
	110	64	-1152.0	-1151.6	1152.5	-1151.1
		128	-1111.0	-1111.1	-1112.2	-1110.5

Table 9: The simulation times (unit: s) for up-and-out Asian call options with payoff function $f = (\bar{S}_T - K)^+ \mathbf{1} \{\bar{S}_T \le H\}$, where H = 120.

Greeks	K	d	MC-MV	QMC-MV	MC-CMV	QMC-CMV
delta	90	64	144	167	128	153
		128	291	350	248	335
	100	64	150	179	125	152
		128	285	383	249	327
	110	64	151	210	146	180
		128	333	401	251	315
gamma	90	64	148	202	170	173
		128	309	340	261	326
	100	64	151	197	143	186
		128	289	356	288	338
	110	64	150	179	141	171
		128	303	358	264	335
vega	90	64	1034	1053	1814	1926
		128	3364	3541	7604	7482
	100	64	981	995	1745	1777
		128	3404	3458	6985	7047
	110	64	1017	1057	1760	1788
		128	3506	3522	7635	7352

- Among all methods, QMC-CMV method is the most effective method to calculate Greeks. This is because
 QMC-CMV method improves the smoothness of the integrand, which is very significant in enhancing the computational efficiency of the QMC method. Furthermore, due to the improved smoothness of the integrands,
 GPCA method can be applied to reduce the effective dimension and increase the efficiency of the QMC method.
- QMC-CMV behaves much better than QMC-MV. However, in MC setting, the VRFs of MC-CMV are nearly the
 same as the VRFs of MC-MV. This is because MC method is not sensitive to the smoothness of the integrand.
 Moreover, CMV and MV in QMC setting behave better than those in MC setting, respectively. However, the
 VRFs of MV in QMC are no more than one hundred, while the VRFs of QMC-CMV can be larger than tens of
 thousands in some cases. This is because the integrands in CMV have good smoothness.
- The VRFs of QMC-CMV in binary Asian option and up-and-out Asian call option (Tables 1 and 7) are more impressive than those in Asian call option (Table 4). The reason is that the payoff function of Asian call option is continuous, whereas the payoff functions of binary Asian option and up-and-out Asian call option are not. Thus the effect of smoothing for binary Asian option and up-and-out Asian option is more significant than that for the Asian call option.
- For *delta* of all options (Tables 1, 4 and 7), the superiority of the QMC-CMV method is more impressive. The VRFs of QMC-CMV are larger than tens of thousands in most cases. For *gamma* of all options (Tables 1, 4 and 7) and *vega* of binary Asian options and Asian call options (Tables 1 and 4), the VRFs of QMC-CMV are no more than ten thousands. However, for *vega* of up-and-out Asian call options (Table 7), the VRFs of QMC-CMV are larger than tens of thousands in most cases. A critical factor that leads to the above results is the different weight functions in different Greeks formulae (12).
- Dimensionality and strike price have some impact on the VRFs. For binary Asian options (Table 1), for example, the VRFs of QMC-CMV for *delta* are smaller when the dimension increases. For Asian call options (Table 4), for example, the VRFs of QMC-CMV are smaller when the strike price is larger for all three Greeks. Generally, the high dimensionality has little impact on all methods in QMC setting since the GPCA method reduces the effective dimension, which lifts the curse of dimensionality.
- From Tables 2, 5 and 8, we can see that there is no significant difference in Greek values between those methods. This implies the consistency of our estimates.
- From Tables 3, 6 and 9, we can see that there is no significant difference in simulation times between those methods when considering *delta* and *gamma*. However, for *vega*, the simulation time of CMV is nearly twice as long as that of MV. A reasonable explanation is that the weight functions of *vega* of the CMV estimates are much more complicated (see Theorem 6.2, 6.4 and 6.6).

Remark 7.1. We only consider the one-dimensional simple Asian options as illustrations in Sections 6 and 7 for convenience. However, the QMC-CMV method can also be implemented in the case of complex Asian options and multiple assets. Due to the complicated calculation, the results of those case are not presented in this paper. Moreover, since the weight functions of simple Asian option Greeks include S_T as well (see Corollary 3.4) and there is no essential difference between the weight functions of simple and complex Asian options, the efficiency of the QMC-CMV in Section 7 still holds when we deal with the complex Asian options.

8 Conclusion

In this paper, we develop an integration by parts formula in the multi-dimensional Malliavin calculus and propose a new method to calculate the Greeks for Asian options. We obtain the Greeks formulae for both simple and complex continuous-time Asian options in the multi-asset Black-Scholes model by the integration by parts formula. By discretization we derive the simulation estimates of the Asian option Greeks. We prove the asymptotic convergence of the simulation estimates when the payoff function is nonnegative, continuous with linear growth. After then, by taking conditional expectations, we get the CMV estimates for Greeks. We show how to calculate the conditional expectations analytically for binary Asian options, Asian call options and up-and-out Asian call options. We find that the smoothness of the CMV estimate is enhanced, leading to a notable efficiency improvement in QMC. Even for options with continuous payoffs, taking conditional expectations is of great significance because the CMV estimates

are smoothed. The numerical results show that, for binary Asian options, Asian call options and up-and-out Asian call options, the VRFs of QMC-CMV methods can be larger than tens of thousands in some cases compared with the MC-MV method.

For further work, the bias of the simulation estimates for Asian option Greeks in general cases need to be studied. More general formulae for CMV estimates of Asian option Greeks are also expected to be developed. Moreover, extending the QMC-CMV method for Greeks to other models is a subject of ongoing research.

acknowledgements

The authors are very grateful to the editors and the anonymous referee for their helpful suggestions and comments. The work is funded by the National Natural Science Foundation of China (No. 72071119).

Conflict of interest

The authors declare that they have no conflict of interest. All authors certify that they have no affiliations with or involvement in any organization or entity with any financial interest or non-financial interest in the subject matter or materials discussed in this manuscript.

References

- [1] Bayer, C., Siebenmorgen, M. and Tempone, R. (2018). Smoothing the payoff for efficient computation of basket option prices. *Quantitative Finance*, **18**, 491–505.
- [2] Benhamou, E. (2000). An application of Malliavin calculus to continuous time Asian Option Greeks. Technical report, London School of Economics.
- [3] Broadie, M. and Glasserman, P. (1996). Estimating security price derivatives using simulation. *Management Science*, **42**, 269–285.
- [4] Caflisch, R.E., Morokoff, W.J. and Owen, A.B. (1997). Valuation of mortgage-backed securities using Brownian bridges to reduce effective dimension. *Journal of Computational Finance*, **1**, 27–46.
- [5] Di Nunno, G., Øksendal, B. and Proske, F. (2009). Malliavin Calculus for Lévy Processes with Applications to Finance, Springer Verlag, Berlin.
- [6] Faure, H. (1982). Discrépance de suites associées à un système de numération (en dimension s). *Acta Arithmetica*, **41**, 337–351.
- [7] Fournié, E., Lasry, J., Lebuchoux, J., Lions, P.L. and Touzi, N. (1999). Applications of Malliavin calculus to Monte Carlo method in finance I. *Finance and Stochastics*, **3**, 391–412.
- [8] Fournié, E., Lasry, J., Lebuchoux, J. and Lions, P.L. (2001). Applications of Malliavin calculus to Monte Carlo method in finance II. *Finance and Stochastics*, **5**, 201–236.
- [9] Fu, M.C. and Hu, J.Q. (1997). Conditional Monte Carlo: Gradient Estimation and Optimization Applications, Kluwer Academic, Norwell, MA.
- [10] Glasserman, P. (2004). Monte Carlo Methods in Financial Engineering, Springer Verlag, New York.
- [11] Glynn, P.W. (1987). Likelihood ratio gradient estimation: An overview, In *Proceedings of the 19th Conference on Winter Simulation*, IEEE Press, New York, pp. 366–375.
- [12] Halton, J.H. (1960). On the efficiency of certain quasi-random sequences of points in evaluating multi-dimensional integrals. *Numerische Mathematik*, **2**, 84–90.
- [13] He, Z. (2019). On the error rate of conditional quasi-Monte Carlo for discontinuous functions. *SIAM Journal on Numerical Analysis*, **57**, 854–874.

- [14] He, Z. and Wang, X. (2014). Good path generation methods in quasi-Monte Carlo for pricing financial derivatives. *SIAM Journal on Scientific Computing*, **36**, B171–B197.
- [15] Ho, Y.C. and Cao, X. (1983). Perturbation analysis and optimization of queueing networks. *Journal of Optimization Theory and Applications*, **40**, 559–582.
- [16] Imai, J. and Tan, K.S. (2004). Minimizing effective dimension using linear transformation. In *Monte Carlo and Quasi-Monte Carlo Methods* 2002, pp. 275-292, Springer, Berlin, Heidelberg.
- [17] Karatzas, I. and Shreve, S. (1991). Brownian Motion and Stochastic Calculus, Springer Verlag, New York.
- [18] L'Ecuyer, P. and Lemieux, C. (2005). Recent advances in randomized quasi-Monte Carlo methods, In *Proceedings of the Modeling Uncertainty: An Examination of Stochastic Theory, Methods, and Applications*, Springer, Boston, MA, pp. 419-474.
- [19] Matoušek, J. (1998). On the L_2 -discrepancy for anchored boxes. *Journal of Complexity*, **14**, 527–556.
- [20] Morokoff, W.J. and Caflisch, R.E. (1995). Quasi-Monte Carlo integration. *Journal of Computational Physics*, **122**, 218–230.
- [21] Niederreiter, H. (1992). Random Number Generation and Quasi-Monte Carlo Methods, SIAM, Philadelphia.
- [22] Nualart, D. (2006). The Malliavin Calculus and Related Topics (2nd ed.), Springer Verlag, Berlin Heidelberg.
- [23] Ocone, D.L. and Karatzas, I. (1991). A generalized clark representation formula, with application to optimal portfolios. *Stochastics: An International Journal of Probability and Stochastic Processes*, **34**, 187–220.
- [24] Owen, A.B. (1998). Scrambling Sobol' and Niederreiter-Xing points. Journal of Complexity, 14, 466-489.
- [25] Sobol, I.M. (1967). On the distribution of points in a cube and the approximate evaluation of integrals. *USSR Computational Mathematics and Mathematical Physics*, **7**, 86–112.
- [26] Wang, X. (2006). On the effects of dimension reduction techniques on some high-dimensional problems in finance. *Operations Research*, **54**, 1063–1078.
- [27] Wang, X. and Fang, K.T. (2003). The effective dimension and quasi-Monte Carlo integration. *Journal of Complexity*, **19**, 101–124.
- [28] Wang, X. and Tan, K.S. (2013). Pricing and hedging with discontinuous functions: Quasi-Monte Carlo methods and dimension reduction. *Management Science*, **59**, 376–389.
- [29] Weng, C., Wang, X. and He, Z. (2017). Efficient computation of option prices and Greeks by quasi-Monte Carlo method with smoothing and dimension reduction. *SIAM Journal on Scientific Computing*, **39**, B298–B322.
- [30] Xiao, Y. and Wang, X. (2018). Conditional quasi-Monte Carlo methods and dimension reduction for option pricing and hedging with discontinuous functions. *Journal of Computational and Applied Mathematics*, 343, 289–308.
- [31] Xiao, Y. and Wang, X. (2019). Enhancing quasi-Monte Carlo simulation by minimizing effective dimension for derivative pricing. *Computational Economics*, **54**, 343–366.
- [32] Xu, Y., Lai, Y. and Yao, H. (2014). Efficient simulation of Greeks of multiasset European and Asian style options by Malliavin calculus and quasi-Monte Carlo methods. *Applied Mathematics and Computation*, **236**, 493–511.
- [33] Zhang, C. and Wang, X. (2020). Quasi-Monte Carlo-based conditional pathwise method for option Greeks. *Quantitative Finance*, **20**, 49–67.