

Johns Hopkins Engineering

625.464 Computational Statistics

Introduction to Function Estimation
Pointwise and Global Properties of Estimators

Module 10 Lecture 10B



JOHNS HOPKINS
WHITING SCHOOL
of ENGINEERING

Pointwise Properties of Function Estimators

Assume \hat{f} estimates $f: D \rightarrow \mathbb{R}$

① Bias

The bias of \hat{f} at x is given by

$$\text{Bias}(\hat{f}(x)) = E[\hat{f}(x)] - f(x)$$

If $\text{Bias}(\hat{f}(x)) = 0$, $\Rightarrow \hat{f}$ is unbiased at x

If \hat{f} is unbiased at every point x in D ,
then we say that the estimator is
pointwise unbiased.

Pointwise Properties of Functions Estimators

② Variance

The variance of \hat{f} at x is given by

$$V[\hat{f}(x)] = E[(\hat{f}(x) - E[\hat{f}(x)])^2]$$

③ Mean Squared Error (MSE)

The MSE of \hat{f} at x is

$$\begin{aligned} \text{MSE}[\hat{f}(x)] &= E[(\hat{f}(x) - f(x))^2] \\ &= V(\hat{f}(x)) + (\text{Bias}(\hat{f}(x)))^2 \end{aligned}$$

Pointwise Properties of Functions Estimators

④ Mean Absolute Error (MAE)

the MAE of \hat{f} at x is given by

$$MAE[\hat{f}(x)] = E(|\hat{f}(x) - f(x)|)$$

⑤ Consistency

\hat{f} is said to be pointwise consistent if

$$E[\hat{f}(x)] \rightarrow f(x)$$

for each x as $n \rightarrow \infty$

Global Properties of Estimators of Functions

- statistical properties of \hat{f} over all of domain D of f .
- written w/o indication on x
- often integration of pointwise prop or defined in terms of norms.

The main tool for comparing \hat{f} & f is

$$\|\hat{f} - f\|_p = \left(\int_D |\hat{f}(x) - f(x)|^p dx \right)^{1/p}$$

Need

- ① \hat{f} defined over all of D ② integral exists

Global Properties of Estimators of Functions

- L_1 : Integrated absolute error (IAE)

$$IAE(\hat{f}) = \int_D |\hat{f}(x) - f(x)| dx$$

- L_2 : Integrated squared error (ISE)

$$ISE(\hat{f}) = \int_D (\hat{f}(x) - f(x))^2 dx$$

- L_∞ : Sup absolute error (SAE)

$$SAE(\hat{f}) = \sup_{x \in D} |\hat{f}(x) - f(x)|$$

Global Properties of Estimators of Functions

Bias

Integrated Absolute Bias (IAB)

$$IAB(\hat{f}) = \int_D |E(\hat{f}(x)) - f(x)| dx$$

Integrated Squared Bias (ISB)

$$ISB(\hat{f}) = \int_D (E(\hat{f}(x)) - f(x))^2 dx$$

If \hat{f} is unbiased then

$$IAB(\hat{f}) = ISB(\hat{f}) = 0$$

$Bias(\hat{f}(x)) = 0$ almost everywhere.

Global Properties of Estimators of Functions

Variance

Integrated Variance (IV)

$$IV(\hat{f}) = \int_D V(\hat{f}(x)) = \int_D E[(\hat{f}(x) - E[\hat{f}(x)])^2] dx$$

Integrated Mean Squared Error (IMSE)

$$IMSE(\hat{f}) = \int_D E(f(x) - \hat{f}(x))^2 dx$$

$$= IV(\hat{f}) + ISB(\hat{f})$$

$$= E\left[\int_D (\hat{f}(x) - f(x))^2 dx\right] = E[IMSE(\hat{f})] = mISE(\hat{f})$$

Global Properties of Estimators of Functions

Integrated Mean Absolute Error (IMAE)

$$\text{IMAE}(\hat{f}) = \int_D E(|\hat{f}(x) - f(x)|)$$

$$= E\left[\int_D |\hat{f}(x) - f(x)| dx\right] = \text{MIAE}(\hat{f})$$

— can also extend the idea of consistency. — See Gentle Ch-10.