

RWorksheet_calatana#3B.Rmd

austine john calatana

2023-10-15

#NUMBER 1 #A.

```
household <- data.frame (  
  Respondents = 1:20,  
  Sex = c(2, 2, 1, 2, 2, 2, 2, 2, 2, 2, 1, 2, 2, 2, 2, 2, 2, 1, 2),  
  Fathers_Occupation = c(1, 3, 3, 3, 1, 2, 3, 1, 1, 1, 3, 2, 1, 3, 3, 1, 3, 1, 2, 1),  
  Person_at_Home = c(5, 7, 3, 8, 5, 9, 6, 7, 8, 4, 7, 5, 4, 7, 8, 8, 3, 11, 7, 6),  
  Siblings_at_school = c(6, 4, 4, 1, 2, 1, 5, 3, 1, 2, 3, 2, 5, 5, 2, 1, 2, 5, 3, 2),  
  Types_of_houses = c(1, 2, 3, 1, 1, 3, 3, 1, 2, 3, 2, 3, 2, 2, 3, 3, 3, 3, 3, 2)  
)  
household
```

##	Respondents	Sex	Fathers_Occupation	Person_at_Home	Siblings_at_school
## 1	1	2	1	5	6
## 2	2	2	3	7	4
## 3	3	1	3	3	4
## 4	4	2	3	8	1
## 5	5	2	1	5	2
## 6	6	2	2	9	1
## 7	7	2	3	6	5
## 8	8	2	1	7	3
## 9	9	2	1	8	1
## 10	10	2	1	4	2
## 11	11	1	3	7	3
## 12	12	2	2	5	2
## 13	13	2	1	4	5
## 14	14	2	3	7	5
## 15	15	2	3	8	2
## 16	16	2	1	8	1
## 17	17	2	3	3	2
## 18	18	2	1	11	5
## 19	19	1	2	7	3
## 20	20	2	1	6	2
##	Types_of_houses				
## 1	1				
## 2	2				
## 3	3				
## 4	1				
## 5	1				
## 6	3				
## 7	3				
## 8	1				
## 9	2				

```
## 10      3
## 11      2
## 12      3
## 13      2
## 14      2
## 15      3
## 16      3
## 17      3
## 18      3
## 19      3
## 20      2
```

#B.

```
str(household)
```

```
## 'data.frame': 20 obs. of 6 variables:
## $ Respondents : int 1 2 3 4 5 6 7 8 9 10 ...
## $ Sex : num 2 2 1 2 2 2 2 2 2 2 ...
## $ Fathers_Occupation: num 1 3 3 3 1 2 3 1 1 1 ...
## $ Person_at_Home : num 5 7 3 8 5 9 6 7 8 4 ...
## $ Siblings_at_school: num 6 4 4 1 2 1 5 3 1 2 ...
## $ Types_of_houses : num 1 2 3 1 1 3 3 1 2 3 ...
```

```
summary(household)
```

```
## Respondents Sex Fathers_Occupation Person_at_Home
## Min. : 1.00 Min. :1.00 Min. :1.00 Min. : 3.0
## 1st Qu.: 5.75 1st Qu.:2.00 1st Qu.:1.00 1st Qu.: 5.0
## Median :10.50 Median :2.00 Median :2.00 Median : 7.0
## Mean :10.50 Mean :1.85 Mean :1.95 Mean : 6.4
## 3rd Qu.:15.25 3rd Qu.:2.00 3rd Qu.:3.00 3rd Qu.: 8.0
## Max. :20.00 Max. :2.00 Max. :3.00 Max. :11.0
## Siblings_at_school Types_of_houses
## Min. :1.00 Min. :1.0
## 1st Qu.:2.00 1st Qu.:2.0
## Median :2.50 Median :2.5
## Mean :2.95 Mean :2.3
## 3rd Qu.:4.25 3rd Qu.:3.0
## Max. :6.00 Max. :3.0
```

#C.

```
mean_siblings <- mean(household$Siblings_at_school)
mean_siblings == 5
```

```
## [1] FALSE
```

#D.

```
subset1 <- household[1:2, ]
subset1
```

```
## Respondents Sex Fathers_Occupation Person_at_Home Siblings_at_school
## 1      1 2      1      5      6
## 2      2 2      3      7      4
## Types_of_houses
## 1      1
## 2      2
```

#E.

```
subset2 <- household[c(3, 5), c(2, 4)]
subset2
```

```
##    Sex Person_at_Home
## 3    1                3
## 5    2                5
```

#F.

```
types_houses <- household$Types_of_houses
types_houses
```

```
## [1] 1 2 3 1 1 3 3 1 2 3 2 3 2 2 3 3 3 3 3 2
```

#G.

```
male_farmer <- subset(household, Sex == 1 & Fathers_Occupation == 1)
male_farmer
```

```
## [1] Respondents      Sex      Fathers_Occupation Person_at_Home
## [5] Siblings_at_school Types_of_houses
## <0 rows> (or 0-length row.names)
```

#H.

```
female_greater_than_5_siblings <- subset(household, Sex == 2 & Siblings_at_school >= 5)
female_greater_than_5_siblings
```

```
##    Respondents Sex Fathers_Occupation Person_at_Home Siblings_at_school
## 1              1  2                  1              5              6
## 7              7  2                  3              6              5
## 13             13  2                  1              4              5
## 14             14  2                  3              7              5
## 18             18  2                  1             11              5
##    Types_of_houses
## 1                  1
## 7                  3
## 13                 2
## 14                 2
## 18                 3
```

#NUMBER 2

```
df <- data.frame(
  Ints = integer(),
  Doubles = double(),
  Characters = character(),
  Logicals = logical(),
  Factors = factor(),
  stringsAsFactors = FALSE
)
cat("Structure of the empty dataframe:\n")
```

```
## Structure of the empty dataframe:
```

```
str(df)
```

```
## 'data.frame':    0 obs. of  5 variables:
## $ Ints      : int
```

```
## $ Doubles : num
## $ Characters: chr
## $ Logicals : logi
## $ Factors : Factor w/ 0 levels:
```

#Output The output shows that the data frame has 0 observations (rows) and 5 variables (columns) with their respective data types. The “Factors” column is empty since there are no levels defined yet.

#NUMBER 3

```
household_data <- data.frame (
  Respondents = 1:10,
  Sex = c("Male", "Female", "Female", "Male", "Male", "Female", "Female", "Male", "Female", "Male"),
  Fathers_Occupation = c(1,2,3,3,1,2,2,3,1,3),
  Person_at_Home = c(5,7,3,8,6,4,4,2,11,6),
  Siblings_at_school = c(2,3,0,5,2,3,1,2,6,2),
  Types_of_houses = c("Wood", "Congrete", "Congrete", "Wood", "Semi-Congrete", "Semi-Congrete", "Wood",
)
household_data
```

```
## Respondents Sex Fathers_Occupation Person_at_Home Siblings_at_school
## 1 1 Male 1 5 2
## 2 2 Female 2 7 3
## 3 3 Female 3 3 0
## 4 4 Male 3 8 5
## 5 5 Male 1 6 2
## 6 6 Female 2 4 3
## 7 7 Female 2 4 1
## 8 8 Male 3 2 2
## 9 9 Female 1 11 6
## 10 10 Male 3 6 2
## Types_of_houses
## 1 Wood
## 2 Congrete
## 3 Congrete
## 4 Wood
## 5 Semi-Congrete
## 6 Semi-Congrete
## 7 Wood
## 8 Semi-Congrete
## 9 Semi-Congrete
## 10 Congrete
```

```
write.csv(household_data, file = "HouseholdData.csv", row.names = FALSE)
```

#A.

```
household_data <- read.csv("HouseholdData.csv")
```

#B.

```
household_data$Sex <- factor(household_data$Sex)
household_data$Sex <- as.integer(factor(household_data$Sex,
  levels = c("Male", "Female"),
  labels = c(1, 2)))
household_data
```

```
## Respondents Sex Fathers_Occupation Person_at_Home Siblings_at_school
```

```
## 1      1  1      1      5      2
## 2      2  2      2      7      3
## 3      3  2      3      3      0
## 4      4  1      3      8      5
## 5      5  1      1      6      2
## 6      6  2      2      4      3
## 7      7  2      2      4      1
## 8      8  1      3      2      2
## 9      9  2      1     11      6
## 10     10  1      3      6      2
##      Types_of_houses
## 1      Wood
## 2      Congrete
## 3      Congrete
## 4      Wood
## 5      Semi-Congrete
## 6      Semi-Congrete
## 7      Wood
## 8      Semi-Congrete
## 9      Semi-Congrete
## 10     Congrete
```

#C.

```
household_data$Types_of_houses <- factor(household_data$Types_of_houses)
household_data$Types_of_houses <- as.integer(factor(household_data$Types_of_houses,
                                                    levels = c("Wood", "Congrete", "Semi-Congrete"),
                                                    labels = c(1, 2, 3)))
print(household_data)
```

```
##      Respondents Sex Fathers_Occupation Person_at_Home Siblings_at_school
## 1      1  1      1      5      2
## 2      2  2      2      7      3
## 3      3  2      3      3      0
## 4      4  1      3      8      5
## 5      5  1      1      6      2
## 6      6  2      2      4      3
## 7      7  2      2      4      1
## 8      8  1      3      2      2
## 9      9  2      1     11      6
## 10     10  1      3      6      2
##      Types_of_houses
## 1      1
## 2      2
## 3      2
## 4      1
## 5      3
## 6      3
## 7      1
## 8      3
## 9      3
## 10     2
```

#D.

```
household_data$Fathers_Occupation <- factor(household_data$Fathers_Occupation)
household_data$Fathers_Occupation <- as.character(factor(household_data$Fathers_Occupation,

                                                         levels = c(1, 2, 3),
                                                         labels = c("Farmer", "Driver", "Others"))))

# Print the updated data frame
print(household_data)
```

```
## Respondents Sex Fathers_Occupation Person_at_Home Siblings_at_school
## 1          1  1          Farmer              5              2
## 2          2  2          Driver              7              3
## 3          3  2          Others              3              0
## 4          4  1          Others              8              5
## 5          5  1          Farmer              6              2
## 6          6  2          Driver              4              3
## 7          7  2          Driver              4              1
## 8          8  1          Others              2              2
## 9          9  2          Farmer             11              6
## 10         10  1          Others              6              2
```

```
## Types_of_houses
## 1          1
## 2          2
## 3          2
## 4          1
## 5          3
## 6          3
## 7          1
## 8          3
## 9          3
## 10         2
```

#E.

```
female_driver <- subset(household_data, Sex == 2 & Fathers_Occupation == "Driver")
female_driver
```

```
## Respondents Sex Fathers_Occupation Person_at_Home Siblings_at_school
## 2          2  2          Driver              7              3
## 6          6  2          Driver              4              3
## 7          7  2          Driver              4              1
```

```
## Types_of_houses
## 2          2
## 6          3
## 7          1
```

#F.

```
greater_than_5_siblings <- subset(household_data, Siblings_at_school >= 5)
greater_than_5_siblings
```

```
## Respondents Sex Fathers_Occupation Person_at_Home Siblings_at_school
## 4          4  1          Others              8              5
## 9          9  2          Farmer             11              6
```

```
## Types_of_houses
## 4          1
```

#4. INTERPRET THE GRAPH The graph you've described appears to be a sentiment analysis over time on Twitter data, specifically for the dates July 14, 15, 17, 18, 20, and 21 in the year 2020. The sentiment analysis categorizes tweets into three sentiment groups: negative, neutral, and positive. Here's an interpretation of the data:

- July 14, 2020:
 - Negative count: Nearly 2,500 tweets were categorized as negative.
 - Neutral count: Around 1,500 tweets were categorized as neutral.
 - Positive count: Approximately 1,750 tweets were categorized as positive.
- July 15, 2020:
 - Negative count: Over 4,000 tweets were categorized as negative.
 - Neutral count: About 2,750 tweets were categorized as neutral.
 - Positive count: Roughly 3,200 tweets were categorized as positive.
- July 17, 2020:
 - Negative count: There were approximately 3,250 tweets categorized as negative.
 - Neutral count: Around 1,800 tweets were categorized as neutral.
 - Positive count: Almost 2,500 tweets were categorized as positive.
- July 18, 2020:
 - Negative count: There were still around 3,250 tweets categorized as negative.
 - Neutral count: About 2,000 tweets were categorized as neutral.
 - Positive count: Approximately 2,500 tweets were categorized as positive.
- July 20, 2020:
 - Negative count: There were nearly 2,500 tweets categorized as negative.
 - Neutral count: Around 1,500 tweets were categorized as neutral.
 - Positive count: Almost 1,750 tweets were categorized as positive.
- July 21, 2020:
 - Negative count: Around 4,000 tweets were categorized as negative.
 - Neutral count: About 2,600 tweets were categorized as neutral.
 - Positive count: Roughly 3,300 tweets were categorized as positive.

From this data, it seems that there were fluctuations in sentiment over the specified dates. July 15th and July 21st had higher numbers of negative and positive tweets compared to the other dates, indicating a potential increase in sentiment-related Twitter activity on those days. July 14th and July 20th had lower counts across all sentiment categories. The data suggests a varying sentiment landscape on Twitter during this period in 2020