



杭州电子科技大学

HANGZHOU DIANZI UNIVERSITY

Online Judge



Online Judge	Online Exercise	Online Teaching	Online Contests	Exercise Author
F.A.Q Hand In Hand Online Acmers Forum Discuss Statistical Charts	Problem Archive Realtime Judge Status Authors Ranklist <div><div></div><div>Search</div></div>	C/C++/Java Exams ACM Steps Go to Job Contest LiveCast ICPC@China	Best Coder <small>beta</small> VIP STD Contests Virtual Contests DIY Web-DIY <small>beta</small> Recent Contests	<div> 用QQ帐号登录</div> <div><div>Author ID</div><div>Password</div><div><div>Sign In</div><div>Register new ID</div></div></div>

Rikka with Number

Time Limit: 8000/4000 MS (Java/Others) Memory Limit: 65536/65536 K (Java/Others)
Total Submission(s): 272 Accepted Submission(s): 76

Problem Description

As we know, Rikka is poor at math. Yuta is worrying about this situation, so he gives Rikka some math tasks to practice. There is one of them:

In radix d , a number $K = (A_1A_2 \dots A_m)_d (A_i \in [0, d), A_1 \neq 0)$ is good if and only $A_1 - A_m$ is a permutation of numbers from 0 to $d - 1$.

A number K is good if and only if there exists at least one $d \geq 2$ and K is good under radix d .

Now, Yuta wants to calculate the number of good numbers in interval $[L, R]$

It is too difficult for Rikka. Can you help her?

Input

The first line contains a number $t(1 \leq t \leq 20)$, the number of the testcases.

For each testcase, the first line contains two decimal numbers $L, R(1 \leq L \leq R \leq 10^{5000})$.

Output

For each testcase, print a single line with a single number -- the answer modulo 998244353.

Sample Input

```
2
5 20
123456 123456789
```

Sample Output

```
3
114480
```

Source

2017 Multi-University Training Contest - Team 5

Recommend

liuyiding

[Statistic](#) | [Submit](#) | [Discuss](#) | [Note](#)

