

Digit DP

Q1 How many numbers in $[a, b]$ are there, where the digit d occurs exactly k times.

Eg- $a = 7$ $b = 58$
 $d = 7$ $k = 1$

$[7, 58]$

ans = 6

7 17 27 37 47 57

Brute force: iterate using for loop from a to b . Check if the number is valid

for ($i = a$; $i \leq b$; $i++$)

{
}
}

y

Iterations of for loop = $b - a + 1$

Now imagine $a = 100$
 $b = 10^{16}$

This will always give TLE

- We need a faster way of solving

ans $[a, b] =$

$0, 1, 2, \dots, a-1, \underline{a}, a+1, a+2, \dots, b$

$0, 1, 2, \dots, a-1,$

$[0, b] - [0, a-1]$

$0, 1, 2, \quad \underline{3, 4, 5, 6, 7, 8}$

$0, 1, 2$

- $[a, b] \Rightarrow [0, b] - [0, a-1]$

$[0, x]$

$\begin{array}{cccc} 5 & 2 & 3 & 4 \\ \hline 5 & - & - & - \end{array} \left. \begin{array}{l} \text{new-d} < \text{cur-d} \\ \text{new-d} > \text{cur-d} \\ \text{new-d} = \text{cur-d} \end{array} \right\} \begin{array}{l} \text{less} \\ \text{bigger} \\ \text{cant say} \end{array}$

Digit DP Build the number digit by digit satisfying all the conditions.

Let $x = \begin{array}{cccccc} & 0 & 1 & 2 & 3 & 4 \\ & 5 & 4 & 3 & 2 & 1 \end{array}$

$\begin{array}{cccccc} 5 & - & - & - & - & - \\ \hline \end{array}$

Info needed \Rightarrow

- 1) cur-digit-index
- 2) cur num vs given no
- 3) How many times digit d has been placed till now.

int dp[pos][is-smaller or bigger]
[count-of-d]

int solve (int pos, bool is-small,
int count-of-d) {

if (count-of-d > k)

return 0

if (pos == num.size()) {

if (count-of-d == k)

return 1

else

return 0

}

if (dp[pos][is-smaller][count-of-d]
already filled)

return dp[pos][is-smaller][count-of-d]

```
int ans = 0
```

```
int max-digit
```

```
if (is-smaller == true)
```

```
    max-digit = 9
```

```
else
```

```
    max-digit = num[pos]
```

```
for (i = 0; i ≤ max-digit; i++) {
```

```
    if (i == d)
```

```
        count-of-d++
```

```
    if (is-smaller == false &&
```

```
        i < num[pos])
```

```
        is-smaller = true
```

```
    ans += solve(pos+1, is-smaller,
```

```
        count-of-d)
```

```
dp[pos][is-smaller][count-of-d]
```

```
    = ans
```

```
return ans
```

main() {

num = b

ans = solve(0, 0, 0)

// solve for [0, b]

num = a-1

ans -= solve(0, 0, 0)

// solve for [0, a-1]

}

Q2 Investigation.

How many nos in range $[a, b]$ are there, whose sum of digits is divisible by K .

Eg - $[10, 28]$ $K = 3$

10 11 12 13 14 15 16 17 18 19

20 21 22 23 24 25 26 27 28

ans = 6

int dp[pos] [is smaller or bigger]
[sum-of-digits % K]

Main logic

```
for (i = 0; i ≤ max-digit; i++) {
```

```
    sum = (sum + i) % K
```

```
    if (is-smaller == false &&  
        i < num[pos])
```

```
        is-smaller = true
```

```
    ans += solve(pos+1, is-smaller,  
                sum)
```

}

Q3 Hard Investigation.

How many nos in range $[a, b]$ are there, whose sum of digits is divisible by K and the number itself is also div by K .

Eg - $[10, 28]$ $K = 2$

10 11 12 13 14 15 16 17 18 19

20 21 22 23 24 25 26 27 28

Ans = 5

int dp[pos][is-smaller]
[sum of dig % K] [num % K]

25

Main logic

for ($i=0$; $i \leq \text{max-digit}$; $i++$)

$\text{dig_sum} = (\text{dig_sum} + i) \% K$

$\text{new_num} = (10 * \text{num} + i) \% K$

if ($\text{is_smaller} == \text{false}$ &&
 $i < \text{num}[\text{pos}]$)

$\text{is_smaller} = \text{true}$

$\text{ans} += \text{solve}(\text{pos}+1, \text{is_smaller},$
 $\text{dig_sum}, \text{new_num})$

y

57321

$$573218 = 10 * 57321 + 8$$

$$635 \\ 6352 = 10 * 635 + 2$$

- new_num = $10 * \text{old_num} + \text{dig}$

TC

How many digits does n have

$$\Rightarrow \log_{10} n$$

$$\text{Iterations} \propto \log_{10} n$$

$$\text{Digit DP} \quad n \rightarrow \log_{10} n$$

{done}

