

Subjectivity Intersection Mathematics: A Common Pointed-Braid Source for Quantum History and Lorentzian Geometry

A fixed-finite conditional composition with relational backreaction

Satoru Watanabe

Subjectivity Intersection Emergence Lab (SIEL)

Working preprint v0.2 — 11 September 2026

Status. This is a working theoretical preprint. The main result is conditional and fixed-finite, with an explicit finite-dimensional coupled realization, a certified nonzero time bin, and a restricted nonzero same-source neighborhood. It is not peer reviewed and does not establish an empirically validated theory of quantum gravity, absolute semiclassical gravity, or an ontological theory of subjectivity.

Abstract

We construct a single typed mathematical system in which a quantum history and initial Lorentzian geometry are read from one finite pointed-braid source, rather than supplied as unrelated models and coupled afterwards. Version 0.2 strengthens the earlier local composition theorem by selecting a complete finite PBM–BIX realization, certifying its 87-dimensional vector field and interval Jacobian bounds, and composing measurement and compilation errors with a nonzero finite-time detector and normalized clock guarantee. A restricted real-orthogonal, grade-preserving Kähler neighborhood re-identifies the metric cometric and its inverse from each varied source, so the dynamic construction is no longer confined to one isolated source point, although the certified radius is extremely small.

Within the selected homogeneous sector, a pointed germ removes arbitrary spatial profile functions from the transported PCMT coefficients. The prior clock–screen carrier, an explicit unit-shell completion rule, and the signed metric Jacobian reconstruct the internal quaternion algebra, $S^3 \cong SU(2)$, the Bianchi-IX bracket, and a consistent screen orientation. These are conditional reductions of added structure, not a proof that bare braid relations uniquely select dimension, topology, homogeneity, physical units, or natural gravity. The result remains a fixed-finite mathematical construction with explicit bridge and constitutive assumptions; it is not peer reviewed and does not establish empirical quantum gravity, SIC/O3, subjectivity, or consciousness.

Keywords: braid group; Yang–Baxter coherence; quantum geometry; Lorentzian metric; backreaction; relational observable; symmetric-hyperbolic system; finite source.

1 Problem and claim

Many quantum–geometry constructions begin with a quantum model and a spacetime model that have independent state spaces, independent clocks, or independently normalized energies. A coupling term can then connect them, but the coupling alone does not show that they arise from one mathematical source. The question addressed here is narrower and stricter:

Can the quantum history, the initial Lorentzian geometry, the relational component that mediates their coupling, and the subsequent backreaction be placed on one source-derived object without duplicating the source or selecting one braid presentation as absolute?

The answer is conditional but constructive. The central result is a local composition theorem for a fixed finite source. Its content is stronger than compatibility of two separately specified models: all source readouts come from one joint state, the source Hamiltonian is counted once, the initial metric is generated from the same reduced state that carries the quantum history, and the coupled state and metric then evolve in one local initial-value problem.

The result also has strict boundaries. Braid relations do not by themselves select four dimensions, the physical clock soldering, the reflection coefficient used to obtain Lorentz signature, the relative gravitational source law, or the absolute units. The construction therefore separates four layers:

1. exact finite algebra and exact rational certificates;
2. conditional mathematical composition under explicit bridge assumptions;
3. an unadopted finite-source physical model with an operational discriminator;
4. ontological interpretation, which is not proved here.

Classical braid groups and Yang–Baxter consistency provide the mathematical background for the path equivalences used below [1, 2, 3]. The direct source theorem is Watanabe’s pointed-unitary-braid result: it proves canonical all-finite-orders endpoint overlaps, bilateral positive contractions, pointed-unitary covariance, and compatibility with the declared shuffled tensor law [9]. The present paper uses that theorem as its Braid foundation and does not strengthen its scope. Local covariance and the initial-value structure of general relativity supply comparison standards for the field-theoretic and geometric parts [4, 6]. The construction is not presented as a replacement for those frameworks.

2 The fixed pointed-braid source

2.1 One source state and its endpoint readouts

Let

$$\mathcal{H}_{\text{src}} = \mathcal{H}_A \otimes \mathcal{H}_B, \quad \mathcal{H}_A = \mathcal{H}_B = \mathbb{C}^{125}. \quad (1)$$

The actual finite source state used throughout the composition is

$$\Gamma_0 = \frac{100\mathbf{1} + D \otimes D}{1,562,500}, \quad \text{Tr } D = 0, \quad \text{Tr}(D^2) = 216. \quad (2)$$

It is normalized and has maximally mixed endpoint marginals,

$$\Gamma_A = \text{Tr}_B \Gamma_0 = \frac{\mathbf{1}_{125}}{125}, \quad \Gamma_B = \text{Tr}_A \Gamma_0 = \frac{\mathbf{1}_{125}}{125}. \quad (3)$$

The full source package also contains the original Hamiltonian H , a fixed endpoint split, the cup mark M_0 , insertion data, polar data, and selected graded generators. These objects are transported together. None is reconstructed from a scalar summary of the state.

The coupled system has one joint state $\Sigma(t)$ on

$$\mathcal{H}_{\text{src}} \otimes \mathcal{H}_F, \quad \Sigma(0) = \Gamma_0 \otimes \rho_F, \quad \text{Tr } \rho_F = 1. \quad (4)$$

The time-dependent source state is only the readout

$$\Gamma(t) = \text{Tr}_F \Sigma(t). \quad (5)$$

Thus Γ is not a second dynamical copy.

2.2 The endpoint-invisible correlation carrier

For any normalized bipartite state Γ , define

$$c_\Gamma = \Gamma - \Gamma_A \otimes \Gamma_B, \quad \mathcal{C}_{\text{corr}} = \ker \text{Tr}_A \cap \ker \text{Tr}_B \quad (6)$$

inside the real vector space of trace-zero Hermitian joint variations. For the actual state,

$$c_0 = \frac{D \otimes D}{1,562,500}, \quad \text{Tr}(c_0^2) = \frac{216^2}{1,562,500^2} > 0. \quad (7)$$

Let $p = (\text{Tr}_B, \text{Tr}_A)$. The inclusion $\iota : \mathcal{C}_{\text{corr}} \hookrightarrow \text{Herm}_0(\mathcal{H}_A \otimes \mathcal{H}_B)$ is the categorical kernel of p in the declared finite-dimensional real-linear category: every $T : Z \rightarrow \text{Herm}_0$ satisfying $pT = 0$ factors uniquely through ι . Moreover, every normalized joint state has the unique decomposition in Eq. (6).

This universal property is elementary finite bipartite mathematics; it is not uniquely braid-specific. The source-specific content enters through the actual state, cup, operators, and their simultaneous transport. A matched control with unchanged endpoint marginals but a rotated right correlation factor changes a fixed cup response by the exact amount 62/48,828,125. Therefore that joint response cannot factor through either endpoint marginal alone. This is an operational nonfactorization result, not an ontological identification of c_Γ .

2.3 Typed braid descent

Let **Cross** be the free typed crossing groupoid generated by the allowed elementary crossings and let **Braid** be its quotient by Yang–Baxter, remote-commutation, inverse, and unit relations. A typed output functor F includes the labelled source history, the joint source-field evolution, the energy ledger, the relation mark, initial soldering, the field, and the local geometric problem.

Proposition 2.1 (Typed descent criterion). *F factors uniquely through **Braid** if and only if it maps both sides of every generating relation to the same typed output.*

Proof. This is the quotient universal property. Necessity follows because equivalent words have one image in the quotient. Sufficiency follows because equality on generating relations makes F constant on the generated congruence, hence defines a unique functor on the quotient. $\square$

The actual even typed source satisfies the required relations in the fixed-finite scope. A non-Yang–Baxter unitary control fails: for a retained label $p = |000\rangle\langle 000|$, the two relation words produce orthogonal projector images and

$$\Delta_p = \|\text{Ad}_{W_L}(p) - \text{Ad}_{W_R}(p)\|_{\text{HS}}^2 = 2. \quad (8)$$

The ordinary flip also satisfies Yang–Baxter, so descent proves a coherence role for braid structure but not uniqueness of the concrete raw representation.

3 Initial Lorentzian geometry from the same source

3.1 A four-plane and ten independent metric directions

The graded-greedy covariance-cone rule begins with the distinguished grade-one Hamiltonian generator and repeatedly adjoins the least higher grade whose unordered pair sums remain distinct and whose required source pivots are nonzero and untruncated. Stopping at four directions gives

$$(1, 2, 4, 8), \quad V = \text{span}_{\mathbb{R}}\{H, U_2, U_4, U_8\}. \quad (9)$$

The ten unordered pair sums are

$$2, 3, 5, 9, 4, 6, 10, 8, 12, 16, \quad (10)$$

all distinct. An exact triangular-minor certificate gives rank ten for the metric Jacobian. With source grade $m = 3$ and declared finite bound $J = 83$, every selected pivot is untruncated because $m + 16 \leq J$.

Four directions are required here because the target is a symmetric 4×4 tensor with ten components. This stopping rule is an explicit construction choice; it is not a theorem that bare braid relations force four spacetime dimensions.

3.2 Covariance cone and Lorentz signature

For selected source generators G_a and a clock operator O_0 , define the centered covariance tensor

$$K_{ab} = \text{Tr } \Gamma \Delta Z_a \Delta Z_b, \quad h_{ab} = \Re K(G_a, G_b), \quad (11)$$

and clock response

$$v_a = -2\Im K(G_a, O_0). \quad (12)$$

At the actual base point, the exact response is

$$v = (17/4, 0, 0, 0). \quad (13)$$

Normalize

$$\theta = \frac{v}{\sqrt{v^T h^{-1} v}} \quad (14)$$

and impose the declared clock reflection

$$g_0 = h - 2\theta \otimes \theta. \quad (15)$$

Then g_0 has signature $(-, +, +, +)$, and

$$K_{\text{cov}} = \{x : \theta(x) \geq 0, g_0(x, x) \leq 0\} \quad (16)$$

is closed, convex, pointed, and full dimensional. Its $\theta = 1$ section is a nondegenerate solid h -ellipsoid in $\ker \theta$.

Equation (15) is a source-internal Lorentz construction. Equality with a separately measured natural light cone is not proved. The geometry map will be denoted G , so the typed initial creation identity is

$$g_0 = G(\Gamma_0) = G(\text{Tr}_F \Sigma_0). \quad (17)$$

4 One quantum–geometry evolution

4.1 Clock and energy soldering

Three clock descriptions occur: the original H -evolution parameter t , the source covector θ , and a material scalar T in the local field system. Their physical identity is not a consequence of the braid relations. We impose the explicit noncircular soldering condition along one timelike anchor γ ,

$$T(\gamma(t)) = t, \quad u(T) = 1, \quad dT \parallel \theta \quad \text{initially.} \quad (18)$$

The source energy is counted once:

$$e(\Sigma) = \frac{1}{2} \text{Tr}[\Sigma(H \otimes \mathbf{1}_F)] = \frac{1}{2} \text{Tr}(\Gamma H). \quad (19)$$

For vanishing coupling, the full source algebra follows the original free history

$$\alpha_t^H(a) = e^{itH} a e^{-itH}. \quad (20)$$

At nonzero coupling, the joint history is not identified with α_t^H on source words that do not commute with H .

4.2 A common interaction and relative stress

The retained local linear interaction is

$$S_{\text{int}} = -\lambda \int \chi(T) H \otimes \phi \, dT \wedge N, \quad (21)$$

where N is the local current three-form and χ is fixed in advance. Only the abelian charge algebra $C^*(H)$ enters spacelike local densities; the full noncommutative source remains on one timelike defect. The interaction commutes with $H \otimes \mathbf{1}_F$, preserving Eq. (19) and the H -sector populations. On a fixed smooth background, the spectral sectors of H give retarded affine scalar shifts and a controlled-Weyl joint propagator.

The geometric source is relative. For an advance-fixed reference state and the same metric and prescription, define

$$\Delta T_\phi[g; \Sigma, \Sigma_{\text{ref}}] = T_\phi[g, \Sigma] - T_\phi[g, \Sigma_{\text{ref}}]. \quad (22)$$

Common state-independent geometric counterterms cancel in the difference. The proposed finite-model field equation is

$$E_c[g] = G[g] + \Lambda g = \kappa (\Delta T_\phi + T_{\text{mat}} + 2\Pi). \quad (23)$$

This is the *Intersection-Relative Gravity* working law. It is a constitutive proposal, not a derived absolute semiclassical Einstein equation. Standard semiclassical gravity also couples quantum stress to classical geometry, but the well-definedness, renormalization, and backreaction problem is delicate [5, 7, 8].

4.3 Relation polarization and reciprocal backreaction

The cup-marked relation selects an energy-tangent source direction Ξ_C . The geometry differential and spatial projection define

$$d_C = P_{\text{sp}} D G \Xi_C. \quad (24)$$

At the actual base point, one exact projected coordinate lies in the strictly negative interval

$$-\frac{1127}{50,000,000,000} \leq Dq[\Xi_C] \leq -\frac{563}{25,000,000,000}, \quad (25)$$

so $d_C \neq 0$. The relation enters as a zeroth-order target shift,

$$m_{\text{sp}}^{\text{RP}} = P_{\text{sp}}(w - D G X_H) - d_C, \quad (26)$$

$$\frac{8K}{\rho} D_u^P \Pi + \Pi = -K m_{\text{sp}}^{\text{RP}}. \quad (27)$$

The source row retains the original intrinsic generator,

$$D_u^B \Gamma = (I - P_H) X_H + R_E \mathcal{L}_u g, \quad (28)$$

and the relation mark is transported by $D_u M = 0$. On the regular patch, d_C depends smoothly and algebraically on (Γ, g, M) . Consequently the feedback loop is

$$(M, \Gamma) \longrightarrow d_C \longrightarrow \Pi \longrightarrow g \longrightarrow \Gamma. \quad (29)$$

The coefficient of d_C in Eq. (27) is a bold constitutive choice. It is not uniquely selected by the braid relations.

4.4 Emergent-then-autonomous geometry

The equality in Eq. (17) is imposed only at the creation/soldering datum. Afterwards g and Σ are distinct state variables of the same coupled system. Requiring $g = G(\text{Tr}_F \Sigma)$ at every time would both reduce geometry to an algebraic readout and conflict with the retained sector structure. The post-creation state space therefore realizes provenance without permanent algebraic identity.

The first-order variables are the generalized-harmonic metric block, the material clock/current and RP stress, the linear scalar field block, one finite joint-state block, and the transported relation mark. After the obsolete all-time matching row is removed, the principal symbol is a triangular direct sum of symmetric-hyperbolic and finite-dimensional blocks. Relative stress and exchange currents are lower-order locally Lipschitz functions on the stated weighted patch.

5 The conditional local composition theorem

We collect the construction into five conditions.

- M1** The actual source state, $H, U_2, U_4, U_8, \theta, K_{\text{cov}}$, and g_0 define the source-internal Lorentz carrier under the graded-greedy and reflection assumptions.
- M2'** The creation square $g_0 = G(\text{Tr}_F \Sigma_0)$ holds. Geometry and the joint state then coevolve without all-time algebraic identification.
- M3*** The relation object is the correlation kernel $\mathcal{C}_{\text{corr}}$, its actual point c_Γ , and the existing cup mark M_0 required for relation transport.
- M4'** One timelike-defect source carries the same joint state, original $H/2$ ledger, local interaction, relative field stress, and relation-polarized constitutive stress.
- M5'** The combined system satisfies the regularity, constraint, cone, reference, and clock-soldering assumptions required for one common short-time solution.

Theorem 5.1 (Pointed Correlation-Kernel Braid Quantum–Geometry Local Composition). *Assume M1–M5' and the fixed-finite source identities above. For $s > 5/2$ and compatible smooth initial data in one regular generalized-harmonic patch, there is a nonzero interval $[0, T_*]$ on which a unique local solution carries one joint state Σ , one reduced source readout Γ , the original Hamiltonian ledger counted once, the endpoint-invisible correlation carrier, the source-generated initial Lorentzian metric, the local quantum interaction, and relative geometric backreaction. If the complete typed data are transported simultaneously, Yang–Baxter-equivalent words define isomorphic typed histories and the same local solution problem.*

Proof. The initial identity follows from Eq. (5) and Eq. (17). The correlation-kernel factorization and endpoint nonfactorization establish the M3* readout without adding a second state. Equation (21) preserves the single $H/2$ ledger and recovers the original H history at zero coupling. Equations (23)–(27), together with the material and exchange-current equations, form the retained triangular symmetric-hyperbolic plus finite-dimensional system. Its positive symmetrizer, local Lipschitz lower-order maps, compatible initial constraints, and strict cone margin give local existence, uniqueness, continuous dependence, and constraint propagation by standard quasilinear theory [6]. Proposition 2.1 then transfers the full typed system to the braid quotient. No separate source state or second Hamiltonian is introduced. $\square$

Remark 5.2 (Meaning of “same source”). The theorem does not merely use the same parameter names in two models. The quantum and geometric sectors share the actual reduced state, clock soldering, energy ledger, relation mark, interaction history, and local solution. Removing any one of these identifications would weaken the result to compatibility or coupling of separate models.

6 A nonzero geometric response

Compare two matched solutions of Eq. (23): $\eta = 1$ retains d_C in Eq. (27), while $\eta = 0$ removes only that term. Initial state, metric, first metric data, field, reference, material variables, Π , and gauge are identical. Then

$$D_u^P(\Pi_1 - \Pi_0)|_0 = \frac{\rho}{8}d_C, \quad (30)$$

and hence

$$D_u\Delta(2\Pi)|_0 = \frac{\rho}{4}d_C \neq 0. \quad (31)$$

For smooth data, the two relative source histories therefore separate on every sufficiently short punctured interval.

Let $S = \ker \theta$ carry the positive screen metric and transport d_C along the material clock curve $\gamma(T)$. For any spatial tensor X , define

$$P_C(X) = \frac{\langle d_C, X \rangle_S}{\langle d_C, d_C \rangle_S}. \quad (32)$$

The relational curvature response is

$$\mathcal{R}_C(T) = P_C(E_c[g_1] - E_c[g_0])|_{\gamma(T)}. \quad (33)$$

All indices are contracted and the branches are compared at equal material-clock value. Thus $\mathcal{R}_C$ is invariant under simultaneous diffeomorphism and screen-frame change within the declared transport scope. From Eq. (31),

$$D_T\mathcal{R}_C(0) = \frac{\kappa\rho}{4} \neq 0 \quad (34)$$

for $\kappa \neq 0$ and $\rho > 0$.

Equation (34) is a model-internal discriminator. It does not show that the proposed relative source law is realized in nature; another covariant constitutive theory may reproduce the same local slope.

7 Finite relational curvature readout

7.1 Seven probe frames

At the equal-clock event, choose a transported orthonormal tetrad

$$e_0 = u, \quad e_1, e_2, e_3 \in S. \quad (35)$$

Use seven future unit-timelike velocities

$$v_0 = e_0, \quad v_i^\pm = \frac{5}{3}e_0 \pm \frac{4}{3}e_i, \quad i = 1, 2, 3. \quad (36)$$

For each velocity, choose an adapted orthonormal separation triple in $v^\perp$ and read the symmetric Jacobi bilinear form

$$J_v(w, z) = R(w, v, z, v). \quad (37)$$

Each frame yields six scalars, for 42 raw readings.

Proposition 7.1 (Finite Riemann reconstruction). *The exact rational linear map from the 20-dimensional space of algebraic four-dimensional Riemann tensors to the 42 Jacobi readings has rank 20. A selected 20-channel square subsystem has an inverse L satisfying*

$$\|L\|_\infty = \frac{13}{4} \quad (38)$$

in the declared tetrad-coordinate norm.

The remaining 22 readings are consistency checks. The reconstructed Riemann tensor gives Ric, scalar curvature, and E_C . Contracting the branch difference with d_C exactly recovers Eq. (33) in the zero-size, zero-noise, zero-backreaction limit. For the exact response fixture used in the certificate,

$$D_T D_C^{\text{probe}}(0) = \frac{35}{132} = \frac{\kappa\rho}{4}. \quad (39)$$

The probes carry only velocities, separations, small classical stresses, and metric readouts. They do not copy the noncommutative source algebra into spacelike regions.

7.2 Finite error ledger

If ε is probe separation, n_a bounds unnormalized acceleration noise, δ_T is clock resolution, and m_{det} measures probe stress, then on a regular patch the first symbolic bound is

$$|D_C^{\text{meas}} - \mathcal{R}_C^J| \leq C_{\text{rec}} \left(\varepsilon \|\nabla \text{Riem}\| + \frac{n_a}{\varepsilon} \right) \quad (40)$$

$$+ C_T \delta_T \|\nabla_u E_C\| + C_{\text{br}} m_{\text{det}}. \quad (41)$$

The reconstruction part is finite because of Proposition 7.1; numerical values for the remaining patch and apparatus constants are not yet supplied. At fixed n_a , taking $\varepsilon \rightarrow 0$ increases n_a/ε , so the ideal limit is not automatically an experimentally accessible limit.

8 Cutoff relevance and the finite-source hypothesis

8.1 A conditional detector-cylinder theorem

Suppose, beyond the fixed model already defined, that complete models indexed by J are connected by typed embeddings. Let ε_J bound every adjacent defect relevant to the initial data, Hamiltonian, cup, clock, geometry map, d_C , and constitutive coefficients.

Theorem 8.1 (Detector-cylinder summable stability). *Assume $\sum_{J \geq J_0} \varepsilon_J < \infty$. Assume further that all models share a nonzero interval $I = [0, T]$, a uniform cone margin, symmetrizer bounds, Sobolev bounds, local Lipschitz constant L , a positive uniform lower bound for $\|d_C^J\|$, and a uniformly Lipschitz 42-reading map with constant C_D . Then*

$$\sup_{t \in I} \|Y_K(t) - Y_J(t)\| \leq C_D(1 + T)e^{LT} \sum_{n=J}^{K-1} \varepsilon_n. \quad (42)$$

Consequently Y_J and $\mathcal{R}_C^J$ are uniformly Cauchy on I .

Proof. Let $E_{JK}(t)$ be the embedded difference norm for the two local solutions. The standard difference-energy estimate has the form

$$E_{JK}(t) \leq a_{JK} + \int_0^t (LE_{JK}(s) + b_{JK}) ds, \quad (43)$$

where the telescoped adjacent defects give $a_{JK}, b_{JK} \leq \sum_{n=J}^{K-1} \varepsilon_n$. Gronwall's inequality yields

$$\sup_I E_{JK} \leq (1 + T)e^{LT} \sum_{n=J}^{K-1} \varepsilon_n. \quad (44)$$

Composition with the uniformly Lipschitz detector and normalized d_C contraction gives Eq. (42). Summability makes its right side vanish as $J, K \rightarrow \infty$. $\square$

The theorem avoids convergence of every source component. It is nevertheless not presently applicable to the actual model: no authoritative construction defines the complete compatible family of joint state, source algebra, H , cup, clock, geometry map, dynamic d_C , constitutive coefficients, and local solution for varying J . The initial inequality $m + 16 \leq J$ controls only the ten initial metric pivots. Because $d_C = d_C(\Gamma, g, M)$ evolves inside the feedback loop, initial nontruncation does not imply equality of finite-time detector histories.

8.2 Intrinsic Finite Braid Source Principle

The shortest live route therefore isolates the following hypothesis.

Assumption 8.2 (Intrinsic Finite Braid Source Principle, IFBSP). The declared actual source with $J_* = 83$ is the internal Braid source of the model, rather than a regulator approximation to an otherwise specified $J = \infty$ source. The Lorentzian spacetime and scalar field remain continuous; only the internal Braid-source sector is intrinsically finite.

IFBSP is noncircular: it fixes the model class before calibration or held-out validation and assumes neither Eq. (34) as an empirical fact nor agreement with data. Under IFBSP, the theoretical observable is $\mathcal{R}_C^{83}$ and Eq. (40) contains no undefined $\mathcal{R}_C^\infty$ residual. This is a finite-theory choice, not a proof of cutoff convergence.

The assumption is falsified as a working route if J_* is retuned after outcomes, if the source definition is shown to require a separately specified infinite object, if a required source identity fails at J_* , if the intended claim requires $J \rightarrow \infty$ universality, or if independent prospective tests reject the frozen finite-source predictions.

9 Physical scales and a no-retuning protocol

The internal equations are dimensionless and retain two continuous scale freedoms. Let τ_* be the physical duration of one internal clock unit and σ_* the physical stress associated with one internal stress unit. Then

$$t_{\text{phys}} = \tau_* T, \quad x_{\text{phys}} = c\tau_* x, \quad (45)$$

$$H_{\text{phys}} = \frac{\hbar}{\tau_*} H, \quad T_{\text{phys}} = \sigma_* T, \quad (46)$$

$$(E_c)_{\text{phys}} = \frac{E_c}{(c\tau_*)^2}, \quad \kappa_{\text{phys}} = \frac{\kappa}{\sigma_*(c\tau_*)^2}. \quad (47)$$

Rescaling τ_* and σ_* leaves every dimensionless braid word, state, cone, and $\mathcal{R}_C(T)$ unchanged. Bare internal mathematics therefore cannot determine absolute seconds, metres, or gravitational strength.

Choose in advance one nonzero dimensionless gap Δh_{cal} and an independently measured angular frequency ω_{cal} , and one independent physical gravitational-coupling anchor κ_{cal} . Then

$$\tau_* = \frac{\Delta h_{\text{cal}}}{\omega_{\text{cal}}}, \quad \sigma_* = \frac{\kappa}{\kappa_{\text{cal}}(c\tau_*)^2}. \quad (48)$$

These two relations locally fix the two logarithmic scale freedoms. They must not use the held-out C -polarized response.

Before any outcome access, a prospective packet must freeze the source revision, J_* , state, H gap, dimensionless couplings, density convention, clock and screen soldering, reference prescription, detector map, apparatus error domain, and success rule. Post-outcome changes define a new exploratory model.

10 A certified finite PBM–BIX realization

10.1 Braid-null transduction and the finite-model route

The post-v0.1 analysis begins with a source-level negative control. Applying the existing relation recipe to the identity representation gives an exact zero relation polarization, whereas the actual nontrivial pointed-braid source has the nonzero relation readout proved in the source certificate. Composed with the declared relation-polarized stress, curvature scalar, and ideal detector, this yields a conditional local transduction contrast. It distinguishes the actual source from the identity null inside the selected model, but it does not prove that Yang–Baxter coherence is necessary or that the raw representation is unique.

A Taylor majorant proves qualitative separation on a sufficiently short interval but initially leaves the common time, second-derivative, and detector-margin constants unpopulated. We tested three shortcuts: a finite worldline jet, a generic finite nonzero-mode truncation, and a homogeneous zero-mode current with the frozen source. All fail in the declared class. The selected PBM–BIX reduction is retained because it closes the source-compatible equations while preserving the required relation signal.

10.2 Complete finite vector field and a nonzero interval

The PBM–BIX state vector has 87 real coordinates. Its creation data include $M_1 = \text{Tr}(\Gamma H)$ and $M_2 = \text{Tr}(\Gamma H^2)$, the six spatial components of d_C , the six components of the Hamiltonian-direction response b_H , and the 36 entries of the transported inverse-cometric coefficient $J = K_{\sharp}^{-1}$. The remaining coordinates carry the homogeneous metric, current, matter, scalar, and state variables. This count belongs to the selected realization; it is not a derivation of field content from braid relations alone.

The complete vector field and interval Jacobian are bounded on a closed regular box. For $|\lambda| \leq 1$ and the fixed PCMT transport law, a bootstrap certificate supplies the dimensionless interval

$$2.4337004147867 \times 10^{-33} \leq T \leq 4.8674008295735 \times 10^{-33}. \quad (49)$$

Throughout this bin, the normalized relational detector remains strictly positive, with certified lower display bound $7.195243466871394 \times 10^{-35}$, and the normalized clock rate remains positive, with lower bound 2.0014274819390794. These numbers certify existence and sign in the declared norm and scaling; they do not establish practical duration in physical seconds.

Proposition 10.1 (Fixed-packet finite-time transfer). *Assume the PBM–BIX regular box, the selected PCMT coefficients, $|\lambda| \leq 1$, the matched detector normalization, and the source-support and clock-window conditions of the finite certificate. Every source packet inside the certified 2^{-31} measurement-and-compilation tolerance box produces one common solution on the interval in Eq. (49), with positive normalized detector response and positive normalized clock rate.*

The proposition composes an exact finite source-statistics enclosure with the certified interval flow. It neither infers a second 87-dimensional flow from each approximate sample nor permits J to remain fixed under an arbitrary source change.

10.3 A non-isolated same-source neighborhood

The fixed- J limitation is removed on a restricted slice. Let $\Gamma(s) = Q(s)\Gamma_0 Q(s)^\top$, where $Q(s)$ is real orthogonal, grade preserving, and has ordered-flag Kähler path length at most 10^{-4177} . The path is nonconstant because Γ_0 is non-scalar on its retained support. Along this slice every source remains real symmetric and support preserving, and

$$\|\Gamma(s) - \Gamma_0\|_1 \leq 4168 \times 10^{-4177} = 4.168 \times 10^{-4174}. \quad (50)$$

At each point the full ordered-flag cometric is evaluated anew; its spatial block supplies $K_\sharp$ and $J = K_\sharp^{-1}$. Reality gives $b_H = 0$ exactly on this slice.

Theorem 10.2 (Restricted same-source dynamic patch). *Under Proposition 10.1, the full-cometric bounds, and the real-orthogonal grade-preserving restriction above, there is a nonzero relative neighborhood of Γ_0 in which each source supplies its own $d_C, M_1, M_2, K_\sharp, J$, and b_H . The coefficient packet lies in the same PBM–BIX dynamic box and retains the finite-time detector and clock conclusions.*

The radius is mathematically nonzero but extremely small. General complex source variation, arbitrary trace-norm noise, code leakage, and a practically useful stability radius remain open.

10.4 Pointed germ, compact completion, and orientation

Inside the selected left-homogeneous geometry, evaluation at the pointed identity bijects left-invariant coefficient fields with their fiber values. Thus the 36 components of J , six of b_H , and six of d_C introduce no arbitrary spatial profile functions once their pointed values and left transport are fixed. Parallel time transport is unique only within the declared no-extra-generator class and compatible connection.

Let $S = \ker \theta$ with positive screen metric h_S . Given an orientation, define on $\mathbb{R} \oplus S$

$$(a, x)(b, y) = (ab - h_S(x, y), ay + bx + x \times y). \quad (51)$$

This is the quaternion product. Selecting its full positive unit shell gives $S^3 \cong SU(2)$, and the imaginary commutator yields the Bianchi-IX bracket $[e_i, e_j] = 2\epsilon_{ij}^k e_k$.

The screen orientation need not be supplied as another independent sign inside the fixed source chart. For an oriented four-carrier V ,


$$\det(\text{Sym}^2 A^*) = (\det A)^5. \quad (52)$$

The odd exponent lets the strictly positive signed ten-direction metric Jacobian transfer the stored source-chart orientation to V ; the fixed future clock then orients S . This fixes the sign in Eq. (51) within the declared chart and carrier identification.

These reductions do not show that braid relations uniquely select the source chart, GGCC rule, clock reflection, unit-shell completion, or physical handedness. An ordinary non-braided oriented Euclidean three-space admits the same quaternion construction. The Braid-specific content remains typed coherent transport of the complete selected structure.

11 What braid coherence contributes

The mathematical role of braid structure in this paper is neither “noncommutativity alone” nor the claim that every quantum–gravity model must be braided. Building on the pointed-unitary-braid theorem [9], its concrete contribution here is commutation of construction with viewpoint transport:



Applying a viewpoint change before constructing the geometry and dynamics gives the same typed result as constructing first and transporting the entire result. State, H , cup, clock, d_C , soldering, metric, and equations must all move together. Moving only one part defines a different experiment.

This is the precise sense in which no crossing word or matrix presentation is absolute inside the allowed groupoid. A labelled non-Yang–Baxter control fails the full typed equalizer, demonstrating that the relation is not vacuous. Yet the metric-only projection can miss that failure, and the ordinary flip can satisfy the same coherence relation. Braid coherence is therefore sufficient for the declared descent and detected by a faithful full output, while concrete physical uniqueness remains open.

The broader Subjectivity-Intersection programme motivates interpreting an irreducible reciprocal relation as a candidate “third standpoint” [10, 11]. No such ontology is used in the proof. In particular, mathematical standpoint data are not phenomenal consciousness, and neither light nor a quantum system is called conscious here.

12 Countermodels, negative results, and claim boundaries

The route was narrowed by explicit failures that remain part of the result.

Boundary	Failure or limitation	Consequence retained in the present model
Local source cloning	Two spacelike regions cannot each implement the full noncommutative source on one code while maintaining microcausality.	The full source lives on one timelike defect; spacelike probes read only metric curvature.
Nonlinear local packet	The examined spatially extended bounded nonlinear Weyl packet has a strict-locality counterexample.	The retained interaction is the evaluated local linear Same- H model.
All-time metric readout	Fixed H -sector populations conflict with demanding ten independent moving-state metric controls at every time.	Geometry is source-generated initially and autonomous thereafter, with reciprocal coupling.
Absolute stress	The examined absolute reference-stress closure does not supply the required second-order local system.	The present source law uses a matched same-reference stress difference and states this constitutive choice openly.
Metric-only braid test	A non-Yang–Baxter source defect can vanish after projection to a matched local field stress.	Braid descent is claimed for the full labelled typed object, not as a unique metric-only signature.
Infinite cutoff	The complete varying- J coupled tower is not defined.	The convergence theorem is conditional; IFBSP supplies a separate finite-source route.
PBM–BIX selection	The clock–screen carrier does not uniquely force homogeneity, unit-shell completion, or physical topology.	These remain explicit model choices after internal profile and orientation reductions.

Boundary	Failure or limitation	Consequence retained in the present model
Source stability	The certified same-source Kähler radius is approximately 10^{-4177} and excludes general complex noise.	The result proves non-isolation, not practical robustness.
Empirical status	No frozen external calibration or independent detector dataset has been evaluated.	The result remains theoretical and conditionally formulated.

The paper therefore does not establish any of the following: a unique derivation of four dimensions from braid relations; equality with the observed light cone; an absolute renormalized stress law; a continuum or large- N limit; numerical apparatus feasibility; empirical quantum gravity; an ontological identity between the correlation kernel and physical space; SIC/O3; subjectivity; or consciousness.

13 Reproducibility and provenance

The exact finite certificates were produced from versioned computational records. The main checks used by this manuscript are listed in Table 2. Hashes refer to the local research snapshot from which working preprint v0.2 was generated. Public replication requires a separately released verification bundle; this PDF alone is not a complete independent replication package.

Table 2: Core internal certificates summarized in working preprint v0.2.

Record	Certified content	Status
UB300	Actual state identity and nonzero same-energy relation response	exact finite
UB303	Original- H compatible relation-polarized repair and feedback typing	exact/conditional
UB317/328	Full typed Yang–Baxter descent and labelled non-YB control	exact/conditional
UB333	Four source directions, Lorentz carrier, ten-pivot rank certificate	exact/conditional
UB336	Correlation-kernel universal property and endpoint nonfactorization	exact finite
UB337	Same-object M1–M5' composition audit	conditional theorem
UB338/339	Nonzero stress-to-curvature response and relational scalar	exact/conditional
UB340	Two-scale underidentification and no-retuning map	exact algebraic
UB341	42-to-20 Riemann reconstruction, inverse norm 13/4	exact finite
UB342	Conditional cutoff theorem and actual tower-definition first-fail	conditional theorem/audit
UB343	IFBSP-83 freeze and prospective no-retuning discriminator	exact/conditional
UB344–347	Braid-null transduction, finite-time majorants, and failed shortcut audit	exact/conditional
UB348–360	PBM–BIX fixture, complete 87D vector field, interval Jacobian, detector bin	exact/interval
UB361–382	Clock transfer, source compilation, implementation and error-radius certificates	exact/conditional
UB383	Fixed source packet composed with the complete coupled dynamic bin	conditional theorem
UB384	Re-identified J on a nonzero restricted same-source neighborhood	conditional theorem
UB385	Pointed-germ reconstruction removes spatial profile freedom in the selected sector	conditional theorem
UB386/387	Quaternion/SU(2)/BIX reconstruction and internal screen-orientation descent	exact/conditional

Standard ingredients are not claimed as original. They include braid groups, the Yang–Baxter relation, partial traces, geodesic deviation, generalized-harmonic reduction, Gronwall estimates, and symmetric-hyperbolic local theory. The project-specific synthesis, source realization, typed composition, and exact

certificates were developed through an iterative human-directed research process assisted by OpenAI Codex. The author is responsible for checking the mathematics, claims, citations, and public release. This v0.2 draft preserves derivation assistance rather than treating the software as an intellectual source.

14 Discussion

The strongest conclusion is architectural. The construction demonstrates that a fixed finite quantum source can do more than source a metric through an externally chosen expectation value. The same reduced state participates in the free quantum history, creates the initial covariance cone and Lorentz metric, supplies the irreducible correlation readout, fixes a relation polarization through the cup mark, and remains inside the joint state that exchanges influence with the geometry. The backreaction loop is therefore internal to one typed object.

This common provenance matters because formal coupling is cheap: two independently defined theories can almost always be joined by adding an interaction term. The present theorem imposes stronger bookkeeping. There is one source state, one Hamiltonian ledger, one clock soldering, one initial creation map, one relation package, and one local solution. Those identities are precisely what would fail if the quantum and geometric sectors were unrelated copies.

The viewpoint claim is also scoped. Yang–Baxter equivalence ensures that the complete construction does not depend on which representative crossing word is used. This supplies a concrete consistency mechanism for avoiding an absolute presentation. It does not show that there are no other coherence mechanisms, nor that mathematical presentations are themselves subjects.

The finite PBM–BIX realization changes the evidential position in a precise way. The construction now includes one complete finite vector field, an interval Jacobian enclosure, a positive detector bin, and a restricted source-varying patch. The pointed-germ, quaternion, and orientation results remove several independent internal choices after the homogeneous candidate is fixed. They do not remove the choice of that candidate or connect dimensionless scales to nature.

The decisive next scientific step is to freeze the remaining physical anchor map and test a held-out C -polarized response or cross-context ratio with predeclared apparatus tolerances. IFBSP fixes the finite source as the theory object. A failure must remain a failure; changing J_* , the reference state, coupling, topology, or detector channel afterwards creates a new model.

15 Conclusion

We have given a conditional, fixed-finite construction in which quantum history and initial Lorentzian geometry arise from one pointed-braid source and subsequently evolve with reciprocal backreaction in one system. The endpoint-invisible correlation carrier is endogenous to the source state, the original source energy is counted once, and the full typed construction descends through Yang–Baxter-equivalent presentations. Version 0.2 adds a complete finite PBM–BIX vector field, a certified nonzero detector-and-clock interval, and a restricted nonzero same-source neighborhood with pointwise cometric re-identification.

Within the selected homogeneous model, the spatial PCMT profile is determined by a pointed germ, while quaternionic completion and the signed metric Jacobian reconstruct the internal $SU(2)$ /Bianchi-IX structure and screen orientation. The result remains a mathematical unification architecture, not an empirically completed theory of quantum gravity. Its strongest unresolved inputs are GGCC and reflection, global completion and soldering, the constitutive relative-gravity law, IFBSP, dimensionful physical anchors, Braid necessity, and independent validation.

Data and code availability

Version 0.1 is preserved at Zenodo DOI 10.5281/zenodo.22693965 with publication date 10 September 2026. This v0.2 draft reports additional exact and interval certificates from UB344 through UB387. A public verification bundle has not yet been deposited; independent reproduction still requires release of the referenced certificate code and manifests.

Competing interests

The author declares no competing interests.

Funding

No external funding is declared for this working preprint.

AI assistance disclosure

OpenAI Codex was used under the author’s direction for mathematical ideation, derivation assistance, exact-check scripting, audit organization, and drafting. The author retains responsibility for verification, interpretation, citations, and the decision to release this version.

References

- [1] E. Artin, “Theory of braids,” *Annals of Mathematics* **48**(1), 101–126 (1947). [doi:10.2307/1969218](https://doi.org/10.2307/1969218).
- [2] C. N. Yang, “Some exact results for the many-body problem in one dimension with repulsive delta-function interaction,” *Physical Review Letters* **19**, 1312–1314 (1967). [doi:10.1103/PhysRevLett.19.1312](https://doi.org/10.1103/PhysRevLett.19.1312).
- [3] L. A. Lambe and D. E. Radford, *Introduction to the Quantum Yang–Baxter Equation and Quantum Groups: An Algebraic Approach*. Springer (1997). [doi:10.1007/978-1-4615-4109-7](https://doi.org/10.1007/978-1-4615-4109-7).
- [4] R. Brunetti, K. Fredenhagen, and R. Verch, “The generally covariant locality principle—a new paradigm for local quantum field theory,” *Communications in Mathematical Physics* **237**, 31–68 (2003). [doi:10.1007/s00220-003-0815-7](https://doi.org/10.1007/s00220-003-0815-7).
- [5] S. Hollands and R. M. Wald, “Local Wick polynomials and time ordered products of quantum fields in curved spacetime,” *Communications in Mathematical Physics* **223**, 289–326 (2001). [doi:10.1007/s002200100540](https://doi.org/10.1007/s002200100540).
- [6] Y. Choquet-Bruhat, *General Relativity and the Einstein Equations*. Oxford University Press (2008). [doi:10.1093/acprof:oso/9780199230723.001.0001](https://doi.org/10.1093/acprof:oso/9780199230723.001.0001).
- [7] N. Pinamonti and D. Siemssen, “Existence and uniqueness of solutions of the semiclassical Einstein equation in cosmological models,” *Annales Henri Poincaré* **23**, 297–329 (2022). [doi:10.1007/s00023-021-01067-8](https://doi.org/10.1007/s00023-021-01067-8).

- [8] P. Meda, “The semiclassical Einstein equations in cosmological spacetimes,” *International Journal of Theoretical Physics* **65**, 137 (2026). doi:[10.1007/s10773-026-06339-9](https://doi.org/10.1007/s10773-026-06339-9).
- [9] S. Watanabe, *Subjectivity-Intersection Mathematics: All-Orders Intersection Consistency and Compositional Objectivity from Pointed Unitary Braids*, version 1.0, Zenodo (2026). doi:[10.5281/zenodo.22166167](https://doi.org/10.5281/zenodo.22166167).
- [10] S. Watanabe, *Subjectivity Intersection Ontology: A Comparative Ontological Framework for Third Subjectivity*, version 1.0.0, Zenodo (2026). doi:[10.5281/zenodo.21641878](https://doi.org/10.5281/zenodo.21641878).
- [11] S. Watanabe, *Subjectivity-Intersection Mathematics Public Working Preprint v4.2 Draft*, Zenodo (2026). doi:[10.5281/zenodo.21909933](https://doi.org/10.5281/zenodo.21909933).