

a
best sample per chain (solid)
t = 0.1 mean (translucent)
random
Sequence recovery (%)
74.9
60.4
69.4
55.1
56.2
51.8
52.2
47.0
53.0
35.8
PACo-Aptamer
RhoDesign
AlignIF
NA-MPNN
gRNAde
b
PACo-Aptamer recovery (%)
RhoDesign recovery (%)
PACo-Aptamer higher
equal recovery
c
PACo-Aptamer recovery (%)
AlignIF recovery (%)
PACo-Aptamer higher
equal recovery
d
Cumulative share of chains (%)
RhoDesign
gRNAde
NA-MPNN
AlignIF
Recovery advantage of PACo-Aptamer
over each baseline (%)
e
Recovery, t = 0.1 mean (%)
Mean log probability of sampled designs
f
ρ > 0
ρ < 0
Chains (%)
Within-chain Spearman ρ
g
+1.69
+1.46
Recovery − pool mean (%)
Designs kept by log-probability ranking (%)
h
PACo-Aptamer
RhoDesign
AlignIF
NA-MPNN
gRNAde
method mean
Recovery, best sample (%)
≤40 nt
40–80 nt
>80 nt
Native chain length
i
Mango-III
8HB1
1MMS
Purine
Theophylline
ZTP
dG
Xanthine
3GCA
Recovery advantage over RhoDesign,
t = 0.1 mean (%)