

a
EGARPS+ native probability
ProteinX ipTM
RhoFold mean pLDDT
0.71 0.71 0.73 0.46 0.17 0.29 0.37 0.58 0.83
0.33 0.25 0.73 0.46 0.54 0.36 0.63 0.83 0.44
0.38 0.50 0.80 0.43 0.62 0.53 0.30 0.17 0.61
Thrombin 7+ / 3−
VEGF 8+ / 3−
PDGF-BB 5+ / 3−
IL-17A 7+ / 4−
Angiopoietin-2 8+ / 3−
SARS-CoV-2 N 12+ / 3−
FGF2 10+ / 3−
swine H3 HA 4+ / 3−
IFN-γ 6+ / 3−
AUROC (binder vs negative)
0.9
0.7
0.5
0.3
0.1
b
EGARPS+ native probability
ProteinX ipTM
RhoFold mean pLDDT
ρ
Thrombin (n = 6) +0.31
VEGF (n = 8) +0.14
PDGF-BB (n = 5) +0.10
IL-17A (n = 6) +0.03
Angiopoietin-2 (n = 3) −0.50
SARS-CoV-2 N (n = 6) +0.09
FGF2 (n = 10) +0.47
swine H3 HA (n = 4) −0.80
IFN-γ (n = 6) +0.71
Thrombin (n = 6) +0.31
VEGF (n = 8) +0.24
PDGF-BB (n = 5) −0.41
IL-17A (n = 6) +0.26
Angiopoietin-2 (n = 3) +1.00
SARS-CoV-2 N (n = 6) −0.43
FGF2 (n = 10) +0.43
swine H3 HA (n = 4) +0.40
IFN-γ (n = 6) +0.66
Thrombin (n = 6) −0.20
VEGF (n = 8) +0.17
PDGF-BB (n = 5) +0.56
IL-17A (n = 6) −0.03
Angiopoietin-2 (n = 3) +0.50
SARS-CoV-2 N (n = 6) +0.09
FGF2 (n = 10) −0.18
swine H3 HA (n = 4) −0.80
IFN-γ (n = 6) +0.60
−1 −0.5 0 0.5 1
Spearman ρ (oracle score vs −log10 Kd)
nominal pass (ρ ≥ 0.6)
c
Native-pose rank
15/21 12/21 17/21 8/21 15/21 18/21 8/21 17/21 16/21 8/20 20/21
TBA HD22 Toggle-25t M08s-1 T1.1 DTRI-177 11F7t ARC1172 BT-100 A58-T10 FY-apt
7/21 14/21 9/21 6/21 14/21 18/21 13/21 14/20
A9g Kato16 p50-apt C13.18 TfR1-apt PfLDH-1 C8N5t pL1
d
EGARPS+ native probability
thrombin series ρ = +0.49 (n = 6)
pegaptanib
Measured Kd (nM)
e
RhoFold 9 11
ProteinX 11 8
EGARPS 11 9
Decision-layer overreach events
ranking objective
screening evaluation
f
Cliff's δ, coordination benefit
permutation null
(10^4 sign flips)
P = 0.06
observed
δ = +0.14
GO gate δ ≥ 0.3
δ = +0.14
P = 2.6e−3
δ = +0.05
P = 0.21
δ = -0.01
P = 0.73
Apparent effect
Budget-matched
Fixed resampling