1

QUESTION: 12.13.6.9

ROLL NO:EE22BTECH11027 NAME: KATARI SIRI VARSHINI

12.13.6.9.An experiment succeeds twice as often as it fails. Find the probability that in the next six trials, there will be at least 4 successes.

Solution: Let p be the probability for the experiment to succeed and q for the failure.

Here, it is given that probability of success is twice that of the failure, so

$$p = 2q$$

$$q = \frac{1}{3}$$

$$p = \frac{2}{3}$$
(1)

Now, let's consider a single trial as a bernuolli random variable $X_i = 1$ represents success and $X_i = 0$ represents failure.

$$X_i$$
 1 success 0 failure

Therefore we have,

$$P(X_i = 1) = p = \frac{2}{3} \tag{2}$$

$$P(X_i = 0) = q = \frac{1}{3} \tag{3}$$

Since we have n=6 trials, the random variable X representing the number of successes in 6 trials follows a binomial distribution. The cumulative distribution function (CDF) of X is given by

$$F_X(k) = P_X(X \le k) = \sum_{k=0}^{n} {}^{n}C_k q^{n-k} p^k$$
 (4)

Here,

$$n = 6, p = \frac{2}{3}, q = \frac{1}{3}, k = 0, 1, 2...6$$

We need to find the probability for the experiment to succeed to at least 4 times i.e. $P_X(X \ge 4)$. Using equation 4 we get,

$$P_X(X \ge 4) = 1 - P_X(X \le 3)$$

$$= 1 - F_X(3)$$

$$= 1 - \frac{233}{36} \ge 0.680$$
(5)

Therefore the probability that in the next six trials, there will be at least 4 successes is 0.680.