

a
12-Axis Non-CNS Clinical Trial Census
Phase II
Phase III
Phase IV
Solid Oncology
Metabolic Endocrine
Hematologic Malignancy
Pulmonary IPF ARDS
Cardiovascular HF Athero
Autoimmune IBD Rheum
Renal CKD AKI
Dermatologic Immune
Musculoskeletal Bone
Ophthalmic Retinal
Hepatic MASH Fibrosis
Infectious Sepsis
0
5000
10000
15000
20000
Unique Non-CNS Trials (n = 41,396)
b
Multi-Agent Stop Deconvolution
Efficacy / Futility
305 (4.6%)
Safety / Toxicity
466 (7.1%)
Unreported / Other
3,118 (47.3%)
Admin / Enrollment
2,706 (41.0%)
0
10
20
30
40
50
60
Share of Stopped Trials (%, n = 6,595)
c
Modality x Phase Completion Matrix
Small Molecule
96.4%
(n=7,454)
97.3%
(n=4,403)
99.2%
(n=2,737)
Antibody / Biologic
95.3%
(n=4,994)
96.7%
(n=2,953)
98.4%
(n=1,096)
Cell / Oligo / Other
96.7%
(n=184)
95.1%
(n=102)
100.0%
(n=44)
Phase II
Phase III
Phase IV
Scientific Completion (%)
95
90
85
80
75
d
OpenTargets Target Coverage by Axis
Target-Resolved Trials (%)
100
90
80
70
60
50
40
30
20
Metabolic
Autoimmune
Solid
Pulmonary
Hepatic MASH
Infectious
Overall 72.0% (29,791)
200
400
600
800
Unique Human Target Genes (n = 1,155 total)