



저작자표시-비영리-변경금지 2.0 대한민국

이용자는 아래의 조건을 따르는 경우에 한하여 자유롭게

- 이 저작물을 복제, 배포, 전송, 전시, 공연 및 방송할 수 있습니다.

다음과 같은 조건을 따라야 합니다:



저작자표시. 귀하는 원저작자를 표시하여야 합니다.



비영리. 귀하는 이 저작물을 영리 목적으로 이용할 수 없습니다.



변경금지. 귀하는 이 저작물을 개작, 변형 또는 가공할 수 없습니다.

- 귀하는, 이 저작물의 재이용이나 배포의 경우, 이 저작물에 적용된 이용허락조건을 명확하게 나타내어야 합니다.
- 저작권자로부터 별도의 허가를 받으면 이러한 조건들은 적용되지 않습니다.

저작권법에 따른 이용자의 권리는 위의 내용에 의하여 영향을 받지 않습니다.

이것은 [이용허락규약\(Legal Code\)](#)을 이해하기 쉽게 요약한 것입니다.

[Disclaimer](#)

Thesis for the Degree of Master of Agriculture

Panax ginseng leaf extracts exert
anti-obesity effects in high fat
diet-induced obese rats

School of Food science and Biotechnology,
Major of Food Application Technology
The Graduate School

Lee Seul-gi

December 2016

The Graduate School
Kyungpook National University

Panax ginseng leaf extracts exert
anti-obesity effects in high fat
diet-induced obese rats

Lee Seul-gi

School of Food science and Biotechnology,
Major of Food Application Technology
The Graduate School

Supervised by Professor Nam Ju-ock

Approved as a qualified thesis of Lee Seul-gi
for the degree of Master of Agriculture
by the Evaluation Committee

December 2016

Chairman _____ (인)

_____ (인)

The Graduate School Council
Kyungpook National University

Contents

1. Introduction	1
2. Materials and Methods	3
2.1. Preparation of ginseng leaf extracts	3
2.2. Animals	3
2.3. Biochemical analysis	5
2.4. Cell culture and differentiation	5
2.5. Oil Red O staining and triglyceride (TG) assay	5
2.6. Reverse transcription-polymerase chain reaction (RT-PCR)	6
2.7. Western blot	8
2.8. Statistical analysis	8
3. Results	9
3.1. Effects of GD and DL on body weight, adipose tissue mass, and food intake in high-fat diet induced obese rats	9
3.2. Effects of GL and DL on activity levels of serum marker enzymes	12
3.3. Effects of GD and DL on the mRNA and protein expression of adipogenic-related genes in adipose tissues	13
3.4. Effects of GD and DL on proliferation of 3T3-L1 adipocytes	16
3.5. Effects of GD and DL on differentiation and lipid accumulation of 3T3-L1 adipocytes	18
3.6. Effect of GD and DL on the mRNA and protein expression of adipogenic-related genes during 3T3-L1 adipocytes	21
4. Discussion	23

5. References	25
6. Abstract in Korean	28

Tables and Figures Index

1. Table Index

Table 1. The composition of experimental diets	4
Table 2. Sequences and accession numbers for primers used in RT-PCR	7
Table 3. Plasma biochemical values in male rats	13
Table 4. Effects of arctiin on the weights of liver, kidney, heart, spleen, pancreas, and muscle tissue in male rats	13

2. Figure Index

Fig. 1. Effects of GD and DL on body weight, adipose tissue mass, and food intake in high-fat diet induced obese rats	10
Fig. 2. Effects of GD and DL on the mRNA and protein expression of adipogenic-related genes in adipose tissues	15
Fig. 3. Effects of GL and DL on cell viability of 3T3-L1 preadipocytes and differentiated adipocytes	17
Fig. 4. Effects of GL and DL on the differentiation and lipid accumulation of 3T3-L1	19
Fig. 5. Effects of GL and DL on mRNA and protein expressions of adipogenesis-related genes in 3T3-L1 adipocytes	22

1. Introduction

Obesity is one of the major public health problems and is associated with increased metabolic diseases worldwide (8, 16). Obesity is caused by an imbalance between energy intake and energy expenditure, generated excessive adipose tissue (5, 13). Adipose tissue mass can be regulated by adipogenesis process or adipocyte differentiation that has been extensively studied for the treatment of prevention of obesity (7, 12). However, As of September 2013, drug approved by the U.S. Food and Drug Administration (FDA) for the long-term treatment of obesity was only one, a gastrointestinal lipase inhibitor (2). Therefore, the strategy for the prevention of obesity using health functional food is considered to be important, as much as treatment.

The root of ginseng has been traditionally used for the prevention and treatment of various diseases for over 2000 years, however, aerial parts of ginseng did not be effectively utilized. Previous studies reported that the leaf of ginseng is rich in bioactive compounds and contents of some ginsenosides are higher than in the roots (10). Also, stem, leaf, and berry of American ginseng (*Panax quinquefolium*) had anti-obesity activities in vivo (15, 17, 20). However, the anti-obesity effect of leaf extracts of Korean ginseng (*Panax ginseng* CA Meyer) has not yet been investigated.

In the present study, we show that leaf extracts of Korean ginseng were significantly reduce body weight in high-fat diet induced obese rats and inhibit the differentiation of 3T3-L1 adipocytes through regulating the

expression of key transcriptional factors.

2. Materials and Methods

2.1 Preparation of ginseng leaf extracts

Leaf extracts of ginseng were provided by Punggi Ginseng Experimental Station (Yeongju, Korea). Two different types of leaf extract from ginseng were prepared with green leaf (GL) and dried leaf (DL) of ginseng. GL and DL were extracted with distilled water and stored at 4 °C for further use.

2.2. Animals

Five-week old, male Sprague-Dawley (SD) rat were purchased from HyoChang Animal (Seoul, Korea). After 1 week adaptation period, animals were randomly grouped as follows: 1) normal diet (ND) 2) high fat diet (HFD) 3) HFD + GL supplementation (3.3 mg/kg) 4) HFD + DL supplementation (3.3 mg/kg) groups. The composition of high-fat diet is shown in Table 1 and all rats were maintained under a 12 h light-dark cycle. GL and DL were orally administered to the rat everyday throughout the experiment period. Each experimental group consisted of 7 animals. After finishing treatment, blood and various adipose tissues for further analysis.

Ingredients (g/kg of diet)	ND	HFD
Corn starch	400	200
Casein	200	200
Sucrose	200	140
Cellulose	50	50
Mineral mix	35	35
Vitamin mix	10	10
D,L-methionine	3.5	3
Choline	1.5	2
Corn oil	60	210
Lard	40	150

Table 1. The composition of experimental diets.

2.3. Biochemical analysis

On the final day of the experiments, the blood was collected into the serum collection tubes and centrifuged at 3000 rpm for 15 min to obtain the plasma supernatant. Plasma marker levels for nephrotoxicity, hepatotoxicity test, and lipid profiling were performed in Food Analysis Research Center in Suwon Women's University (Korea).

2.4 Cell culture and differentiation

The 3T3-L1 mouse preadipocyte cells were purchased from the Korea Cell Line Bank (KCLB, Seoul, Korea) and cultured in Dulbecco's modified Eagle medium (DMEM; GIBCO, NY, USA) supplemented with 10% bovine calf serum (BCS; GIBCO). To differentiation the 3T3-L1 adipocytes were cultured to post-confluence (designated as day 0) for 2 days and the media was replaced with DMEM containing 10% fetal bovine serum (FBS; GIBCO) and MDI solution (0.5 mM IBMX, 1 μ M DEXA, 0.125 mM indomethacin, and 10 μ g/mL insulin) for 2 days. After, the cells were maintained in media supplemented with 10% FBS and 10 μ g/mL insulin for 4 days. To examine the anti-adipogenic effect of leaf extracts of ginseng, the cells undergoing differentiation were treated with GL and DL during all stages of differentiation.

2.5. Oil Red O staining and triglyceride (TG) assay

Oil Red O solution (ORO) and TG assay were performed as previously described (submitted for publication). Briefly, for staining of lipid droplets, the cells undergo differentiation into mature adipocytes were washed, fixed and stained with Oil red O solution to detect lipid droplets. All images were obtained on microscope (Leica, Germany). TG assay was performed according to the manufacturer's instructions.

2.6. Reverse transcription-polymerase chain reaction (RT-PCR)

RNA was isolated using RNAiso Plus reagent (TaKaRa Bio, Japan). Complementary DNA was synthesized using the Prime-Script RT reagent Kit (TaKaRa Bio) according to the manufacturer's protocol. The specific rat and mouse primers were designed as shown in Table 2. The expressions of mRNA in 3T3-L1 adipocytes and epididymis adipose tissues were normalized to b-actin or GAPDH using Image J software, respectively (National Institutes of Health, Bethesda, MD, USA). The sequences of the designed primers are shown in Table 2.

Gene name	Accession no.	Forward primer	Reverse primer
aP2 (M)	NM_024406	AACACCGAGATTCCTTCAA	TCACGCCTTTCATAACACAT
Adiponectin (M)	NM_009605	ACCTACGACCAGTATCAGGAAAAG	ACTAAGCTGAAAGTGTGTCGACTG
C/EBP α (M)	NM_001287523	TTACAACAGGCCAGGTTTCC	GGCTGGCGACATACAGATCA
C/EBP α (R)	NM_001287579	GCCAAGAAGTCGGTGGATAA	CCTTGACCAAGGAGCTCTCA
C/EBP β (M)	NM_009883	GACTACGCAACACACGTGTAAC	CAAAACCAAAAACATCAACAACCC
C/EBP δ (M)	NM_007679	GATCTGCACGGCCTGTTGTA	CTCCACTGCCACCTGTC
LPL (M)	NM_008509	TCCTCTGACATTTG CAGGTCTATC	TCACGCCTTTCATAACACAT
PPAR γ (M)	AB644275	TTTTCAAGGGTGCCAGTTTC	AATCCTTGGCCCTCTGAGAT
PPAR γ (R)	NM_013124	CTTGCCATATTTATAGCTGTCATTATT	TGTCCTCGATGGGCTTCAC
β -actin (M)	EF095208	GACAACGGCTCCGGCATGTGCAAAG	TTCACGGTTGGCCTTAGGGTTCAG
GAPDH (R)	NM_017008	TCTGACATGCCGCTGGAGAA	TGGAGGCCATGTAGGCCATGA

M, Mouse; R, Rat

Table 2. Sequences and accession numbers for primers used in RT-PCR.

2.7. Western blot

Total proteins were extracted using with pro-prep lysis buffer (iNtRON BIOTECHNOLOGY) containing phosphatase inhibitors and protease inhibitor cocktail. Lysates were centrifuged at 13,000 rpm for 15 min. Protein samples were separated by SDS-PAGE, transferred onto nitrocellulose membranes, blocked using 5% non-fat skim milk and incubated with antibodies against PPAR γ , LPL, Ap2, Adiponectin (Abcam plc, Cambridge, UK), C/EBP α (Cell Signaling Technology, Beverly, MA, USA), and β -actin (Santa Cruz Biotechnology, CA, USA) overnight at 4°C.

2.8. Statistical analysis

Data are expressed as mean \pm SD. Statistical comparisons were performed using 1-way ANOVA. Values of $P < 0.05$ were considered statistically significant (* $P < 0.05$, ** $P < 0.01$).



3. Results

3.1. Effects of GD and DL on body weight, adipose tissue mass, and food intake in high-fat diet induced obese rats



In order to examine the biological effectiveness of GL and DL, we confirmed the anti-obesity effects of GL and DL in diet induced obese rats. Final body weights were slightly lower in GL and DL supplementation rats (HFD+GL and HFD+DL rats), compared to the HFD rats without supplementation (HFD-fed rats), although there was no significant difference (Fig. 1A). Adipose tissues mass (epididymis, abdominal) were significantly decreased in the HFD+GL and HFD+DL rats, compared to the HFD-fed rats (Fig. 1B).

Also, the HFD+GL and HFD+DL rats were decreased the food efficiency ratio, although there was no significant difference (Fig. 1C).

(A)



(B)



(C)




Fig. 1. Effects of GD and DL on body weight, adipose tissue mass, and food intake in high-fat diet induced obese rats. **(A)**The body weight of rat was measured every 3 days (n=7 rat/group). Rats were fed with ND or HFD. The HFD-fed rats were treated with or without GL or DL at a dose of 3.3mg/kg/day.**(B)**Total epididymal WAT and abdominal WAT weight. **(C)** Food intake and Food efficiency ratio. Significantly different from the high-fat diet group (HFD), P values of <0.01(**) or<0.05(*).

3.2. Effects of GL and DL on activity levels of serum marker enzymes

To evaluate the safety of GL and DL, We confirmed the nephrotoxic and hepatotoxic effects of GL and DL. The levels of ALB, TBIL, BUN, and CRE were no significant differences between HFD-fed rats and HFD+GL or HFD+DL rats (Table 3). As expected, GL and DL did not affect to the other organs mass, compared to the HFD rats (Table 4). Furthermore, GL and DL reduces HDL-cholesterol and increases LDL-cholesterol and triglyceride levels.

	ND	HFD	HFD+GL	HFD+DL
ALB (g/dL)	2.2±0.06	2.3±0.18	2.1±0.28	2.2±0.29
Tbil (mg/dl)	0.2±0.02	0.2±0.02	0.2±0.02	0.2±0.02
BUN (mg/dl)	28.0±3.80	20.5±4.91	19.2±5.59	22.9±4.77
CRE (mg/dl)	0.3±0.07	0.3±0.06	0.3±0.01	0.3±0.03
TG (mg/dl)	43.8±27.13	72.2±24.13	24.1±12.04 ^{**}	32.1±17.28 [*]
HDL (mg/dl)	31.0±4.05	34.3±1.83	33.5±5.31	41.3±6.41 [*]
LDL (mg/dl)	10.3±1.51	19.2±3.65	12.5±1.83 ^{**}	12.6±0.94 ^{**}

Table 3. Plasma biochemical values in male rats. Significantly different from the high-fat diet group (HFD), P values of <0.01(**) or <0.05(*).

	ND	HFD	HFD+GL	HFD+DL
Liver weight (g)	4.9±0.51	7.3±1.40	4.7±0.67 [*]	5.2±0.49 [*]
Kidney weight (g)	1.7±0.07	1.9±0.12	1.7±0.11	1.8±0.18
Heart weight (g)	0.6±0.03	0.8±0.07	0.7±0.13	0.7±0.05
Spleen weight (g)	0.3±0.04	0.4±0.03	0.3±0.08	0.3±0.06
Pancreas (g)	0.6±0.10	0.4±0.17	0.8±0.16	1.0±0.10
Muscle weight				
Gastrocnemius (g)	0.1±0.02	0.2±0.01	0.1±0.01	0.2±0.01
Soleus (g)	2.0±0.13	2.0±0.28	1.9±0.32	1.8±0.21

Table 4. Effects of GL and DL on the weights of liver, kidney, heart, spleen, pancreas, and muscle tissue in male rats. Significantly different from the high-fat diet group (HFD), P values of <0.01(**) or <0.05(*).

3.3. Effects of GD and DL on the mRNA and protein expression of adipogenic-related genes in adipose tissues

To investigate the mechanisms underlying the anti-obesity effects of GL and DL, we measured the mRNA and protein expression of adipogenesis-related genes, such as key adipogenic regulators (C/EBP α and PPAR γ). The mRNA and protein expression of C/EBP α and PPAR γ were significantly lower in the epididymal adipose tissues of HFD+GL and HFD+DL rats compared to the HFD-fed rats (Fig. 2A-B). Also, the protein expression of downstream target genes of C/EBP α and PPAR γ , including aP2 and adiponectin, was decreased in both HFD+GL and HFD+DL rats (Fig. 2B).



(A)**(B)**

Fig. 2. Effects of GD and DL on the mRNA and protein expression of adipogenic-related genes in adipose tissues. Rats were fed with a ND or HFD. The HFD-fed rats were treated with or without GL or DL at a dose of 3.3 mg/kg/day. Epididymal WAT was acquired at the final day of the experiments. **(A)** The mRNA expression of PPAR γ , C/EBP α , and GAPDH in epididymal WAT. **(B)** The protein expression of the indicated adipogenic genes in epididymal WAT.

3.4. Effects of GD and DL on proliferation of 3T3-L1 adipocytes

In order to investigate whether the anti-obesity effects of the GL and DL *in vivo* were occurred by inhibiting adipogenesis of adipocytes, we conducted an *in vitro* experiment using 3T3-L1 adipocytes.

First, we determine the cytotoxicity effects of GL and DL. 3T3-L1 adipocytes were treated with GL or DL in preadipocytes or fully differentiated adipocyte (mature adipocytes). GL and DL did not affect to viability of 3T3-L1 preadipocytes and fully differentiated adipocyte (mature adipocytes) (Fig. 3).





Fig. 3. Effects of GL and DL on cell viability of 3T3-L1 preadipocytes and differentiated adipocytes. 3T3-L1 preadipocytes were treated with GL or DL for 48 h or were left untreated. 3T3-L1 preadipocytes were differentiated into adipocytes and treated with GL or DL for entire differentiation period or after induction of differentiation. Cell viability was determined by MTT assay and measured as absorbance at 495 nm.


3.5. Effects of GD and DL on differentiation and lipid accumulation of 3T3-L1 adipocytes

3T3-L1 adipocytes were treated with GL or DL throughout the period of differentiation. DL significantly decreases the lipid accumulation in a dose-dependent manner (Fig. 4A-B). Furthermore, DL at a dose of 0.15 and 0.3 mg/ml markedly decreased the intracellular TG levels by 15.0% and 30.1%, respectively (Fig. 4C). However, GL had little effect on the differentiation of 3T3-L1 adipocytes, suggesting that anti-obesity effects of GL in vivo were caused by specific factor released other organs or increased energy expenditure.

(A)



(B)



(C)




Fig. 4. Effects of GL and DL on the differentiation and lipid accumulation of 3T3-L1. 3T3-L1 preadipocytes were treated with GL and DL for 8 days or were left untreated. (A) The differentiated cells were stained by Oil Red O

and photographed with a microscope at $400 \times$ magnification. (B) The stained cells by Oil Red O measured as absorbance at 495 nm. (C) Intracellular triglyceride concentrations measured as absorbance at 570 nm. Significantly different from the high-fat diet group (HFD), P values of $<0.01(**)$ or $<0.05(*)$.

3.6. Effect of GD and DL on the mRNA and protein expression of adipogenic-related genes during 3T3-L1 adipocytes

To investigate the mechanisms underlying the anti-differentiation effects of GL and DL, we measured the mRNA and protein expression levels of adipogenic-related genes, such as key adipogenic regulators (C/EBPs and PPAR γ). Treatment of DL significantly decreased the mRNA and protein expression levels of C/EBP α , PPAR γ , and adiponectin and also decreased the mRNA expression of fatty acid-binding protein 4 (aP2), lipoprotein lipase (LPL), adiponectin, and C/EPBs (β and δ) (Fig. 5A-B). GL only slightly reduced the mRNA and protein expressions of adipogenesis-related genes. Remarkably, GL greatly increased the protein expression level of PPAR γ than untreated cells.




Fig. 5. Effects of GL and DL on mRNA and protein expressions of adipogenesis-related genes in 3T3-L1 adipocytes. Cells were treated with GL and DL for 8 days during differentiation and total RNA and protein was isolated. **(A)** The mRNA expression of the adipogenic genes in 3T3-L1 cells. **(B)** The protein expression of PPAR γ , C/EBP α , and Adiponectin in 3T3-L1 cells.

4. Discussion

A previous review indicated that the root and leaf-stem extracts of ginseng have similar pharmacological effects (15). Also, a number of studies found that the berry extracts of ginseng could also be used to treat diabetes in obese mice (3, 4, 18, 19).

In this study, we found that two kinds of leaf extracts from ginseng have anti-obesity effects *in vivo* through different mechanisms. GL and DL similarly reduced the body weight gain and adipose tissue mass in high-fat induced obese rats. In the case of DL, these effects were thought to be caused by a decreased differentiation of adipocytes. The expression of mRNA and protein analyses revealed that DL regulated adipogenesis process in 3T3-L1 adipocytes by modulating adipogenic gene expression. On the other hands, GL had little effects on the differentiation of adipocytes. Interestingly, cells treated GL were significantly increased the protein expression of PPAR γ than untreated cells.

PPAR γ is mainly expressed in adipose tissues (WAT, BAT) and a master regulator of adipogenesis as well as a potent metabolic modulator of whole-body lipid metabolism and insulin sensitivity (1, 6, 14). Previous studies report that the mice, a lack of skeletal-muscle PPAR γ , developed excess adiposity and hepatic insulin resistance (1, 11). Another study indicates that PPAR γ hypomorphic mice have mild lipodystrophy because of an efficient compensation by other organs such as, muscle (9). Therefore, from a dynamic point of view, we hypothesises that anti-obesity

effects of GL may be caused by improved glucose homeostasis or energy expenditure through greatly increased expression of PPAR α . However, further experiments such as, glucose homeostasis changes by supplement with GL are required to establish this hypothesis.

In summary, our results demonstrate that the leaf extracts of Korean ginseng have anti-obesity effects by selectively reducing the adipose tissue mass. DL suppressed the adipogenesis of 3T3-L1 adipocytes by modulating the expression of central transcription factors, but not GL. These results were probably due to metabolism modulation or increased energy expenditure. However, additional experimentation is needed to further clarify this issue. To our knowledge, these findings are the first to provide anti-obesity effects of the leaf extracts of Korean ginseng, it can be used to a potential therapeutic agent or dietary supplement against obesity.

5. References

1. Ahmadian M, Suh JM, Hah N, Liddle C, Atkins AR, Downes M, and Evans RM. 2013. PPARgamma signaling and metabolism: the good, the bad and the future. *Nat Med.* **19**:557-566.
2. Annamalai S, Mohanam L, Alwin D, and Prabhu V. 2016. Effect of combination therapy of melatonin and orlistat on high fat diet induced changes in lipid profiles and liver function parameters in serum of rats. *Obesity Medicine.* **2**:41-45.
3. Attele AS, Zhou YP, Xie JT, Wu JA, Zhang L, Dey L, *etal.* 2002. Antidiabetic effects of Panax ginseng berry extract and the identification of an effective component. *Diabetes.* **51**:1851-1858.
4. Cho WC, Chung WS, Lee SK, Leung AW, Cheng CH, and Yue KK. 2006. Ginsenoside Re of Panax ginseng possesses significant antioxidant and antihyperlipidemic efficacies in streptozotocin-induced diabetic rats. *Eur J Pharmacol.* **550**:173-179.
5. Esparza J, Fox C, Harper IT, Bennett PH, Schulz LO, Valencia ME, and Ravussin E. 2000. Daily energy expenditure in Mexican and USA Pima indians: low physical activity as a possible cause of obesity. *Int J Obes Relat Metab Disord.* **24**:55-59.
6. Evans RM, Barish GD, and Wang YX. 2004. PPARs and the complex journey to obesity. *Nat Med.* **10**:355-361.
7. Gregoire FM. 2001. Adipocyte differentiation: from fibroblast to

- endocrine cell. *Exp Biol Med*(Maywood). **226**:997-1002.
8. Kopelman PG. 2000. Obesity as a medical problem. *Nature*.404:635-643.
 13. Spiegelman BM and Flier JS. 2001. Obesity and the regulation of energy balance. *Cell*.104:531-543.
 9. Koutnikova H, Cock TA, Watanabe M, Houten SM, Champy MF, Dierich A, and Auwerx J. 2003. Compensation by the muscle limits the metabolic consequences of lipodystrophy in PPAR gamma hypomorphic mice. *Proc Natl Acad Sci USA*.**100**: 14457-14462.
 10. Lee HS, Lee HJ, Cho HJ, Park SS, Kim JM, and Suh HJ. 2010. Cosmetic Potential of Enzymatic Treated Ginseng Leaf. *Journal of ginseng research*. **34**: 227-236.
 11. Norris AW, Chen L, Fisher SJ, Szanto I, Ristow M, Jozsi AC,*etal*.2003. Muscle-specific PPARgamma-deficient mice develop increased adiposity and insulin resistance but respond to thiazolidinediones. *J Clin Invest*. **112**: 608-618.
 12. Senol-Cosar O, Flach RJ, DiStefano M, Chawla A, Nicoloro S, Straubhaar J,*etal*. 2016. Tenomodulin promotes human adipocyte differentiation and beneficial visceral adipose tissue expansion. *Nat Commun*. **7**: 10686.
 13. Spiegelman BM and Flier JS. 2001. Obesity and the regulation of energy balance. *Cell*. **104**: 531-543.
 14. Tontonoz P and Spiegelman BM. 2008. Fat and beyond: the diverse biology of PPARgamma. *Annu Rev Bio chem*.**77**: 289-312.

15. Wang H, Peng D, and Xie J. 2009. Ginseng leaf-stem: bioactive constituents and pharmacological functions. *Chin Med.* **4**:20.
16. Williams EP, Mesidor M, Winters K, Dubbert PM, and Wyatt SB. 2015. Overweight and Obesity: Prevalence, Consequences, and Causes of a Growing Public Health Problem. *Curr Obes Rep.* **4**:363-370.
17. Xie JT, Mehendale SR, Wang A, Han AH, Wu JA, Osinski J, and Yuan CS. 2004. American ginseng leaf: ginsenoside analysis and hypoglycemic activity. *Pharmacol Res.* **49**:113-117.
18. Xie JT, Wang CZ, Ni M, Wu JA, Mehendale SR, Aung HH, *et al.* 2007. American ginseng berry juice intake reduces blood glucose and body weight in ob/ob mice. *J Food Sci.* **72**: S590-594.
19. Xie JT, Wu JA, Mehendale S, Aung HH, and Yuan CS. 2004. Anti-hyperglycemic effect of the polysaccharides fraction from American ginseng berry extract in ob/ob mice. *Phytomedicine.* **11**: 182-187.
20. Xie JT, Zhou YP, Dey L, Attele AS, Wu JA, Gu M, *et al.* 2002. Ginseng berry reduces blood glucose and body weight in db/db mice. *Phytomedicine.* **9**: 254-258.

고려인삼 잎 추출물에 대한 항비만 효과

이슬기

경북대학교 대학원 식품공학부 식품응용공학전공

(지도교수 남주옥)

초록

최근 한 연구는 인삼의 지상부 (열매, 잎, 줄기) 가 뿌리만큼 다양하고 많은 사포닌을 포함하고 있다고 보고한 바 있으며 이러한 사포닌의 항산화, 항염증 및 항비만 효과는 밝혀진 바 있다. 그러나, 고려인삼 잎 추출물의 항비만효과는 아직 밝혀진 바 없다. 그러므로, 본 연구에서는 고려인삼의 생잎 (Green leaf of ginseng, GL) 및 건조된 잎 (Dried leaf of ginseng, DL) 열수 추출물의 항비만 효과를 생체 내, 외에서 조사하였다. GL 및 DL의 섭취는 고지방식이에 의한 비만유도 쥐의 몸무게 및 부고환 지방, 복강 후 지방 조직을 감소시켰다. 또한, 생체 외에서 DL은 PPAR γ 및 C/EBP α 와 같은 지방생성 주요 조절인자의 발현을 조절함으로써 3T3-L1 지방전구세포의 지방형성과정 (adipogenesis) 을 억제하였다. 반면에, GL은 3T3-L1 세포의 지방형성과정에 미미한 영향을 미쳤으며, 또한 대조군에 비해 PPAR γ 의 발현을 확실하게 증가시켰다. 이러한 결과는 생체 내에서의 GL의 항비만 효과와는 일치하지 않다. 그러므로, GL의 항비만 효과는 지방조직이 아닌 다른 조직에서 분비된 특정 인자로 인한 혹은 생체 에너지 소비율 증가 유도를 통해 도출된 결과라고

사료된다. 본 연구에서는, 고려인삼 잎 추출물의 항비만 효과를 처음으로 검증하였으며, 이를 활용한 비만 치료제 및 기능성식품 개발 가능성을 제공한다.