

Class10

Sabrina Koldinger (A16368238)

Intro to the RCSB Protein Data Bank (PDB)

Here we examine the size and composition of the main database of biomolecular structures- the PDB.

```
pdldata = read.csv("Data_Export_Summary.csv", row.names = 1)
pdldata
```

	X.ray	EM	NMR	Multiple.methods	Neutron	Other
Protein (only)	161,663	12,592	12,337	200	74	32
Protein/Oligosaccharide	9,348	2,167	34	8	2	0
Protein/NA	8,404	3,924	286	7	0	0
Nucleic acid (only)	2,758	125	1,477	14	3	1
Other	164	9	33	0	0	0
Oligosaccharide (only)	11	0	6	1	0	4
Total						
Protein (only)	186,898					
Protein/Oligosaccharide	11,559					
Protein/NA	12,621					
Nucleic acid (only)	4,378					
Other	206					
Oligosaccharide (only)	22					

Q1: What percentage of structures in the PDB are solved by X-Ray and Electron Microscopy.

The data has commas, which may be a problem... treats them like characters.

```
pdldata$X.ray
```

```
[1] "161,663" "9,348" "8,404" "2,758" "164" "11"
```

```
as.numeric(pdbdata$X.ray)
```

Warning: NAs introduced by coercion

```
[1] NA NA NA NA 164 11
```

We could use `gsub()`

```
as.numeric(gsub(",", "", pdbdata$X.ray))
```

```
[1] 161663 9348 8404 2758 164 11
```

```
x=pdbdata$X.ray
commasum= function(x){sum(as.numeric(gsub(",", "", x)))}
commasum(pdbdata$X.ray)
```

```
[1] 182348
```

```
totals=apply(pdbdata, 2, commasum)
totals
```

X.ray	EM	NMR	Multiple.methods
182348	18817	14173	230
Neutron	Other	Total	
79	37	215684	

```
totals/totals["Total"]
```

X.ray	EM	NMR	Multiple.methods
0.8454405519	0.0872433746	0.0657118748	0.0010663749
Neutron	Other	Total	
0.0003662766	0.0001715473	1.0000000000	

About 84.5% of structures are solved by Xray and 8.7% is solved by EM.

Q2: What proportion of structures in the PDB are protein?

```
proteintotal= as.numeric(gsub(",","", pdbdata[1,7]))  
proteintotal/totals["Total"]
```

```
Total  
0.8665362
```

Proteins are about 86.7% (or 0.867) of the total structures in the database.

Visualizing Protein Structure

We will learn the basics of Mol* (mol-star) We will play with 1HSG.




Figure 1: 1HSG




Figure 2: HIV-Pr with inhibitor and the water 308 and ASP 25 molecules highlighted in green

Back to R and working with PDB structures

Predict the dynamics (flexibility) of an important protein:

```
library(bio3d)
hiv= read.pdb("1hsg")
```

Note: Accessing on-line PDB file

```
hiv
```

```
Call: read.pdb(file = "1hsg")
```

```
Total Models#: 1
Total Atoms#: 1686, XYZs#: 5058 Chains#: 2 (values: A B)

Protein Atoms#: 1514 (residues/Calpha atoms#: 198)
Nucleic acid Atoms#: 0 (residues/phosphate atoms#: 0)
```

Non-protein/nucleic Atoms#: 172 (residues: 128)
 Non-protein/nucleic resid values: [HOH (127), MK1 (1)]

Protein sequence:

PQITLWQRPLVTIKIGGQLKEALLDTGADDTVLEEMSLPGRWKPKMIGGIGGFIKVRQYD
 QILIEICGHKAIGTVLVGPTPVNIIGRNLLTQIGCTLNFPQITLWQRPLVTIKIGGQLKE
 ALLDTGADDTVLEEMSLPGRWKPKMIGGIGGFIKVRQYDQILIEICGHKAIGTVLVGPTP
 VNIIGRNLLTQIGCTLNF

+ attr: atom, xyz, seqres, helix, sheet,
 calpha, remark, call

`head(hiv$atom)`

	type	eleno	elety	alt	resid	chain	resno	insert	x	y	z	o	b
1	ATOM	1	N	<NA>	PRO	A	1	<NA>	29.361	39.686	5.862	1	38.10
2	ATOM	2	CA	<NA>	PRO	A	1	<NA>	30.307	38.663	5.319	1	40.62
3	ATOM	3	C	<NA>	PRO	A	1	<NA>	29.760	38.071	4.022	1	42.64
4	ATOM	4	O	<NA>	PRO	A	1	<NA>	28.600	38.302	3.676	1	43.40
5	ATOM	5	CB	<NA>	PRO	A	1	<NA>	30.508	37.541	6.342	1	37.87
6	ATOM	6	CG	<NA>	PRO	A	1	<NA>	29.296	37.591	7.162	1	38.40

	segid	elesy	charge
1	<NA>	N	<NA>
2	<NA>	C	<NA>
3	<NA>	C	<NA>
4	<NA>	O	<NA>
5	<NA>	C	<NA>
6	<NA>	C	<NA>

`pdbseq(hiv)`

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
"P"	"Q"	"I"	"T"	"L"	"W"	"Q"	"R"	"P"	"L"	"V"	"T"	"I"	"K"	"I"	"G"	"G"	"Q"	"L"	"K"
21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40
"E"	"A"	"L"	"L"	"D"	"T"	"G"	"A"	"D"	"D"	"T"	"V"	"L"	"E"	"E"	"M"	"S"	"L"	"P"	"G"
41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60
"R"	"W"	"K"	"P"	"K"	"M"	"I"	"G"	"G"	"I"	"G"	"G"	"F"	"I"	"K"	"V"	"R"	"Q"	"Y"	"D"
61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80
"Q"	"I"	"L"	"I"	"E"	"I"	"C"	"G"	"H"	"K"	"A"	"I"	"G"	"T"	"V"	"L"	"V"	"G"	"P"	"T"

```

81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 1
"P" "V" "N" "I" "I" "G" "R" "N" "L" "L" "T" "Q" "I" "G" "C" "T" "L" "N" "F" "P"
 2  3  4  5  6  7  8  9 10 11 12 13 14 15 16 17 18 19 20 21
"Q" "I" "T" "L" "W" "Q" "R" "P" "L" "V" "T" "I" "K" "I" "G" "G" "Q" "L" "K" "E"
22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41
"A" "L" "L" "D" "T" "G" "A" "D" "D" "T" "V" "L" "E" "E" "M" "S" "L" "P" "G" "R"
42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61
"W" "K" "P" "K" "M" "I" "G" "G" "I" "G" "G" "F" "I" "K" "V" "R" "Q" "Y" "D" "Q"
62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81
"I" "L" "I" "E" "I" "C" "G" "H" "K" "A" "I" "G" "T" "V" "L" "V" "G" "P" "T" "P"
82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99
"V" "N" "I" "I" "G" "R" "N" "L" "L" "T" "Q" "I" "G" "C" "T" "L" "N" "F"

```

Here we will do a Normal Mode Analysis(NMA) to predict functional motions of a kinase.

```
adk= read.pdb("6s36")
```

Note: Accessing on-line PDB file

PDB has ALT records, taking A only, rm.alt=TRUE

```
adk
```

```
Call: read.pdb(file = "6s36")
```

```
Total Models#: 1
```

```
Total Atoms#: 1898, XYZs#: 5694 Chains#: 1 (values: A)
```

```
Protein Atoms#: 1654 (residues/Calpha atoms#: 214)
```

```
Nucleic acid Atoms#: 0 (residues/phosphate atoms#: 0)
```

```
Non-protein/nucleic Atoms#: 244 (residues: 244)
```

```
Non-protein/nucleic resid values: [ CL (3), HOH (238), MG (2), NA (1) ]
```

```
Protein sequence:
```

```

MRIILLGAPGAGKGTQAQFIMEKYGIPQISTGDMRLRAAVKSGSELGKQAKDIMDAGKLV
TDELVIALVKERIAQEDCRNGFLLDGFPRTIPQADAMKEAGINVDYVLEFDVPDELIVDKI
VGRRVHAPSGRVYHVKFNPVKVEGKDDVTGEELTTRKDDQEETVRKRLVEYHQMTAPLIG
YYSKEAEAGNTKYAKVDGTPVAEVRADLEKILG


```

```
+ attr: atom, xyz, seqres, helix, sheet,  
      calpha, remark, call
```

```
modes= nma(adk)
```

```
Building Hessian...      Done in 0.02 seconds.  
Diagonalizing Hessian... Done in 0.448 seconds.
```

```
plot(modes)
```



Make a “movie” called a trajectory of the predicted motions.

```
mktrj(modes, file="adk_m7.pdb")
```

Then can open in Mol*