

	EMERGENCY MEDICAL SERVICES AGENCY	POLICY NUMBER	728
	A Division of the Merced County Department of Public Health	Effective Date	11/2025
Authority: Health and Safety Code, Division 2.5, and California Code of Regulations, Title 22, Division 9, Chapter 4, 1797.220		Initial Date:	10/2011
		Review Date:	01/2028

SEIZURES	
ADULT	PEDIATRIC
BLS	
Assess Vitals / Obtain SpO ₂ / Oxygen - Titrate to SpO ₂ of 94% or higher / Secure Airway - Assist Ventilations if needed.	
Position – If not contraindicated by injuries, place patient in left lateral decubitus position.	
Blood Sugar Check – For patients who have ALOC.	
ALS	
Follow BLS procedures if applicable.	
Obtain 4-Lead ECG.	
Vascular Access - IV/IO rate as appropriate.	
Midazolam – 2 mg IV/IO <u>OR</u> 5 mg IN/IM • Titrate IV/IO in 1 mg increments for seizure control, max dose 5 mg • May repeat IN/IM once after 10 minutes if seizures continue	Midazolam – 0.1 mg/kg IV/IO <u>OR</u> 0.2 mg/kg IN/IM • Max dose 5 mg • May repeat once after 10 minutes if seizures continue
<u>For patients ≥ 24 weeks pregnant:</u> Magnesium Sulfate – 2 gms in 10 ml of NS slow IVP over 2 minutes.	
Special Considerations	
Manage airway with aggressive BLS techniques (suction, airway adjuncts, BVM), if necessary, providers may use approved advanced airway to secure airway.	

Signatures on File

Approved By: Tim Williams
EMS Administrator

Ajinder Singh, MD
EMS Medical Director