

	EMERGENCY MEDICAL SERVICES AGENCY	POLICY NUMBER	732
	A Division of the Merced County Department of Public Health	Effective Date:	11/2025
Authority: Health and Safety Code, Division 2.5, and California Code of Regulations, Title 22, Division 9, Chapter 4, 1797.220		Initial Date:	10/2011
		Review Date:	04/2028

HYPERTHERMIA	
ADULT	PEDIATRIC
BLS	
Assess Vitals / Obtain SpO ₂ / Oxygen - Titrate to SpO ₂ of 94% or higher / Secure Airway - Assist Ventilations if needed. Cooling Measures - Move patient to a cool environment and provide shade when possible, removing constricted or tight clothing including shoes and socks. Blood Sugar Check – For patients with ALOC Temperature – Assess temperature	
Heat Exhaustion	
Fluid replacement with water or non-caffeinated sports drink if patient is conscious and able to swallow without difficulty.	
Heat Stroke	
Splash or sponge patient with water. Place cool packs on neck, axillary, and groin areas. Promote cooling by fanning (evaporation methods).	
ALS	
Follow BLS procedures if applicable. Obtain 4-Lead ECG. Vascular Access - IV/IO rate as appropriate – If needed, Refer to EMS Policy #738 – Non-Traumatic Shock.	
For Seizure Control:	
Midazolam – 2 mg IV/IO <u>OR</u> 5 mg IN/IM - Titrate IV/IO in 1 mg increments for seizure control, max dose 5 mg - May repeat IN/IM once after 10 minutes if seizures continue	Midazolam – 0.1 mg/kg IV/IO <u>OR</u> 0.2 mg/kg IN/IM - Max dose 5 mg - May repeat once after 10 minutes if seizures continue

Signatures on File

Approved By: Tim Williams
EMS Administrator

Ajinder Singh, MD
EMS Medical Director