

	EMERGENCY MEDICAL SERVICES AGENCY	POLICY NUMBER	749
	A Division of the Merced County Department of Public Health	Effective Date	11/2025
Authority: Health and Safety Code, Division 2.5, and California Code of Regulations, Title 22, Division 9, Chapter 4, 1797.220		Initial Date:	11/2018
		Review Date:	07/2028

NEWBORN RESUSCITATION
PEDIATRIC
BLS
Assess Vitals / Obtain SpO ₂ – Right Hand (Pre-Ductal) / Oxygen – See Special Considerations below and apply as appropriate / Secure Airway - Assist Ventilations if needed. Stimulate – Vigorously rub and dry infant. Place infant in sniffing position by padding the shoulders. Suction – Suction airway as needed. Temperature – Keep infant warm.
Heart Rate 100 BPM or Greater
Oxygen – Administer Oxygen as appropriate (See Special Considerations). Assess Vitals – Assess respirations and pulse every minute.
Heart Rate 80 - 100 BPM
Oxygen – Administer oxygen via non-rebreather mask. Assess Vitals - If heart rate remains less than 100 BPM after 30 seconds Begin assisted ventilation with 100% oxygen via bag-valve mask at 40 breaths per minute
Heart Rate 60 - 80 BPM
Oxygen / Secure Airway – Assist ventilation with 100% oxygen via bag-valve mask at 40 breaths per minute Consider placing OPA to secure airway Assess Vitals - If heart rate remains less than 80 BPM after 1 minute Begin CPR at 120 compressions per minute until heart improves > 80 BPM
Heart Rate Less Than 60 BPM
CPR - Begin chest compressions at rate of 120 compressions per minute Oxygen / Secure Airway – Assist ventilation with 100% oxygen via bag-valve mask at 40 breaths per minute Secure airway with OPA until heart rate > 80 BPM, until infant begins to cry, or has gag reflex Assess Vitals – Reassess vitals every minute until heart > 100 BPM

Signatures on File

Approved By: Tim Williams
EMS Administrator

Ajinder Singh, MD
EMS Medical Director

ALS

Follow BLS procedures.
Obtain 4-Lead ECG.

Heart Rate 60 - 80 BPM

Advanced Airway - Secure with an ALS airway if BLS interventions are ineffective.

Heart Rate Less Than 60 BPM

Advanced Airway - Secure with an ALS airway if BLS interventions are ineffective.

Vascular Access – Establish IV/IO, TKO rate.

Epinephrine 1:10,000 - 0.01 mg/kg IV/IO

- Repeat every 5 minutes until heart rate > 80 BPM

Special Considerations

NRP – Target Pre-ductal SPO₂ After Birth

- 1 Minute: 60 – 65%
- 2 Minutes: 65 – 70%
- 3 Minutes: 70 – 75%
- 4 Minutes: 75 – 80%
- 5 Minutes: 80 – 85%
- 10 Minutes: 85 – 95%

NRP Corrective Ventilation Steps; Mr. Sopa:

- M: Mask Adjustment
- R: Reposition the head and neck
- S: Suction the mouth and nose
- O: Open the mouth
- P: Positive pressure ventilation
- A: Advanced Airway

APGAR Sign	0	1	2
Appearance (Skin Color)	Bluish gray or pale skin	Pink body with blue hands and/or feet	Entire body is pink
Pulse (Heart Rate)	No pulse	<100 bpm	At least 100 bpm
Grimace (Reflexes)	No response	Grimaces	Grimaces and coughs; cries; pulls away or sneezes
Activity (Muscle Tone)	Limp or weak movements	Some movement of arms and legs	Active movement
Respiration (Breathing)	Not breathing	Slow or irregular breathing; a weak cry	Normal breathing rate and effort; strong cry