



# Causal prediction for medical decision making: Methods and practice

Introduction to causal prediction  
Ruth Keogh

[Day 1, afternoon]



# Types of investigation

Hernan, Hsu, Healy. A second chance to get causal inference right: a classification of data science tasks. Chance 2019; 32:42-49.

Schmueli. To explain or to predict? Statistical Science 2010; 25: 289-310



Description

Prediction

Causation

# Types of investigation

Hernan, Hsu, Healy. A second chance to get causal inference right: a classification of data science tasks. *Chance* 2019; 32:42-49.

Schmueli. To explain or to predict? *Statistical Science* 2010; 25: 289-310



Description

Prediction

Using data to map some features of the world to other features of the world

Causation

Using data to predict certain features of the world as if the world had been different

# Clinical prediction models

Collins G S, Moons K G M, Dhiman P, Riley R D, Beam A L, Van Calster B et al. **TRIPOD+AI statement: updated guidance for reporting clinical prediction models that use regression or machine learning methods** BMJ 2024; 385 :e078378 doi:10.1136/bmj-2023-078378

- “Prediction models are used across different healthcare settings. They are used to estimate an outcome value or risk. Most models estimate the probability of the presence of a particular health condition (diagnostic) or whether a particular outcome will occur in the future (prognostic).


# Clinical prediction models

Collins G S, Moons K G M, Dhiman P, Riley R D, Beam A L, Van Calster B et al. **TRIPOD+AI statement: updated guidance for reporting clinical prediction models that use regression or machine learning methods** BMJ 2024; 385 :e078378 doi:10.1136/bmj-2023-078378

- “Prediction models are used across different healthcare settings. They are used to estimate an outcome value or risk. Most models estimate the probability of the presence of a particular health condition (diagnostic) or whether a particular outcome will occur in the future (prognostic).
- Their primary use is to support clinical decision making, such as whether to refer patients for further testing, monitor disease deterioration or treatment effects, or initiate treatment or lifestyle changes.”

# Use of prediction models for decision making


Their primary use is to support clinical decision making, such as whether to refer patients for further testing, monitor disease deterioration or treatment effects, or initiate treatment or lifestyle changes



What would my risk of the outcome be  
... if I get referred for more tests?  
...if I do not get referred for more tests?

# Use of prediction models for decision making

Their primary use is to support clinical decision making, such as whether to refer patients for further testing, monitor disease deterioration or treatment effects, or initiate treatment or lifestyle changes




What would my risk of the outcome be  
... if I get referred for more tests?  
...if I do not get referred for more tests?

What would my risk of the outcome be  
... if I initiate treatment?  
...if I do not initiate treatment?

# Use of prediction models for decision making

Their primary use is to support clinical decision making, such as whether to refer patients for further testing, monitor disease deterioration or treatment effects, or initiate treatment or lifestyle changes



What would my risk of the outcome be  
... if I get referred for more tests?  
...if I do not get referred for more tests?


What would my risk of the outcome be  
... if I initiate treatment?  
...if I do not initiate treatment?

What would my risk of the outcome be  
... if I start exercising 3 times a week?  
...if I continue my current activity levels?



# Use of prediction models for decision making

Their primary use is to support clinical decision making, such as whether to refer patients for further testing, monitor disease deterioration or treatment effects, or initiate treatment or lifestyle changes



What would my risk of the outcome be  
... if I get referred for more tests?  
...if I do not get referred for more tests?


What would my risk of the outcome be  
... if I initiate treatment?  
...if I do not initiate treatment?

What would my risk of the outcome be  
... if I start exercising 3 times a week?  
...if I continue my current activity levels?

These are causal questions, so causal thinking and techniques are needed...but prediction models are not usually developed using causal considerations

# Use of prediction models for decision making

Their primary use is to support clinical decision making, such as whether to refer patients for further testing, monitor disease deterioration or treatment effects, or initiate treatment or lifestyle changes



What would my risk of the outcome be  
... if I get referred for more tests?  
...if I do not get referred for more tests?

What would my risk of the outcome be  
... if I initiate treatment?  
...if I do not initiate treatment?

What would my risk of the outcome be  
... if I start exercising 3 times a week?  
...if I continue my current activity levels?

## What do we call this?

- Causal prediction
- Counterfactual prediction
- Prediction under interventions

## Using standard prediction models to inform treatment decisions

Van Geloven, Keogh, van Amsterdam, et al. **The risks of risk assessment: causal blind spots when using prediction models for treatment decisions.** <https://arxiv.org/abs/2402.17366>

Three ways in which standard prediction models incorporate treatments:

1. Including treatment as a predictor in the model
2. Restricting to untreated individuals when developing the model
3. Ignoring treatment (even though it may be used in the population)

# Standard prediction models:

## 1. Including treatment as a predictor in the model

Data used to develop the prediction model (training data):

- Includes treated and untreated individuals.

Prediction model:

- Includes treatment status  $A$  and additional predictors  $X_1, \dots, X_p$

Prediction formula (e.g. if the prediction model was developed using a Cox model):

- $Risk(t|A, X) = 1 - S_0(t)^{\exp(\beta A + \gamma_1 X_1 + \gamma_2 X_2 + \dots + \gamma_p X_p)}$

# Standard prediction models:

## 1. Including treatment as a predictor in the model

Data used to develop the prediction model (training data):

- Includes treated and untreated individuals.

Prediction model:

- Includes treatment status  $A$  and additional predictors  $X_1, \dots, X_p$

Prediction formula (e.g. if the prediction model was developed using a Cox model):

- $Risk(t|A, X) = 1 - S_0(t)^{\exp(\beta A + \gamma_1 X_1 + \gamma_2 X_2 + \dots + \gamma_p X_p)}$

**Predicted risk if I take treatment?**

Set  $A = 1$  and using individual values for  $X_1, \dots, X_p$

**Predicted risk if I do not take treatment?**

Set  $A = 0$  and using individual values for  $X_1, \dots, X_p$

Are these valid  
estimates of risk  
under the two  
treatment strategies?

# Standard prediction models:

## 1. Including treatment as a predictor in the model

Prediction formula (e.g. if the prediction model was developed using a Cox model):

- $Risk(t|A, X) = 1 - S_0(t)^{\exp(\beta A + \gamma_1 X_1 + \gamma_2 X_2 + \dots + \gamma_p X_p)}$

**Predicted risk if I take treatment?**

Set  $A = 1$  and using individual values for  $X_1, \dots, X_p$

**Predicted risk if I do not take treatment?**

Set  $A = 0$  and using individual values for  $X_1, \dots, X_p$

Are these valid  
estimates of risk  
under the two  
treatment strategies?

## Conditions under which these risks have a causal interpretation

- If  $X_1, \dots, X_p$  includes all confounders of the association between treatment and outcome, and it does not include any mediators
- ...and the model is correctly specified

# Standard prediction models:

## 1. Including treatment as a predictor in the model

### Example: PREDICT study

- Used routinely collected primary care data to develop a model for predicting 5-year risk of cardiovascular disease.
- Model includes BP-lowering medication use, and a range of other predictors.


Predictions under interventions for a particular individual who has high BP

- **Setting BP med=1:** 11% risk of cardiovascular disease in the next five years.
  - **Setting BP med=0:** 8% risk of cardiovascular disease in the next five years.
- According to the model, use of blood pressure lowering medication at baseline results in a higher predicted risk of cardiovascular disease.
  - But this is unlikely to have a causal interpretation.

# Standard prediction models:

## 1. Including treatment as a predictor in the model

### A. PREDICT





# Standard prediction models:

## 2. Restricting to treated individuals when developing the model

Data used to develop the prediction model (training data):

- Includes only treated individuals.

Prediction model:

- Includes predictors  $X_1, \dots, X_p$

Prediction formula (e.g. if prediction model was developed using a Cox model):

- $Risk(t|X) = 1 - S_0(t)^{\exp(\gamma_1 X_1 + \gamma_2 X_2 + \dots + \gamma_p X_p)}$

# Standard prediction models:

## 2. Restricting to treated individuals when developing the model

Data used to develop the prediction model (training data):

- Includes only treated individuals.

Prediction model:

- Includes predictors  $X_1, \dots, X_p$

Prediction formula (e.g. if prediction model was developed using a Cox model):

- $Risk(t|X) = 1 - S_0(t)^{\exp(\gamma_1 X_1 + \gamma_2 X_2 + \dots + \gamma_p X_p)}$

**Predicted risk if I take treatment?**

Set  $A = 1$  and using individual values for  $X_1, \dots, X_p$

Are these valid  
estimates of risk  
under treatment?

# Standard prediction models:

## 2. Restricting to treated individuals when developing the model

Data used to develop the prediction model (training data):

- Includes only treated individuals.

Prediction model:

- Includes predictors  $X_1, \dots, X_p$

Prediction formula (e.g. if prediction model was developed using a Cox model):

- $Risk(t|X) = 1 - S_0(t)^{\exp(\gamma_1 X_1 + \gamma_2 X_2 + \dots + \gamma_p X_p)}$

**Predicted risk if I take treatment?**

Set  $A = 1$  and using individual values for  $X_1, \dots, X_p$

Are these valid  
estimates of risk  
under treatment?

We are still conditioning on treatment  
so similar issues arise as in the previous  
scenario

# Standard prediction models:

## 2. Restricting to treated individuals when developing the model

### Example: Cardiac surgery

- The EuroSCORE model predicts mortality after cardiac surgery
- The model was developed only using individuals who actually received surgery
- It has been suggested to use this to inform the decision about whether to proceed with the surgery

### Example: Organ transplantation

- In organ transplantation prediction models have been developed for post-transplant survival
- It has been suggested to use this to inform the decision about whether to list a person for transplant or to inform the allocation of organs

# Standard prediction models:

## 3. Ignoring treatment (even though it may be used in the population)

Data used to develop the prediction model (training data):

- Includes treated and untreated individuals.

Prediction model:

- Predictors  $X_1, \dots, X_p$ , not including treatment status  $A$

Prediction formula:

- $Risk(t|A, X) = 1 - S_0(t)^{\exp(\gamma_1 X_1 + \gamma_2 X_2 + \dots + \gamma_p X_p)}$

# Standard prediction models:

## 3. Ignoring treatment (even though it may be used in the population)

Data used to develop the prediction model (training data):

- Includes treated and untreated individuals.

Prediction model:

- Predictors  $X_1, \dots, X_p$ , not including treatment status  $A$

Prediction formula:

- $Risk(t|A, X) = 1 - S_0(t)^{\exp(\gamma_1 X_1 + \gamma_2 X_2 + \dots + \gamma_p X_p)}$

- This arguably provides estimates of risk under the “current standard of care”
- If risk is “high” then this may support a decision to offer some treatment

# Standard prediction models:

## 3. Ignoring treatment (even though it may be used in the population)

Data used to develop the prediction model (training data):

- Includes treated and untreated individuals.

Prediction model:

- Predictors  $X_1, \dots, X_p$ , not including treatment status  $A$

Prediction formula:

- $Risk(t|A, X) = 1 - S_0(t)^{\exp(\gamma_1 X_1 + \gamma_2 X_2 + \dots + \gamma_p X_p)}$

- This arguably provides estimates of risk under the “current standard of care”
- If risk is “high” then this may support a decision to offer some treatment

Some people in the development data may be at low risk because they received treatment under standard care – that doesn’t mean we should not offer treatment to those with low risk estimates

# Standard prediction models:

## 3. Ignoring treatment (even though it may be used in the population)

Sperrin M, Martin GP, Pate A, Van Staa T, Peek N, Buchan I. Using marginal structural models to adjust for treatment drop-in when developing clinical prediction models. *Statistics in Medicine*. 2018; 37: 4142–4154. <https://doi.org/10.1002/sim.7913>

- The **QRISK model** is used to inform whether a person should be prescribed statins, based on 10-year risk of CVD >10%.
- Restricted to people not taking statins at baseline.
- People who contributed to the model could start statins during follow-up.



# Standard prediction models:

## 3. Ignoring treatment (even though it may be used in the population)

Sperrin M, Martin GP, Pate A, Van Staa T, Peek N, Buchan I. Using marginal structural models to adjust for treatment drop-in when developing clinical prediction models. *Statistics in Medicine*. 2018; 37: 4142–4154. <https://doi.org/10.1002/sim.7913>

- The **QRISK model** is used to inform whether a person should be prescribed statins, based on 10-year risk of CVD >10%.
- Restricted to people not taking statins at baseline.
- People who contributed to the model could start statins during follow-up.
- Interpretation of risk derived from this model is difficult.
- A patient's predicted risk of lower than 10% may be driven by similar patients in the derivation cohort taking statins shortly after baseline.

# Clinical prediction models

Collins G S, Moons K G M, Dhiman P, Riley R D, Beam A L, Van Calster B et al. **TRIPOD+AI statement: updated guidance for reporting clinical prediction models that use regression or machine learning methods** BMJ 2024; 385:e078378 doi:10.1136/bmj-2023-078378

Their primary use is to support clinical decision making, such as whether to refer patients for further testing, monitor disease deterioration or treatment effects, or initiate treatment or lifestyle changes.”

Van Amsterdam, Cina, Didelez et al. **Rapid Response: Prognostic models for decision support need to report their targeted treatments and the expected changes in treatment decisions.** 2024 <https://www.bmj.com/content/385/bmj-2023-078378/rr-1>

# Clinical prediction models

Collins G S, Moons K G M, Dhiman P, Riley R D, Beam A L, Van Calster B et al. **TRIPOD+AI statement: updated guidance for reporting clinical prediction models that use regression or machine learning methods** BMJ 2024; 385:e078378 doi:10.1136/bmj-2023-078378

Their primary use is to support clinical decision making, such as whether to refer patients for further testing, monitor disease deterioration or treatment effects, or initiate treatment or lifestyle changes.”

Van Amsterdam, Cina, Didelez et al. **Rapid Response: Prognostic models for decision support need to report their targeted treatments and the expected changes in treatment decisions.** 2024 <https://www.bmj.com/content/385/bmj-2023-078378/rr-1>

**From tomorrow: how we can develop prediction models in a way that supports clinical decision making through combining causal thinking and methods with prediction modelling techniques**