

Homework 2

Due Mar 11 at 8:29pm	Points 20	Questions 10	Time Limit None
Allowed Attempts Unlimited			

Take the Quiz Again

Attempt History

	Attempt	Time	Score
KEPT	Attempt 2	2 minutes	20 out of 20
LATEST	Attempt 2	2 minutes	20 out of 20
	Attempt 1	13,399 minutes	18.5 out of 20

! Correct answers are hidden.

Score for this attempt: **20** out of 20
Submitted Mar 8 at 9:03pm
This attempt took 2 minutes.

Question 1

2 / 2 pts

[Q28-01] Which of the following vectors are valid quantum states.

☐

$\left(\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}\right)$

☒

$\left(\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}\right)$

☒

$\left(\frac{2}{3}, \frac{1}{3}, \frac{2}{3}, 0\right)$

☒

$\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}\right)$

☐

$\left(\frac{1}{2}, 0, \frac{1}{2}, 0\right)$

☒

$\left(\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}, 0, 0\right)$

Question 2

1 / 1 pts

[Q24-01] What is the quantum state obtained after applying H operator to state $|1\rangle$?

☐ $|+\rangle$
☒ $|-\rangle$
☐ $|0\rangle$
☐ $|1\rangle$

Question 3

2 / 2 pts

[Q28-02] Which of the following vectors are *not* valid quantum states.

☒ $\left(\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}\right)$
☒ $\left(\frac{1}{4}, -\frac{1}{4}, -\frac{1}{4}, \frac{1}{4}\right)$
☐ $(0, -1, 0, 0)$
☐ $\left(\frac{2}{3}, -\frac{1}{3}, \frac{2}{3}, 0\right)$
☐ $\left(\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}, 0, 0\right)$
☒ $\left(\frac{1}{2}, 0, \frac{1}{2}, 0\right)$

Question 4

2 / 2 pts

[Q32-01] If the angle of a real valued qubit is x , what is the probability of observing state ?

$$\cos(x) + \sin(x)$$

☐

$$\cos(x)$$

☐

$$\sin(x)$$

☐

$$\sin^2(x)$$

☐

$$\cos^2(x)$$

☒

Question 5

2 / 2 pts

[Q12-01] We have a circuit with a single qubit created with the code given below. What should replace "#Your code here" if we want to apply a NOT operator to the qubit?

```
q = QuantumRegister(1)
c = ClassicalRegister(1)
qc = QuantumCircuit(q,c)

#Your code here

qc.measure(q[0],c[0])
job = execute(qc,Aer.get_backend('qasm_simulator'),shots=1024)
counts = job.result().get_counts(qc)
print(counts) # counts is a dictionary
```

Question 6

2 / 2 pts

[Q12-02] What will be the output of the code shown below?

```
q = QuantumRegister(1)
c = ClassicalRegister(1)
qc = QuantumCircuit(q,c)

qc.x(q[0])

qc.measure(q[0],c[0])
job = execute(qc,Aer.get_backend('qasm_simulator'),shots=1024)
```

```
counts = job.result().get_counts(qc)
print(counts) # counts is a dictionary
```

- ☐ {'11': 1024}
- ☐ {'1': 502, '0': 522}
- ☐ {'0': 1024}
- ☒ {'1': 1024}

Question 7

2 / 2 pts

[P20-01] In the quantum coin flipping experiment, what happens when a photon is send through the beam splitter?

- ☒ It is reflected with probability 1/2 and transmitted with probability 1/2.
- ☐ It is transmitted with probability 1.
- ☐ It is reflected with probability 1.
- ☐ It is reflected with probability 1/4 and transmitted with probability 3/4.

Question 8

3 / 3 pts

[Q36-01] In the following code, what should replace "#Your code here", if we want to create the quantum state $\frac{1}{\sqrt{2}}(|0\rangle - |1\rangle)$?

(Hint: Think about the vector representation of this state to start with if you are stuck)

```
q2 = QuantumRegister(1,"qreg")
c2 = ClassicalRegister(1,"creg")
qc2 = QuantumCircuit(q2,c2)

qc2.x(q2[0])
#Your code here

qc2.measure(q2,c2)
job = execute(qc2,Aer.get_backend('qasm_simulator'),shots=100)
counts = job.result().get_counts(qc2)
print(counts) # counts is a dictionary
```

```
qc2.h(q2[0])
```

Question 9**2 / 2 pts****[Q12-03]** What will be the output of the following code?

```
q2 = QuantumRegister(2, "qreg")
c2 = ClassicalRegister(2, "creg")
qc2 = QuantumCircuit(q2, c2)

qc2.x(q2[0])

qc2.measure(q2, c2)
job = execute(qc2, Aer.get_backend('qasm_simulator'), shots=100)
counts = job.result().get_counts(qc2)
print(counts) # counts is a dictionary
```

☐ {'10': 1024}☐ {'10': 100}☐ {'11': 100}☒ {'01': 100}**Question 10****2 / 2 pts****[Q24-02]** Suppose a qubit is in state $\begin{pmatrix} 0.43 \\ -0.90 \end{pmatrix}$. What is the amplitude of being in state $|1\rangle$?☒ -0.90☐ 0.81☐ 0.43☐ 0.90

Quiz Score: **20** out of 20

