

Mathematical Methods for Economic Dynamics (ECON 696)

Lecture 5: Static Optimization (Ch. 3)

Example:

$$\max_{x,y,z} f(x,y,z) = x+2z \text{ subject to } \begin{cases} g^1(x,y,z) = x+y+z = 1 \\ g^2(x,y,z) = x^2+y^2+z = 7/4 \end{cases}$$

We setup the Lagrangian:

$$\mathcal{L}(x,y,z) = x + 2z - \lambda_1(x + y + z - 1) - \lambda_2(x^2 + y^2 + z - 7/4).$$

The first-order conditions are:

$$\mathcal{L}'_1 = 1 - \lambda_1 - 2\lambda_2x = 0 \quad (\text{i})$$

$$\mathcal{L}'_2 = -\lambda_1 - 2\lambda_2y = 0 \quad (\text{ii})$$

$$\mathcal{L}'_3 = 2 - \lambda_1 - \lambda_2 = 0 \quad (\text{iii})$$

$$(iii) \Rightarrow \lambda_2 = 2 - \lambda_1$$

Plugging $\lambda_2 = 2 - \lambda_1$ into (ii), we obtain

$$-\lambda_1 - 4y + 2\lambda_1 y = 0 \Leftrightarrow \lambda_1(2y - 1) = 4y \Rightarrow y \neq 1/2.$$

So,

$$\lambda_1 = \frac{4y}{2y - 1}.$$

Plugging $\lambda_1 = 4y/(2y - 1)$ and $\lambda_2 = 2 - \lambda_1$ into (i), we obtain

$$y = 2x - 1/2.$$

Plugging $y = 2x - 1/2$ into the constraints, we translate the constraint equalities into:

$$3x + z = 3/2 \text{ and } 5x^2 - 2x + z = 3/2.$$

$$3x + z = 3/2 \Rightarrow z = -3x + 3/2.$$

Plugging $z = -3x + 3/2$ into $5x^2 - 2x + z = 3/2$, we obtain

$$5x(x - 1) = 0 \Rightarrow x = 0 \text{ or } 1.$$

Case 1: $x = 0$

We obtain $y = -1/2$ and $z = 3/2$. In this case,

$$f(0, -1/2, 3/2) = 3.$$

Case 2: $x = 1$

We obtain $y = 3/2$ and $z = -3/2$. In this case,

$$f(1, 3/2, -3/2) = -2.$$

Hence, $(x, y, z) = (0, -1/2, 3/2)$ is the only possible candidate for the solution. And the associated Lagrange multipliers are $\lambda_1 = \lambda_2 = 1$.

When $\lambda_1 = \lambda_2 = 1$,

$$\mathcal{L}(x, y, z) = -x^2 - y^2 - y + 11/4.$$

The associated Hessian matrix is

$$D^2\mathcal{L} = \begin{pmatrix} -2 & 0 & 0 \\ 0 & -2 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

We check its principal minors: $\Delta_{(1)}^2\mathcal{L} = -2, -2, 0$;

$$\Delta_{(2)}^2\mathcal{L} = \begin{vmatrix} -2 & 0 \\ 0 & 0 \end{vmatrix} = 0; \begin{vmatrix} -2 & 0 \\ 0 & 0 \end{vmatrix} = 0; \begin{vmatrix} -2 & 0 \\ 0 & -2 \end{vmatrix} = 4.$$

and $\Delta_{(3)}^2\mathcal{L} = 0$. Therefore, \mathcal{L} is a concave function given $\lambda_1 = \lambda_2 = 1$. By the sufficiency result of the Lagrangian approach, we confirm $(0, -1/2, 3/2)$ is the solution.

Sufficiency for Local Extreme Points

(Ch. 3.4)

$$\max_{x \in S} f(x) \quad \text{subject to} \quad g^j(x) = 0, j = 1, \dots, m \quad (m < n).$$

In general, we define the bordered Hessian determinants, for $r = m + 1, \dots, n$:

$$B_r(x^*) = \begin{vmatrix} 0 & \dots & 0 & \frac{\partial g^1(x^*)}{\partial x_1} & \dots & \frac{\partial g^1(x^*)}{\partial x_r} \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ 0 & \dots & 0 & \frac{\partial g^m(x^*)}{\partial x_1} & \dots & \frac{\partial g^m(x^*)}{\partial x_r} \\ \frac{\partial g^1(x^*)}{\partial x_1} & \dots & \frac{\partial g^m(x^*)}{\partial x_1} & \mathcal{L}''_{11}(x^*) & \dots & \mathcal{L}''_{1r}(x^*) \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ \frac{\partial g^1(x^*)}{\partial x_r} & \dots & \frac{\partial g^m(x^*)}{\partial x_r} & \mathcal{L}''_{r1}(x^*) & \dots & \mathcal{L}''_{rr}(x^*) \end{vmatrix}$$

The determinant B_r is the $(m + r)$ th leading principal minor of $(m + n) \times (m + n)$ bordered matrix

$$\begin{pmatrix} \mathbf{0}_{m \times m} & \underbrace{Dg(x^*)}_{m \times n} \\ \underbrace{(Dg(x^*))^T}_{n \times m} & \underbrace{D^2\mathcal{L}(x^*)}_{n \times n} \end{pmatrix} = \begin{pmatrix} 0 & \cdots & 0 & \frac{\partial g^1(x^*)}{\partial x_1} & \cdots & \frac{\partial g^1(x^*)}{\partial x_n} \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & 0 & \frac{\partial g^m(x^*)}{\partial x_1} & \cdots & \frac{\partial g^m(x^*)}{\partial x_n} \\ \frac{\partial g^1(x^*)}{\partial x_1} & \cdots & \frac{\partial g^m(x^*)}{\partial x_1} & \mathcal{L}''_{11}(x^*) & \cdots & \mathcal{L}''_{1n}(x^*) \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ \frac{\partial g^1(x^*)}{\partial x_n} & \cdots & \frac{\partial g^m(x^*)}{\partial x_n} & \mathcal{L}''_{n1}(x^*) & \cdots & \mathcal{L}''_{nn}(x^*) \end{pmatrix}.$$

Theorem (Sufficiency for Local Maximum): Suppose there is a point $x^* \in S \subseteq \mathbb{R}^n$ satisfying $g(x^*) = 0$, and a $\lambda = (\lambda_1, \dots, \lambda_m) \in \mathbb{R}^m$ such that

$$\nabla f(x^*) - \sum_{j=1}^m \lambda_j \nabla g^j(x^*) = \underbrace{\mathbf{0}}_{n \times 1} = \begin{pmatrix} 0 \\ \vdots \\ 0 \end{pmatrix}.$$

$(-1)^r B_r(x^*) > 0$ for $r = m+1, \dots, n \Rightarrow x^*$ is the local maximum to (*).

$(-1)^m B_r(x^*) > 0$ for $r = m+1, \dots, n \Rightarrow x^*$ is the local minimum to (*).

Proof: We skip the proof. ■

Example:

$$\max_{x,y,z} f(x,y,z) = x^2 + y^2 + z^2 \text{ subject to } \begin{cases} g^1(x,y,z) = x + 2y + z = 30 \\ g^2(x,y,z) = 2x - y - 3z = 10 \end{cases}$$

We setup the Lagrangian:

$$\mathcal{L}(x,y,z) = x^2 + y^2 + z^2 - \lambda_1(x + 2y + z - 30) - \lambda_2(2x - y - 3z - 10).$$

The first-order conditions are:

$$\begin{aligned} \mathcal{L}'_1 &= 2x - \lambda_1 - 2\lambda_2 = 0 \\ \mathcal{L}'_2 &= 2y - 2\lambda_1 + \lambda_2 = 0 \\ \mathcal{L}'_3 &= 2z - \lambda_1 + 3\lambda_2 = 0. \end{aligned}$$

Considering two equality constraints, the unique solution to the first-order conditions is $(x,y,z) = (10,10,0)$. The associated Lagrange multipliers are $\lambda_1 = 12$ and $\lambda_2 = 4$.

We compute the Bordered Hessian determinant B_3 :

$$B_3(x, y, z) = \begin{vmatrix} 0 & 0 & \partial g^1/\partial x & \partial g^1/\partial y & \partial g^1/\partial z \\ 0 & 0 & \partial g^2/\partial x & \partial g^2/\partial y & \partial g^2/\partial z \\ \partial g^1/\partial x & \partial g^2/\partial x & \mathcal{L}_{xx}'' & \mathcal{L}_{xy}'' & \mathcal{L}_{xz}'' \\ \partial g^1/\partial y & \partial g^2/\partial y & \mathcal{L}_{yx}'' & \mathcal{L}_{yy}'' & \mathcal{L}_{yz}'' \\ \partial g^1/\partial z & \partial g^2/\partial z & \mathcal{L}_{zx}'' & \mathcal{L}_{zy}'' & \mathcal{L}_{zz}'' \end{vmatrix}$$

$$B_3(10, 10, 0) = \begin{vmatrix} 0 & 0 & 1 & 2 & 1 \\ 0 & 0 & 2 & -1 & -3 \\ 1 & 2 & 2 & 0 & 0 \\ 2 & -1 & 0 & 2 & 0 \\ 1 & -3 & 0 & 0 & 2 \end{vmatrix} = 150 > 0$$

This implies $(-1)^2 B_3(10, 10, 0) > 0$. Hence, $(10, 10, 0)$ is the local minimum point.

Optimization with Inequality Constraints

Constrained Optimization: Inequality Constraints (Ch. 3.5)

Consider the following optimization problem with inequality constraints.

$$\max_{x \in S} f(x) \text{ subject to } \begin{cases} g^1(x_1, \dots, x_n) \leq 0 \\ g^2(x_1, \dots, x_n) \leq 0 \\ \vdots \\ g^m(x_1, \dots, x_n) \leq 0 \end{cases}$$

A vector $x = (x_1, \dots, x_n)$ that satisfies all the constraints is called **feasible** (or admissible).

Throughout we assume that $f(\cdot)$ and all the g^j functions are C^1 .

An inequality constraint $g^j(x) \leq 0$ is said to be **binding (active)** at x if $g^j(x) = 0$ and **non-binding (inactive)** at x if $g^j(x) < 0$.

Minimizing $f(x)$ is equivalent to maximizing $-f(x)$. Moreover, $g^j(x) \geq 0$ can be rewritten as $-g^j(x) \leq 0$. In this way, most constrained optimization problems can be expressed as the above form.

We define the Lagrangian exactly the same as before.

$$\mathcal{L}(x) = f(x) - \lambda \cdot g(x) = f(x) - \sum_{j=1}^m \lambda_j g^j(x),$$

where $\lambda = (\lambda_1, \dots, \lambda_m) \in \mathbb{R}^m$ are the Lagrangian multipliers.

Again, the first-order partial derivatives of the Lagrangian are equated to 0:

$$\begin{aligned}\nabla \mathcal{L}(x) &= \nabla f(x) - \sum_{j=1}^m \lambda_j \nabla g^j(x) = 0 \\ \Leftrightarrow \frac{\partial \mathcal{L}(x)}{\partial x_i} &= \frac{\partial f(x)}{\partial x_i} - \sum_{j=1}^m \lambda_j \frac{\partial g^j(x)}{\partial x_i} = 0, \quad \forall i = 1, \dots, n \quad (KT-1)\end{aligned}$$

In addition, we introduce the **complementary slackness conditions**. For all $j = 1, \dots, m$,

$$\lambda_j \geq 0 \text{ and } \lambda_j = 0 \text{ if } g^j(x) < 0 \quad (KT-2)$$

An alternative formulation of KT-2 is that for any $j = 1, \dots, m$,

$$\lambda_j \geq 0 \text{ and } \lambda_j g^j(x) = 0$$

In particular, if $\lambda_j > 0$, one must have $g^j(x) = 0$. However, it is perfectly possible to have both $\lambda_j = 0$ and $g^j(x) = 0$.

Conditions (KT-1) and (KT-2) together are often called the **Kuhn-Tucker conditions**.

Suppose one can find a point x^* at which $f(\cdot)$ is stationary and $g^j(x^*) < 0$ for all $j = 1, \dots, m$. Then, the Kuhn-Tucker conditions will automatically be satisfied by x^* together with all the Lagrangian multipliers $\lambda_j = 0$ for all $j = 1, \dots, m$.

Theorem (Sufficiency for the Kuhn-Tucker Conditions I):

Consider the maximization problem and suppose that x^* is feasible and satisfies conditions (KT-1) and (KT-2). If the Lagrangian $\mathcal{L}(x) = f(x) - \lambda \cdot g(x)$ (with the λ values obtained from solving KT-1 and KT-2) is concave, then x^* is a solution to the maximization problem.

Proof: Since $\mathcal{L}(x, \lambda)$ is concave by assumption and $\nabla \mathcal{L}(x^*) = 0$ from $(KT - 1)$, by the sufficiency result on unconstrained optimization, we have that for all $x \in S$,

$$f(x^*) - \sum_{j=1}^m \lambda_j g^j(x^*) \geq f(x) - \sum_{j=1}^m \lambda_j g_j(x)$$

Rearranging gives the equivalent inequality:

$$f(x^*) - f(x) \geq \sum_{j=1}^m \lambda_j \left(g^j(x^*) - g^j(x) \right).$$

Thus, it suffices to show

$$\sum_{j=1}^m \lambda_j (g^j(x^*) - g^j(x)) \geq 0$$

for every feasible x , because this will imply that x^* solves the maximization problem.

Suppose that $g^j(x^*) < 0$. Then (KT-2) shows that $\lambda_j = 0$.

Suppose that $g^j(x^*) = 0$, we have $\lambda_j(g^j(x^*) - g^j(x)) = -\lambda_j g^j(x) \geq 0$ because x is feasible, i.e., $g^j(x) \leq 0$ and $\lambda_j \geq 0$. Then, we have $\sum_{j=1}^m \lambda_j (g^j(x^*) - g^j(x)) \geq 0$ as desired. ■

Theorem (Sufficiency for the Kuhn-Tucker Conditions II):

Consider the maximization problem and suppose that x^* is feasible and satisfies conditions $(KT - 1)$ and $(KT - 2)$. If $f(\cdot)$ is concave and each $\lambda_j g^j(x)$ (with the λ values obtained from the recipe) is quasiconvex, then x^* is the solution to the maximization problem.

Proof: We want to show that $f(x) - f(x^*) \leq 0$ for all feasible x . Since $f(\cdot)$ is concave, then, according to the first derivative characterization of concavity of $f(\cdot)$,

$$f(x) - f(x^*) \leq \nabla f(x^*) \cdot (x - x^*) \underset{(KT-1)}{\equiv} \sum_{j=1}^m \lambda_j \nabla g^j(x^*) \cdot (x - x^*)$$

where we use the first order condition $(KT - 1)$.

It therefore suffices to show that for all $j = 1, \dots, m$, and all feasible x ,

$$\lambda_j \nabla g^j(x^*) \cdot (x - x^*) \leq 0.$$

The above inequality is satisfied for those j such that $g^j(x^*) < 0$, because then $\lambda_j = 0$ from the complementary slackness condition ($KT - 2$).

For those j such that $g^j(x^*) = 0$, we have $g^j(x) \leq g^j(x^*)$ (because x is feasible), and hence $-\lambda_j g^j(x) \geq -\lambda_j g^j(x^*)$ because $\lambda_j \geq 0$.

Since the function $-\lambda_j g^j(x)$ is quasiconcave (because $\lambda_j g^j(x)$ is quasiconvex), it follows from the first derivative characterization of quasiconcavity that $\nabla(-\lambda_j g^j(x^*)) \cdot (x - x^*) \geq 0$, and thus, $\lambda_j \nabla g^j(x^*) \cdot (x - x^*) \leq 0$. ■

Necessity of the Kuhn and Tucker Condition and Constraint Qualification (Ch. 3.5)

Consider the following maximization problem with inequality constraints.

$$\max_{x \in S} f(x) \text{ subject to } g^j(x) \leq 0, \quad j = 1, \dots, m$$

The following condition plays an important role when one uses the Kuhn-Tucker condition.

Definition: A solution x^* to the constrained maximization problem satisfies the **constraint qualification** if the gradient vectors $\nabla g^j(x^*)$ ($1 \leq j \leq m$) corresponding those constraints that are active (binding) at x^* , are linearly independent.

Theorem (Necessity for Kuhn-Tucker Conditions): Suppose that $x^* = (x_1^*, \dots, x_n^*)$ solves the constrained maximization problem where $f(\cdot)$ and $g^1(\cdot), \dots, g^m(\cdot)$ are C^1 functions. Suppose furthermore that x^* satisfies the constraint qualification. Then, there exist unique numbers $\lambda_1, \dots, \lambda_m$ such that $(KT - 1)$ and $(KT - 2)$ hold at $x = x^*$.

Proof: We skip the proof. ■

Example:

$$\max_{x,y} f(x,y) = xy + x^2 \text{ subject to } \begin{cases} g^1(x,y) = x^2 + y \leq 2 \\ g^2(x,y) = -y \leq -1 \end{cases}$$

We set up the Lagrangian:

$$\mathcal{L}(x,y) = xy + x^2 - \lambda_1(x^2 + y - 2) - \lambda_2(-y + 1).$$

The K-T conditions are:

- (i) $\mathcal{L}'_x = y + 2x - 2\lambda_1 x = 0$
- (ii) $\mathcal{L}'_y = x - \lambda_1 + \lambda_2 = 0$
- (iii) $\lambda_1 \geq 0$, and $\lambda_1(x^2 + y - 2) = 0$
- (iv) $\lambda_2 \geq 0$, and $\lambda_2(-y + 1) = 0$.

Case 1: Both constraints are binding

Then, $x^2 + y = 2$ and $y = 1 \Rightarrow x = \pm 1$ and $y = 1$.

When $x = y = 1$, (i) and (ii) yield $\lambda_1 = 3/2$ and $\lambda_2 = 1/2$.

Thus, $(x, y) = (1, 1)$ with $\lambda_1 = 3/2$ and $\lambda_2 = 1/2$ is a solution candidate.

When $x = -1$ and $y = 1$, (i) and (ii) yield $\lambda_1 = 1/2$ and $\lambda_2 = 3/2$.

Since $f(1, 1) = 2$ and $f(-1, 1) = 0$, $(x, y) = (1, 1)$ with $\lambda_1 = 3/2$ and $\lambda_2 = 1/2$ is a solution candidate.

Case 2: g^1 is binding but g^2 is not.

Then, $x^2 + y = 2$ and $y > 1$. From (iv), $\lambda_2 = 0$. From (ii), $x = \lambda_1$.

Plugging $\lambda_1 = x$ into (i), we obtain

$$y + 2x - 2x^2 = 0 \quad \underbrace{\Rightarrow}_{y=2-x^2} \quad 3x^2 - 2x - 2 = 0.$$

The solutions are $x = (1 \pm \sqrt{7})/3$. But $x = \lambda_1 \geq 0$, only $x = (1 + \sqrt{7})/3$ is admissible.

However, plugging this into $y = 2 - x^2$, we obtain

$$y = \frac{2}{9}(5 - \sqrt{7}) < \frac{2}{9}(5 - 2) = \frac{2}{3} < 1,$$

which contradicts $y > 1$.

So, there is no solution candidate in this case.

Case 3: g^1 is not binding but g^2 is binding.

Then, $x^2 + y < 2$ and $y = 1$. From (iii), $\lambda_1 = 0$.

Then (i) gives $x = -1/2$ and (ii) gives $\lambda_2 = 1/2$.

Thus, $(x, y) = (-1/2, 1)$ with $\lambda_1 = 0$ and $\lambda_2 = 1/2$ is a solution candidate.

Case 4: Both constraints are not binding.

Then, $x^2 + y < 2$ and $y > 1$.

(iii) and (iv) gives $\lambda_1 = \lambda_2 = 0$.

However, plugging $\lambda_1 = \lambda_2 = 0$ into (i) and (ii), we have $y = 0$, which contradicts $y \geq 1$.

So, there is no solution candidate in this case.

The two solution candidates are $f(1, 1) = 2$ and $f(-1/2, 1) = -1/4$ and the objective function is highest at $(1, 1)$.

Weierstraas' theorem applies here.

Define

$$D = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y \leq 2\} \cap \{(x, y) \in \mathbb{R}^2 \mid y \geq 1\}$$

as the set of all feasible points in this question.

We claim that D is a closed set. Basically, we need to show the following two facts: (1) any set involving a weak inequality is closed and (2) the intersection of two closed sets is closed.

Define $B = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 \leq 5\}$ as a closed ball around $(0, 0)$ with radius $\sqrt{5}$.

We claim $D \subseteq B$, which implies that D is a bounded set.

Since $x^2 \leq 2 - y$ and $y \geq 1$, we have $x^2 \leq 1$.

Fix $(x, y) \in D$. Then,

$$\begin{aligned} x^2 + y^2 &\stackrel{\underbrace{\leq}_{x^2 \leq 1}}{\leq} 1 + y^2 \stackrel{\underbrace{\leq}_{y \leq 2 - x^2}}{\leq} 1 + (2 - x^2)^2 \\ &= x^4 - 4x^2 + 5 = (x^2 - 2)^2 + 1 \stackrel{\underbrace{\leq}_{0 \leq x^2 \leq 1}}{\leq} 5. \end{aligned}$$

This implies $(x, y) \in B$. Thus, D is a bounded set.

Since the objective function f is continuous and the set of all feasible points is compact, by Weierstrass theorem, this constrained maximization problem has a solution.

Checking the constraint qualification

The gradients of the two constraints are $\nabla g^1(x, y) = (2x, 1)$ and $\nabla g^2(x, y) = (0, -1)$.

In Case 1, suppose, by way of contradiction, that there exist $\alpha \neq 0$ and a feasible point (x, y) that falls into Case 1 such that $\nabla g^1(x, y) = \alpha \nabla g^2(x, y)$. This implies that $(2x, 1) = (0, -\alpha)$. This equality holds only when $x = 0$ and $\alpha = -1$.

However, in Case 1, we have $x^2 + y = 2$ and $y = 1$, which further imply that $x = \pm 1$. This contradicts the previous conclusion that $x = 0$. Therefore, these two vectors are linearly independent.

In Case 2, only g^1 is binding so that we only look at $\nabla g^1(x, y) = (2x, 1)$, which is linearly independent because it is not the zero vector.

In Case 3, only g^2 is binding so that we only look at $\nabla g^2(x, y) = (0, -1)$, which is linearly independent because it is not the zero vector.

In Case 4, the constraint qualification trivially holds.

So, the constraint qualification holds.

We conclude that $(1, 1)$ is the solution to the problem.

Concave Programming Problems (Ch. 3.9)

The constrained maximization problem is said to be a **concave programming program** when the objective function $f(\cdot)$ is concave and each constraint $g^j(\cdot)$ is a convex function.

$$\max_{x \in S} f(x) \text{ subject to } g(x) \leq 0.$$

One mild condition is needed for the theorem below:

Definition: The constrained optimization problem satisfies the **Slater qualification** if there exists a vector $z \in S$ such that $g^j(z) < 0$ for all $j = 1, \dots, m$.

Theorem (Necessary and Sufficient Condition for Concave Programming): Let $f : S \rightarrow \mathbb{R}$ be a concave C^1 objective function, where $S \subseteq \mathbb{R}^n$ is open and convex. For each $j = 1, \dots, m$, let $g^j : S \rightarrow \mathbb{R}$ be a convex C^1 constraint function. Suppose that the constrained optimization problem satisfies the Slater constraint qualification. Then, $x^* \in S$ is a solution to the constrained optimization problem if and only if there exists $\lambda \in \mathbb{R}^m$ that satisfies:

$$[\text{KT-1}] \quad \nabla f(x^*) - \sum_{j=1}^m \lambda_j \nabla g^j(x^*) = 0; \text{ and}$$

$$[\text{KT-2}] \quad \lambda_j \geq 0 \text{ and } \lambda_j g^j(x^*) = 0 \text{ for each } j = 1, \dots, m.$$

Proof: We skip the proof. ■

Example:

$$\max_{x,y,z} f(x,y,z) = \frac{1}{2}x - y \text{ subject to } \begin{cases} g^1(x,y) = x + e^{-x} + z^2 \leq y \\ g^2(x,y) = -x \leq 0 \end{cases}$$

We setup the Lagrangian:

$$\mathcal{L}(x,y,z) = \frac{1}{2}x - y - \lambda_1(x + e^{-x} - y + z^2) - \lambda_2(-x).$$

The K-T conditions are:

- (i) $\mathcal{L}'_x = \frac{1}{2} - \lambda_1(1 - e^{-x}) + \lambda_2 = 0$
- (ii) $\mathcal{L}'_y = -1 + \lambda_1 = 0$
- (iii) $\mathcal{L}'_z = -2\lambda_1 z = 0$
- (iv) $\lambda_1 \geq 0$, and $\lambda_1(x + e^{-x} + z^2 - y) = 0$
- (v) $\lambda_2 \geq 0$, and $\lambda_2(-x) = 0$.

From (ii), we have $\lambda_1 = 1$. Plugging this into (iii), we obtain $z = 0$.

$\lambda_1 = 1$, $z = 0$, and (iv) together imply $x + e^{-x} = y$.

Plugging $\lambda_1 = 1$ into (i), we obtain

$$e^{-x} = \frac{1}{2} - \lambda_2 \leq \frac{1}{2} \Rightarrow x \geq \ln 2 > 0.$$

Then, $x \geq \ln 2 > 0$ and (v) imply that $\lambda_2 = 0$.

Thus, the only candidate for the solution would be $(x, y, z) = (x, x + e^{-x}, z) = (\ln 2, \ln 2 + 1/2, 0)$.

With $\lambda_1 = 1$ and $\lambda_2 = 0$, the Lagrangian becomes

$$\mathcal{L}(x, y, z) = -\frac{1}{2}x - e^{-x} - z^2.$$

Since the sum of concave functions is concave, $\mathcal{L}(x, y, z)$ is concave as well.

Quasiconcave Programming (Ch. 3.6)

Theorem (Sufficiency for Quasiconcave Programming): Let $f : S \rightarrow \mathbb{R}$ be a quasiconcave C^1 function and $S \subseteq \mathbb{R}^n$ be open and convex. Assume that there exist numbers $\lambda_1, \dots, \lambda_m$ and a vector $x^* \in S$ such that

- (1) x^* is feasible and satisfies [KT-1] and [KT-2];
- (2) $\lambda_j g_j(\mathbf{x})$ is quasiconvex for each $j = 1, \dots, m$; and
- (3) $\nabla f(x^*) \neq 0$ or $f(\cdot)$ is concave.

Then, x^* is a solution to the constrained optimization problem.

Proof: We first prove that for all x ,

$$f(x) > f(x^*) \Rightarrow \nabla f(x^*) \cdot (x - x^*) > 0. \quad (*)$$

Assume $f(x) > f(x^*)$. Due to the continuity of $f(\cdot)$, we choose $\alpha > 0$ small enough so that $f(x - \alpha \nabla f(x^*)) \geq f(x^*)$.

Using the characterization of quasiconcave functions via first derivatives, we have

$$\nabla f(x^*) \cdot (x - \alpha \nabla f(x^*) - x^*) \geq 0$$

This is equivalent to

$$\nabla f(x^*) \cdot (x - x^*) \geq \alpha \|\nabla f(x^*)\|^2 > 0 \quad (\because \nabla f(x^*) \neq 0).$$

This establishes (*).

Let x be any feasible vector, i.e., $g^j(x) \leq 0$ for each $j = 1, \dots, m$.

Let $J = \{j \in \{1, \dots, m\} \mid g^j(x^*) = 0\}$.

$$j \in J \Rightarrow \lambda_j g^j(x) \leq \lambda_j g^j(x^*) \Leftrightarrow -\lambda_j g^j(x) \geq -\lambda_j g^j(x^*).$$

$$j \notin J \Rightarrow \lambda_j = 0 \Rightarrow -\lambda_j g^j(x) \geq -\lambda_j g^j(x^*).$$

Since each $-\lambda_j g^j(x)$ is quasiconcave (because $\lambda_j g^j(x)$ is quasiconvex), we use the characterization of quasiconcave functions via first derivatives,

$$-\lambda_j \nabla g^j(x^*) \cdot (x - x^*) \geq 0 \Leftrightarrow \lambda_j \nabla g^j(x^*) \cdot (x - x^*) \leq 0$$

for each $j \in \{1, \dots, m\}$.

So,

$$0 \geq \sum_{j=1}^m \lambda_j \nabla g^j(x^*) \cdot (x - x^*) \underbrace{=}_{KT-1} \nabla f(x^*) \cdot (x - x^*)$$

By (*), we conclude $f(x) \leq f(x^*)$. Hence, x^* is the solution. ■