

Pre-University Mathematics

Titus Lim

June 2025

Contents

1	Sequences and Series	2
1.1	Sequence	2
1.2	Series	3
1.3	Arithmetic Progression	3
1.3.1	Sum of AP Series	4
1.4	Geometric Progression	5
1.4.1	Sum of GP Series	6
1.5	p-Series	6
1.6	Convergence and Divergence of Series	7
1.6.1	Absolute Convergence	7
1.6.2	Divergence Test	8
1.6.3	Geometric Series Test	8
1.6.4	Comparison Test	9

Introduction

This paper consists of many different pre-university topics primarily from Cambridge A-Level H2 and H3 mathematics such as Fourier, Maclaurin Series, Conics, Equation of Planes etc. It covers the derivation and applications of the different topics. Some examples are also given.

1 Sequences and Series

Consider the following infinite series of elements:

$$\frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \dots, \frac{1}{2^i}, \dots$$

Can we assign this infinite sum to a numerical value? Indeed, we can. We must first understand what are **series** and **sequences**.

1.1 Sequence

A sequence is any number of elements arranged in a specific order.

Example 1. An infinite sequence of ascending odd numbers.

$$1, 3, 5, 7, \dots$$

Sequences can be both infinite and finite. An example of a finite sequence is:

$$2, 4, 8, 16, \dots, n$$

where n is the final element in the sequence.

More formally, the algebraic notation for a sequence is expressed as:

$$a_1, a_2, a_3, \dots, a_n$$

where a_1 is the first term, a_2 is the second term, and a_n is the n -th term.

1.2 Series

A series is the total sum of all elements in a sequence. In the first section, we posed the question of obtaining a numerical value from an infinite sum. What we are really asking is: *How do we evaluate a series?*

With the initial sequence in **Section 1**, let's first change it to a finite sequence. We can rewrite it into this series:

$$S_n = \frac{1}{2} + \frac{1}{4} + \frac{1}{6} + \cdots + \frac{1}{2^n}$$

We can also write the above in summation notation:

$$S_n = \sum_{k=1}^n \frac{1}{2^k}$$

This describes the sum of elements of $\frac{1}{2^k}$ where $k = 1$ to $k = n$.

Suppose we want to find the sum of 10 elements in the series. We can write it as:

$$\begin{aligned} S_{10} &= \sum_{k=1}^{10} \frac{1}{2^k} \\ &= \frac{1}{2^1} + \frac{1}{2^2} + \frac{1}{2^3} + \cdots + \frac{1}{2^{10}} \\ &= 0.99902 \end{aligned}$$

1.3 Arithmetic Progression

The first type of sequence is known as an **arithmetic progression**.

Example 2. Consider the following sequence:

$$2, 5, 8, 11, \dots$$

To obtain the above sequence, we start with the first term and add a fixed value to each term successively.

$$2, 2 + 3 = 5, 5 + 3 = 8, 8 + 3 = 11, \dots$$

This fixed value is called the **common difference**. In the above example, the common difference would be 3.

More formally, an **arithmetic progression** or **AP** is a sequence whereby the difference between the preceding and succeeding terms is common. In algebraic notation, it can be written as:

$$a, a + d, a + 2d, a + 3d, \dots, a + (n - 1)d$$

where a is the first term, d is the common difference, and n is the number of terms in the sequence.

From the above, we can observe that the n -th term is:

$$a_n = a + (n - 1)d$$

1.3.1 Sum of AP Series

An AP series is simply the total sum of terms in a given arithmetic progression. For a general arithmetic progression:

$$a, (a + d), (a + 2d), \dots, (\ell - 2d), (\ell - d), \ell$$

where a is the first term, d is the common difference and ℓ is the n -th term.

The AP series is written as:

$$S_n = a + (a + d) + (a + 2d) + \dots + (\ell - 2d) + (\ell - d) + \ell$$

To derive a general formula for the AP series, let's first reverse the order of S_n :

$$S_n = \ell + (\ell - d) + (\ell - 2d) + \dots + (a + 2d) + (a + d) + a$$

Then, let's add the original AP series to the reverse AP series by vertically summing each term.

$$2S_n = (a + \ell) + (a + \ell) + (a + \ell) + \dots + (a + \ell) + (a + \ell) + (a + \ell)$$

Notice how all the terms have become $(a + \ell)$. There are now n number of $(a + \ell)$ terms. Now, we can make S_n the subject:

$$S_n = \frac{1}{2}n(a + \ell)$$

As seen in the previous section, the $n - th$ term, ℓ , can also be written as $a + (n - 1)d$.

Derivation. Hence, the general formula for the sum of any AP series is:

$$S_n = \frac{1}{2}n(2a + (n - 1)d)$$

All AP series with common difference, $d \neq 0$ are **divergent**. For any AP series,

If $d > 0$, then the sum of any AP series diverges to $+\infty$ as $n \rightarrow \infty$.

If $d < 0$, then the sum of any AP series diverges to $-\infty$ as $n \rightarrow \infty$.

If $d = 0$ and $a \neq 0$, then the sum of any AP series diverges to $+\infty$ as $n \rightarrow \infty$ depending on the sign of a .

If $d = 0$ and $a = 0$, then the sum of any AP series is 0.

1.4 Geometric Progression

The second type of sequence is known as a **geometric progression**.

Example 3. Consider the following sequence:

$$2, 6, 18, 54, \dots$$

To obtain the above sequence, we start with the first term and multiply it by a fixed value to each term successively.

$$2, 2 \times 3 = 6, 6 \times 3 = 18, 18 \times 3 = 54, \dots$$

This fixed value is called the **common ratio**. In the above example, the common ratio would be 3.

More formally, a **geometric progression** or **GP** is a sequence whereby each term is produced by multiplying each preceding term by a constant value. In algebraic notation, it can be written as:

$$a, ar, ar^2, ar^3, \dots, ar^{n-1}$$

where a is the first term, r is the common ratio, and n is the number of terms in the sequence.

From the above, we can observe that the n -th term is:

$$a_n = ar^{n-1}$$

1.4.1 Sum of GP Series

Similar to a series, a GP Series is simply the total sum of terms in a given geometric progression. For a general geometric progression, its GP series can be written as:

$$S_n = a + ar + ar^2 + ar^3 + \dots + ar^{n-1}$$

We can evaluate the series by multiplying both sides by r and then subtracting:

$$\begin{aligned} rS_n &= ar + ar^2 + ar^3 + ar^4 + \dots + ar^{n-1} + ar^n \\ S_n - rS_n &= a - ar^n \\ S_n(1 - r) &= a - ar^n \end{aligned}$$

Now, we can make S_n the subject.

Derivation. Hence, the general formula for the sum of any GP series is:

$$S_n = \frac{a - ar^n}{1 - r} \quad \text{for } r \neq 1$$

A GP series can either be **convergent** or **divergent**.

1.5 p-Series

The **p-Series** is a type of series in the form:

$$\sum_{n=1}^{\infty} \frac{1}{n^p} = 1 + \frac{1}{2^p} + \frac{1}{3^p} + \dots + \frac{1}{n^p} + \dots$$

where p is an exponent and $p \geq 0$.

Interestingly, when $p = 1$, it forms a special type of series called a **Harmonic Series**:

$$\sum_{n=1}^{\infty} \frac{1}{n} = 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n} + \dots$$

In general, a p-series converges if $p > 1$ as the partial sum grows smaller fast enough. Conversely, it diverges if $p \leq 1$.

1.6 Convergence and Divergence of Series

A **convergent** series is any series where the sum converges to a numerical value. For a given series

$$\sum_{n=1}^{\infty} a_n$$

we say it is convergent if the series converges to the sum, $S \in \mathbb{R}$, if

$$\lim_{n \rightarrow \infty} S_n = S$$

where the partial sum

$$S_n = a_1 + a_2 + a_3 + \cdots + a_n = \sum_{k=1}^n a_k$$

Otherwise, the series is **divergent**.

Distributive and associative arithmetic laws still apply for any series.

1.6.1 Absolute Convergence

The series

$$\sum_{n=1}^{\infty} a_n$$

converges **absolutely** if

$$\sum_{n=1}^{\infty} |a_n| \text{ converges}$$

Otherwise, it converges **conditionally** if

$$\sum_{n=1}^{\infty} a_n \text{ converges, but } \sum_{n=1}^{\infty} |a_n| \text{ diverges}$$

In the next few sections, we will explore how we can test if a given series is convergent or divergent.

1.6.2 Divergence Test

The divergence test states that a series is divergent if the limit of the term $a_n \neq 0$ or does not exist.

Definition. The series diverges if

$$\lim_{n \rightarrow \infty} a_n \neq 0 \text{ or}$$
$$\lim_{n \rightarrow \infty} a_n \text{ does not exist}$$

If the limit is 0, more tests would need to be carried out to determine if the series is truly convergent.

Example 4.1 Determine if the following series $\sum_{n=1}^{\infty} \frac{n-1}{n}$ is convergent or divergent.

$$\begin{aligned} & \lim_{n \rightarrow \infty} \frac{n-1}{n} \\ &= \lim_{n \rightarrow \infty} \left(1 - \frac{1}{n}\right) \\ &= \lim_{n \rightarrow \infty} (1 - 0) \\ &= 1 \end{aligned}$$

\therefore Since the limit $\neq 0$, the series is divergent.

1.6.3 Geometric Series Test

The geometric series test applies to any geometric series in the form $\sum_{n=1}^{\infty} ar^{n-1}$.

Definition. The series converges or diverges based on the following

$$\sum_{n=1}^{\infty} ar^{n-1} = \begin{cases} \frac{a}{1-r} & \text{for } |r| < 1 \\ \text{diverges} & \text{for } |r| \geq 1 \end{cases}$$

Example 4.2 Determine if the following series $\sum_{n=1}^{\infty} \frac{5^{n-1}}{3^{n+1}}$ is convergent or divergent. If it

converges, find the value it converges to.

$$\begin{aligned} & \sum_{n=1}^{\infty} \frac{3^{n-1}}{5^{n+1}} \\ &= \sum_{n=1}^{\infty} \frac{1}{25} \left(\frac{3}{5}\right)^{n-1} \\ &= \frac{\frac{1}{25}}{1 - \frac{3}{5}} \\ &= \frac{1}{10} \end{aligned}$$

1.6.4 Comparison Test