

Optimization Methods (CS1.404), Spring 2024

Lecture 17

Naresh Manwani

Machine Learning Lab, IIIT-H

March 14th, 2024



Quasi-Newton Method (Rank One Correction)

- 1: **Initialize:** The starting point \mathbf{x}_0 , Symmetric positive definite matrix B_0 and the tolerance parameter $\epsilon > 0$, Set $k = 0$
- 2: **while** $\|\mathbf{g}_k\| > \epsilon$ **do**
- 3: $\mathbf{d}_k = -B_k \mathbf{g}_k$
- 4: Find α_k along \mathbf{d}_k such that
 - $f(\mathbf{x}_k + \alpha_k \mathbf{d}_k) < f(\mathbf{x}_k)$
 - α_k satisfies Armijo-Wolfe condition
- 5: $\mathbf{x}_{k+1} = \mathbf{x}_k + \alpha_k \mathbf{d}_k$
- 6: Find B_{k+1} as

$$B_{k+1} = B_k + \frac{(\delta_k - B_k \gamma_k)(\delta_k - B_k \gamma_k)^T}{(\delta_k - B_k \gamma_k)^T \gamma_k}$$

- 7: $k = k + 1$
- 8: **end while**
- 9: **Output:** $\mathbf{x}^* = \mathbf{x}_k$, a stationary point of f .

Convergence of SR1 Applied on Quadratic Functions

Theorem1: For Quadratic Functions

Consider SR1 quasi-Newton algorithm applied to a quadratic function with positive definite Hessian H . Then, for any starting point \mathbf{x}_0 and any symmetric starting matrix B_0 , the sequence of iterates \mathbf{x}_k generated by SR1 converges to the minimizer in n -steps, provided

$(\delta_k - B_k \gamma_k)^T \gamma_k \neq 0, \forall k$. Moreover, if n -steps are performed and $\delta_0, \delta_1, \dots, \delta_{n-1}$ are linearly independent, then $B_n = H^{-1}$.

Convergence of SR1 Applied on General Functions

- For general nonlinear functions, the SR1 update continues to generate good Hessian approximations under certain conditions.

Theorem

Suppose that f is twice continuously differentiable, and that its Hessian is bounded and Lipschitz continuous in a neighborhood of a point \mathbf{x}^* . Let \mathbf{x}_k , $k = 0, 1, \dots$ be any sequence of iterates such that $\mathbf{x}_k \rightarrow \mathbf{x}^*$ for some $\mathbf{x}^* \in \mathbb{R}^n$. Suppose in addition that the $|(\boldsymbol{\delta}_k - B_k \boldsymbol{\gamma}_k)^T \boldsymbol{\gamma}_k| \geq r \|\boldsymbol{\delta}_k - B_k \boldsymbol{\gamma}_k\| \cdot \|\boldsymbol{\gamma}_k\|$ holds for all k , for some $r \in (0, 1)$, and that $\boldsymbol{\delta}_0, \boldsymbol{\delta}_1, \dots, \boldsymbol{\delta}_{n-1}$ are linearly independent. Then, the matrices B_k generated by the SR1 updating formula satisfy

$$\lim_{k \rightarrow \infty} \|B_k - \nabla^2 f(\mathbf{x}^*)\| = 0$$

SR1: Some Remarks

- SR1 is a simple and elegant way to use the information gathered during two consecutive iterations to update B_k .
- B_{k+1} is positive definite if $(\delta_k - B_k \gamma_k)^T \gamma_k > 0$ which can not be guaranteed for every k .
- Numerical difficulties happen if $(\delta_k - B_k \gamma_k)^T \gamma_k$ is close to 0.

Rank Two Correction Quasi newton Method

- Given that B_k is symmetric and positive definite matrix, let

$$B_{k+1} = B_k + \alpha \mathbf{u} \mathbf{u}^T + \beta \mathbf{v} \mathbf{v}^T$$

- B_{k+1} is required to satisfy Quasi-Newton condition. Thus,

$$\alpha \mathbf{u}^T \gamma_k \mathbf{u} + \beta \mathbf{v}^T \gamma_k \mathbf{v} = \delta_k - B_k \gamma_k$$

- Letting $\alpha \mathbf{u}^T \gamma_k = \beta \mathbf{v}^T \gamma_k = 1$, we get $\mathbf{u} + \mathbf{v} = \delta_k - B_k \gamma_k$. Taking $\mathbf{u} = \delta_k$ and $\mathbf{v} = -B_k \gamma_k$, we get

$$\alpha^{-1} = \mathbf{u}^T \gamma_k = \delta_k^T \gamma_k$$

$$\beta^{-1} = \mathbf{v}^T \gamma_k = -\gamma_k^T B_k \gamma_k$$

- Therefore, we get the following update for B_{k+1} :

$$B_{k+1} = B_k + \frac{\delta_k^T \delta_k}{\delta_k^T \gamma_k} - \frac{B_k \gamma_k \gamma_k^T B_k}{\gamma_k^T B_k \gamma_k}$$

- This update is called DFP named after Davidson, Fletcher and Powell.

Theorem

Given that B_k is symmetric and positive definite, B_{k+1} generated by DFP is symmetric and positive definite.

DFP Quasi-Newton Algorithm

- 1: **Initialize:** The starting point \mathbf{x}_0 , Symmetric positive definite matrix B_0 and the tolerance parameter $\epsilon > 0$, Set $k = 0$
- 2: **while** $\|\mathbf{g}_k\| > \epsilon$ **do**
- 3: $\mathbf{d}_k = -B_k \mathbf{g}_k$
- 4: Find α_k along \mathbf{d}_k such that
 - $f(\mathbf{x}_k + \alpha_k \mathbf{d}_k) < f(\mathbf{x}_k)$
 - α_k satisfies Armijo-Wolfe condition
- 5: $\mathbf{x}_{k+1} = \mathbf{x}_k + \alpha_k \mathbf{d}_k$
- 6: Find B_{k+1} as

$$B_{k+1} = B_k + \frac{\delta_k^T \delta_k}{\delta_k^T \gamma_k} - \frac{B_k \gamma_k \gamma_k^T B_k}{\gamma_k^T B_k \gamma_k}$$

- 7: $k = k + 1$
- 8: **end while**
- 9: **Output:** $\mathbf{x}^* = \mathbf{x}_k$, a stationary point of f .

- Quasi Newton condition requires $B_{k+1}\gamma_k = \delta_k$ to hold for all k .
- Assume that we want to approximate the Hessian H_{k+1} rather than its inverse. Let $G_{k+1} = B_{k+1}^{-1}$ which approximates Hessian H_{k+1} .
- Then, Quasi-Newton condition would result into $G_{k+1}\delta_k = \gamma_k$.
- Rank two update of G_{k+1} will have the form

$$G_{k+1} = G_k + \alpha \mathbf{u}\mathbf{u}^T + \beta \mathbf{v}\mathbf{v}^T$$

- G_{k+1} is required to satisfy Quasi-Newton condition. Thus,

$$\alpha \mathbf{u}^T \delta_k \mathbf{u} + \beta \mathbf{v}^T \delta_k \mathbf{v} = \gamma_k - G_k \delta_k$$

- Letting $\alpha \mathbf{u}^T \delta_k = \beta \mathbf{v}^T \delta_k = 1$, we get $\mathbf{u} + \mathbf{v} = \gamma_k - G_k \delta_k$. Taking $\mathbf{u} = \gamma_k$ and $\mathbf{v} = -G_k \delta_k$, we get

$$\alpha^{-1} = \mathbf{u}^T \delta_k = \gamma_k^T \delta_k$$

$$\beta^{-1} = \mathbf{v}^T \delta_k = -\delta_k^T G_k \delta_k$$

- Therefore, we get the following update for B_{k+1} :

$$G_{k+1} = G_k + \frac{\gamma_k^T \gamma_k}{\gamma_k^T \delta_k} - \frac{G_k \delta_k \delta_k^T G_k}{\delta_k^T G_k \delta_k}$$

- Next step is to find B_{k+1} as G_{k+1}^{-1} .
- We use Sherman-Morrison Formula to find G_{k+1}^{-1} .
$$(A + \mathbf{u}\mathbf{v}^T)^{-1} = A^{-1} - \frac{A^{-1}\mathbf{u}\mathbf{v}^T A^{-1}}{1 + \mathbf{v}^T A^{-1}\mathbf{u}}$$
- Applying this formula twice to G_{k+1} , we get

$$B_{k+1}^{BFGS} = B_k^{BFGS} + \left(1 + \frac{\gamma_k^T B_k^{BFGS} \gamma_k}{\delta_k^T \gamma_k}\right) \frac{\delta_k \delta_k^T}{\delta_k^T \gamma_k} - \left(\frac{\delta_k \gamma_k^T B_k^{BFGS} + B_k^{BFGS} \gamma_k \delta_k^T}{\delta_k^T \gamma_k}\right)$$

- $B_{k+1}(\phi) = \phi B_{k+1}^{BFGS} + (1 - \phi) B_{k+1}^{DFP}$, where $\phi \in [0, 1]$

Broyden Family Quasi-Newton Algorithm

- 1: **Initialize:** The starting point \mathbf{x}_0 , Symmetric positive definite matrix B_0 and the tolerance parameter $\epsilon > 0$, Set $k = 0$
- 2: **while** $\|\mathbf{g}_k\| > \epsilon$ **do**
- 3: $\mathbf{d}_k = -B_k \mathbf{g}_k$
- 4: Find α_k along \mathbf{d}_k such that
 - $f(\mathbf{x}_k + \alpha_k \mathbf{d}_k) < f(\mathbf{x}_k)$
 - α_k satisfies Armijo-Wolfe condition
- 5: $\mathbf{x}_{k+1} = \mathbf{x}_k + \alpha_k \mathbf{d}_k$
- 6: Find B_{k+1} as

$$B_{k+1}(\phi) = \phi B_{k+1}^{BFGS} + (1 - \phi) B_{k+1}^{DFP}$$

where $\phi \in [0, 1]$.

- 7: $k = k + 1$
- 8: **end while**
- 9: **Output:** $\mathbf{x}^* = \mathbf{x}_k$, a stationary point of f .