

Games in Logic



Goal: check which properties / languages are *definable* in a logic (e.g. FO)

Games in Logic



Goal: check which properties / languages are *definable* in a logic (e.g. FO)

Examples

- Is the property “Universe has even cardinality” definable in $\text{FO}(\mathcal{E})$?
- Is the class of “Strongly connected graphs” definable in $\text{FO}(\mathcal{E})$?
- Is the language $L=(AA)^*$ definable in $\text{FO}(\leq, A, B)$?

Warmup — The evaluation game

Goal: check whether $M \models \phi$

Model-check(φ , M)

```
if  $\varphi = R(x_1, \dots, x_k)$  then
    if  $(x_1^M, \dots, x_k^M) \in R^M$  then
        return true
    else
        return false
else if  $\varphi = \varphi_1 \vee \varphi_2$  then
    return Model-check( $\varphi_1$ ,  $M$ ) OR
        Model-check( $\varphi_2$ ,  $M$ )
else if ...
...
else if  $\varphi = \exists x \varphi'$  then
    for  $u \in U^M$  do
        if Model-check( $\varphi'$ ,  $M[x:=u]$ ) then
            return true
        return false
else if  $\varphi = \forall x \varphi'$  then
    for  $u \in U^M$  do
        if NOT Model-check( $\varphi'$ ,  $M[x:=u]$ ) then
            return false
    return true
```

Warmup – The evaluation game

Goal: check whether $M \models \phi$

Model-check(ϕ , M)

```
if  $\phi = R(x_1, \dots, x_k)$  then
    if  $(x_1^M, \dots, x_k^M) \in R^M$  then
        return true
    else
        return false
else if  $\phi = \varphi_1 \vee \varphi_2$  then
    return Model-check( $\varphi_1$ ,  $M$ ) OR
        Model-check( $\varphi_2$ ,  $M$ )
else if ...
...
else if  $\phi = \exists x \varphi'$  then
    for  $u \in U^M$  do
        if Model-check( $\varphi'$ ,  $M[x:=u]$ ) then
            return true
        return false
else if  $\phi = \forall x \varphi'$  then
    for  $u \in U^M$  do
        if NOT Model-check( $\varphi'$ ,  $M[x:=u]$ ) then
            return false
    return true
```

Construct a two-player game $G_{\phi, M}$
whose winner determines whether
 $M \models \phi$

Warmup — The evaluation game

Goal: check whether $M \models \phi$



Players: Eve, Adam

Construct a two-player game $G_{\phi, M}$
whose winner determines whether
 $M \models \phi$

Warmup — The evaluation game

Goal: check whether $M \models \phi$



Players: Eve, Adam

Arena: subformulas α of ϕ

+ binding $\lambda : \text{FreeVars}(\alpha) \rightarrow U^M$

Construct a two-player game $G_{\phi, M}$
whose winner determines whether
 $M \models \phi$

Warmup — The evaluation game

Goal: check whether $M \models \phi$



Players: Eve, Adam

Arena: subformulas α of ϕ

+ binding $\lambda : \text{FreeVars}(\alpha) \rightarrow U^M$

Construct a two-player game $G_{\phi, M}$ whose winner determines whether $M \models \phi$

(assume w.l.o.g. that ϕ is in Negation Normal Form)

Recall: negations pushed inside

$$\neg \forall \phi \rightsquigarrow \exists \neg \phi \quad \neg \exists \phi \rightsquigarrow \forall \neg \phi$$

$$\neg(\phi \wedge \psi) \rightsquigarrow \neg \phi \vee \neg \psi \quad \dots$$

Warmup — The evaluation game

Goal: check whether $M \models \phi$



Players: Eve, Adam

Arena: subformulas α of ϕ

+ binding $\lambda : \text{FreeVars}(\alpha) \rightarrow U^M$

Construct a two-player game $G_{\phi, M}$ whose winner determines whether $M \models \phi$

At each position (α, λ) of the arena

- if $\alpha = R(x_1, \dots, x_k)$ then game ends, Eve wins if $(\lambda(x_1), \dots, \lambda(x_k)) \in R^M$, otherwise Adam wins
- if $\alpha = \neg R(x_1, \dots, x_k)$ then game ends, Adam wins if $(\lambda(x_1), \dots, \lambda(x_k)) \in R^M$, otherwise Eve wins

Warmup — The evaluation game

Goal: check whether $M \models \phi$



Players: Eve, Adam

Arena: subformulas α of ϕ

+ binding $\lambda : \text{FreeVars}(\alpha) \rightarrow U^M$

Construct a two-player game $G_{\phi, M}$ whose winner determines whether $M \models \phi$

At each position (α, λ) of the arena

- if $\alpha = R(x_1, \dots, x_k)$ then game ends, Eve wins if $(\lambda(x_1), \dots, \lambda(x_k)) \in R^M$, otherwise Adam wins
- if $\alpha = \neg R(x_1, \dots, x_k)$ then game ends, Adam wins if $(\lambda(x_1), \dots, \lambda(x_k)) \in R^M$, otherwise Eve wins
- if $\alpha = \alpha_1 \vee \alpha_2$ then Eve can choose $\alpha' \in \{\alpha_1, \alpha_2\}$, game continues at position (α', λ)

Warmup — The evaluation game

Goal: check whether $M \models \phi$



Players: Eve, Adam

Arena: subformulas α of ϕ

+ binding $\lambda : \text{FreeVars}(\alpha) \rightarrow U^M$

Construct a two-player game $G_{\phi, M}$ whose winner determines whether $M \models \phi$

At each position (α, λ) of the arena

- if $\alpha = R(x_1, \dots, x_k)$ then game ends, Eve wins if $(\lambda(x_1), \dots, \lambda(x_k)) \in R^M$, otherwise Adam wins
- if $\alpha = \neg R(x_1, \dots, x_k)$ then game ends, Adam wins if $(\lambda(x_1), \dots, \lambda(x_k)) \in R^M$, otherwise Eve wins
- if $\alpha = \alpha_1 \vee \alpha_2$ then Eve can choose $\alpha' \in \{\alpha_1, \alpha_2\}$, game continues at position (α', λ)
- if $\alpha = \alpha_1 \wedge \alpha_2$ then Adam can choose $\alpha' \in \{\alpha_1, \alpha_2\}$, game continues at position (α', λ)

Warmup — The evaluation game

Goal: check whether $M \models \phi$



Players: Eve, Adam

Arena: subformulas α of ϕ

+ binding $\lambda : \text{FreeVars}(\alpha) \rightarrow U^M$

Construct a two-player game $G_{\phi, M}$ whose winner determines whether $M \models \phi$

At each position (α, λ) of the arena

- if $\alpha = R(x_1, \dots, x_k)$ then game ends, Eve wins if $(\lambda(x_1), \dots, \lambda(x_k)) \in R^M$, otherwise Adam wins
- if $\alpha = \neg R(x_1, \dots, x_k)$ then game ends, Adam wins if $(\lambda(x_1), \dots, \lambda(x_k)) \in R^M$, otherwise Eve wins
- if $\alpha = \alpha_1 \vee \alpha_2$ then Eve can choose $\alpha' \in \{\alpha_1, \alpha_2\}$, game continues at position (α', λ)
- if $\alpha = \alpha_1 \wedge \alpha_2$ then Adam can choose $\alpha' \in \{\alpha_1, \alpha_2\}$, game continues at position (α', λ)
- if $\alpha = \exists x \alpha'(x)$ then Eve can choose any element $u \in U^M$ to be bound to x , game continues at position (α', λ') where $\lambda' = \lambda[x := u]$

Warmup — The evaluation game

Goal: check whether $M \models \phi$



Players: Eve, Adam

Arena: subformulas α of ϕ

+ binding $\lambda : \text{FreeVars}(\alpha) \rightarrow U^M$

Construct a two-player game $G_{\phi, M}$ whose winner determines whether $M \models \phi$

At each position (α, λ) of the arena

- if $\alpha = R(x_1, \dots, x_k)$ then game ends, Eve wins if $(\lambda(x_1), \dots, \lambda(x_k)) \in R^M$, otherwise Adam wins
- if $\alpha = \neg R(x_1, \dots, x_k)$ then game ends, Adam wins if $(\lambda(x_1), \dots, \lambda(x_k)) \in R^M$, otherwise Eve wins
- if $\alpha = \alpha_1 \vee \alpha_2$ then Eve can choose $\alpha' \in \{\alpha_1, \alpha_2\}$, game continues at position (α', λ)
- if $\alpha = \alpha_1 \wedge \alpha_2$ then Adam can choose $\alpha' \in \{\alpha_1, \alpha_2\}$, game continues at position (α', λ)
- if $\alpha = \exists x \alpha'(x)$ then Eve can choose any element $u \in U^M$ to be bound to x , game continues at position (α', λ') where $\lambda' = \lambda[x := u]$
- if $\alpha = \forall x \alpha'(x)$ then Adam can choose any element $u \in U^M$ to be bound to x , game continues at position (α', λ') where $\lambda' = \lambda[x := u]$

Warmup — The evaluation game

Goal: check whether $M \models \phi$



Players: Eve, Adam

Arena: subformulas α of ϕ (assume w.l.o.g. that ϕ is in Negation Normal Form)
+ binding $\lambda : \text{FreeVars}(\alpha) \rightarrow U^M$

At each position (α, λ) of the arena

- if $\alpha = R(x_1, \dots, x_k)$ then game ends, Eve wins if $(\lambda(x_1), \dots, \lambda(x_k)) \in R^M$, otherwise Adam wins
- if $\alpha = \neg R(x_1, \dots, x_k)$ then game ends, Adam wins if $(\lambda(x_1), \dots, \lambda(x_k)) \in R^M$, otherwise Eve wins
- if $\alpha = \alpha_1 \vee \alpha_2$ then Eve can choose $\alpha' \in \{\alpha_1, \alpha_2\}$, game continues at position (α', λ)
- if $\alpha = \alpha_1 \wedge \alpha_2$ then Adam can choose $\alpha' \in \{\alpha_1, \alpha_2\}$, game continues at position (α', λ)
- if $\alpha = \exists x \alpha'(x)$ then Eve can choose any element $u \in U^M$ to be bound to x , game continues at position (α', λ') where $\lambda' = \lambda[x := u]$
- if $\alpha = \forall x \alpha'(x)$ then Adam can choose any element $u \in U^M$ to be bound to x , game continues at position (α', λ') where $\lambda' = \lambda[x := u]$

Lemma

$M \models \phi$ iff Eve has a strategy to win $G_{\phi, M}$

Definability vs elementary equivalence vs n -equivalence

Notation P : property (i.e. set of models), M : model, ϕ : FO formula

Definability vs elementary equivalence vs n -equivalence

Notation

P : property (i.e. set of models), M : model, ϕ : FO formula

1) P defined by ϕ if for every M

$M \in P$ iff $M \models \phi$

Definability vs elementary equivalence vs n -equivalence

Notation

P : property (i.e. set of models), M : model, ϕ : FO formula

intuitively,

no formula can distinguish M from M'

1) P defined by ϕ if for every M

$M \in P$ iff $M \models \phi$

2) M, M' elementary equivalent if for every ϕ

$M \models \phi$ iff $M' \models \phi$

Definability vs elementary equivalence vs n -equivalence

Notation

P : property (i.e. set of models), M : model, ϕ : FO formula

intuitively,

no formula can distinguish M from M'

1) P defined by ϕ if for every M

$M \in P$ iff $M \models \phi$

2) M, M' elementary equivalent if for every ϕ

$M \models \phi$ iff $M' \models \phi$

Lemma

If there are M, M' such that

$M \in P$, $M' \notin P$, and M, M' elementary equivalent

then P is *not* definable in FO

Definability vs elementary equivalence vs n -equivalence

- 1) P defined by ϕ if for every M $M \in P$ iff $M \models \phi$
- 2) M, M' elementary equivalent if for every ϕ $M \models \phi$ iff $M' \models \phi$

Lemma If there are M, M' such that
 $M \in P$, $M' \notin P$, and M, M' elementary equivalent
then P is *not* definable in FO

Definability vs elementary equivalence vs n -equivalence

- 1) P defined by ϕ if for every M $M \in P$ iff $M \models \phi$
- 2) M, M' elementary equivalent if for every ϕ $M \models \phi$ iff $M' \models \phi$
- 3) ϕ has quantifier rank n if it has at most n nested quantifiers
Example $\phi = \forall x \forall y (\neg E(x,y) \vee (\exists z E(x,z)) \vee (\exists t E(t,y)))$
has quantifier rank 3 (q.r. can be \ll # quantifiers)

Lemma If there are M, M' such that
 $M \in P$, $M' \notin P$, and M, M' elementary equivalent
then P is *not* definable in FO

Definability vs elementary equivalence vs n -equivalence

- 1) P defined by ϕ if for every M $M \in P$ iff $M \models \phi$
- 2) M, M' elementary equivalent if for every ϕ $M \models \phi$ iff $M' \models \phi$
- 3) ϕ has quantifier rank n if it has at most n nested quantifiers
- 4) M, M' are n -equivalent if for every ϕ with q.r. n $M \models \phi$ iff $M' \models \phi$

Lemma If there are M, M' such that
 $M \in P$, $M' \notin P$, and M, M' elementary equivalent
then P is *not* definable in FO

Definability vs elementary equivalence vs n -equivalence

- 1) P defined by ϕ if for every M $M \in P$ iff $M \models \phi$
- 2) M, M' elementary equivalent if for every ϕ $M \models \phi$ iff $M' \models \phi$
- 3) ϕ has quantifier rank n if it has at most n nested quantifiers
- 4) M, M' are n -equivalent if for every ϕ with q.r. n $M \models \phi$ iff $M' \models \phi$

Lemma If for every n there are M, M' such that
 $M \in P$, $M' \notin P$, and M, M' n -equivalent
then P is *not* definable in FO

Definability vs elementary equivalence vs n -equivalence

- 1) P defined by ϕ if for every M $M \in P$ iff $M \models \phi$
- 2) M, M' elementary equivalent if for every ϕ $M \models \phi$ iff $M' \models \phi$
- 3) ϕ has quantifier rank n if it has at most n nested quantifiers
- 4) M, M' are n -equivalent if for every ϕ with q.r. n $M \models \phi$ iff $M' \models \phi$

Lemma If for every n there are M, M' such that
 $M \in P$, $M' \notin P$, and M, M' n -equivalent
then P is *not* definable in FO

New goal: check whether
 M, M' are n -equivalent

Construct a new game $G_{M,M'}$
whose winner determines whether
 M, M' are n -equivalent


Ehrenfeucht-Fraïssé games

Duplicator

Spoiler

Ehrenfeucht-Fraïssé games

M, M' are
 n -equivalent!




Duplicator

Spoiler

Ehrenfeucht-Fraïssé games

M, M' are
 n -equivalent!



Duplicator


No they're
NOT!!!!



Spoiler

Ehrenfeucht-Fraïssé games

M, M' are
 n -equivalent!



Duplicator

No they're
NOT!!!!



Spoiler

Play for n rounds on the arena whose positions are tuples

$$(u_1, \dots, u_i, v_1, \dots, v_i) \in U^M \times \dots \times U^M \times U^{M'} \times \dots \times U^{M'}$$

At each round i

Spoiler chooses an element u_i from U^M (or v_i from $U^{M'}$)


Duplicator responds with an element v_i from $U^{M'}$ (resp. u_i from U^M)

Duplicator survives if $M \models \{u_1, \dots, u_i\}$ and $M' \models \{v_1, \dots, v_i\}$ are isomorphic


Ehrenfeucht-Fraïssé games

Example

How many rounds can **Duplicator** survive ?



M




M'


Ehrenfeucht-Fraïssé games

Example

How many rounds can Duplicator survive ?



M




M'


Ehrenfeucht-Fraïssé games

Example

How many rounds can Duplicator survive ?



M




M'


Ehrenfeucht-Fraïssé games

Example

How many rounds can Duplicator survive ?



M




M'


Ehrenfeucht-Fraïssé games

Example

How many rounds can Duplicator survive ?



M




M'


Ehrenfeucht-Fraïssé games

Example

How many rounds can Duplicator survive ?



M




M'


Ehrenfeucht-Fraïssé games

Example

How many rounds can Duplicator survive ?



M




M'


Ehrenfeucht-Fraïssé games

Example

How many rounds can Duplicator survive ?



M





M'



Ehrenfeucht-Fraïssé games

Example



How many rounds can Duplicator survive ?



Ehrenfeucht-Fraïssé games

Example



How many rounds can Duplicator survive ?



Ehrenfeucht-Fraïssé games

Example



How many rounds can Duplicator survive ?



Ehrenfeucht-Fraïssé games

Example



How many rounds can Duplicator survive ?



Ehrenfeucht-Fraïssé games

Example



How many rounds can Duplicator survive ?



Ehrenfeucht-Fraïssé games

Example


How many rounds can **Duplicator** survive ?




Ehrenfeucht-Fraïssé games

Example

How many rounds can **Duplicator** survive ?



$$M = (\mathbb{Z}, <)$$



$$M' = (\mathbb{R}, <)$$



Ehrenfeucht-Fraïssé games

On non-isomorphic *finite* models, **Spoiler** always wins, eventually...

Why?

Ehrenfeucht-Fraïssé games

On non-isomorphic *finite* models, **Spoiler** always wins, eventually...

Why?

...and he often wins very quickly!



$2^n - 1$ nodes





2^n nodes

Ehrenfeucht-Fraïssé games

On non-isomorphic *finite* models, **Spoiler** always wins, eventually...

Why?

...and he often wins very quickly!





Ehrenfeucht-Fraïssé games

On non-isomorphic *finite* models, **Spoiler** always wins, eventually...

Why?

...and he often wins very quickly!





Ehrenfeucht-Fraïssé games

On non-isomorphic *finite* models, **Spoiler** always wins, eventually...

Why?

...and he often wins very quickly!





Ehrenfeucht-Fraïssé games

On non-isomorphic *finite* models, **Spoiler** always wins, eventually...

Why?

...and he often wins very quickly!





Ehrenfeucht-Fraïssé games

On non-isomorphic *finite* models, **Spoiler** always wins, eventually...

Why?

...and he often wins very quickly!




Ehrenfeucht-Fraïssé games


On non-isomorphic *finite* models, **Spoiler** always wins, eventually...

Why?

...and he often wins very quickly!



$2^n - 1$ nodes



2^n nodes



But there are non-isomorphic *infinite* models where
Duplicator can survive for *arbitrarily many rounds* (but not necessarily forever!)

Ehrenfeucht-Fraïssé games



On non-isomorphic *finite* models, **Spoiler** always wins, eventually...

Why?

...and he often wins very quickly!



But there are non-isomorphic *infinite* models where
Duplicator can survive for *arbitrarily many rounds* (but not necessarily forever!)





Ehrenfeucht-Fraïssé games



On non-isomorphic *finite* models, **Spoiler** always wins, eventually...

Why?

...and he often wins very quickly!



But there are non-isomorphic *infinite* models where
Duplicator can survive for *arbitrarily many rounds* (but not necessarily forever!)





Ehrenfeucht-Fraïssé games



On non-isomorphic *finite* models, **Spoiler** always wins, eventually...

Why?

...and he often wins very quickly!



But there are non-isomorphic *infinite* models where
Duplicator can survive for *arbitrarily many rounds* (but not necessarily forever!)





Ehrenfeucht-Fraïssé games



On non-isomorphic *finite* models, **Spoiler** always wins, eventually...

Why?

...and he often wins very quickly!



But there are non-isomorphic *infinite* models where
Duplicator can survive for *arbitrarily many rounds* (but not necessarily forever!)





Ehrenfeucht-Fraïssé games



On non-isomorphic *finite* models, **Spoiler** always wins, eventually...

Why?

...and he often wins very quickly!



But there are non-isomorphic *infinite* models where
Duplicator can survive for *arbitrarily many rounds* (but not necessarily forever!)





Ehrenfeucht-Fraïssé games



On non-isomorphic *finite* models, **Spoiler** always wins, eventually...

Why?

...and he often wins very quickly!



But there are non-isomorphic *infinite* models where
Duplicator can survive for *arbitrarily many rounds* (but not necessarily forever!)





Ehrenfeucht-Fraïssé games



On non-isomorphic *finite* models, **Spoiler** always wins, eventually...

Why?

...and he often wins very quickly!



But there are non-isomorphic *infinite* models where
Duplicator can survive for *arbitrarily many rounds* (but not necessarily forever!)





Ehrenfeucht-Fraïssé games



On non-isomorphic *finite* models, **Spoiler** always wins, eventually...

Why?

...and he often wins very quickly!





But there are non-isomorphic *infinite* models where
Duplicator can survive for *arbitrarily many rounds* (but not necessarily forever!)



Given n ,
at each round $i = 1, \dots, n$,
pairs of marked nodes in M and M'
must be either at *equal distance*
or at *distance* $\geq 2^{n-i}$



Ehrenfeucht-Fraïssé games

Duplicator can survive for *arbitrarily many rounds* (but not necessarily forever!)



Ehrenfeucht-Fraïssé games



Duplicator can survive for *arbitrarily many rounds* (but not necessarily forever!)



Lemma If for every n there are M, M' such that
 $M \in P$, $M' \notin P$, and M, M' n -equivalent
then P is *not* definable in FO

Ehrenfeucht-Fraïssé games

Duplicator can survive for *arbitrarily many rounds* (but not necessarily forever!)





Lemma If for every n there are M, M' such that
 $M \in P$, $M' \notin P$, and M, M' n -equivalent
then P is *not* definable in FO

Theorem M, M' n -equivalent iff Duplicator survives n rounds in $G_{M, M'}$
[Fraïssé '50, Ehrenfeucht '60]

Ehrenfeucht-Fraïssé games

Duplicator can survive for *arbitrarily many rounds* (but not necessarily forever!)



Lemma If for every n there are M, M' such that
 $M \in P$, $M' \notin P$, and M, M' n -equivalent
then P is *not* definable in FO

Theorem M, M' n -equivalent iff Duplicator survives n rounds in $G_{M,M'}$
[Fraïssé '50, Ehrenfeucht '60]

In particular, $P = \{\text{discrete orders}\}$ is *not* definable in FO,
since $\mathbb{Z} \in P$ and $\{1,2\} \times \mathbb{Z} \notin P$

Ehrenfeucht-Fraïssé games

Lemma If for every n there are M, M' such that
 $M \in P$, $M' \notin P$, and M, M' n -equivalent
then P is *not* definable in FO

Theorem M, M' n -equivalent iff Duplicator survives n rounds in $G_{M, M'}$
[Fraïssé '50, Ehrenfeucht '60]

Ehrenfeucht-Fraïssé games

Lemma If for every n there are M, M' such that $M \in P$, $M' \notin P$, and M, M' n -equivalent then P is *not* definable in FO

Theorem M, M' n -equivalent iff Duplicator survives n rounds in $G_{M, M'}$
[Fraïssé '50, Ehrenfeucht '60]

Example $P = \{\text{connected graphs}\}$. Given n , find $M \in P, M' \notin P$ where Duplicator survives n rounds


Ehrenfeucht-Fraïssé games

Lemma

If for every n there are M, M' such that
 $M \in P$, $M' \notin P$, and M, M' n -equivalent
then P is *not* definable in FO

Theorem M, M' n -equivalent iff Duplicator survives n rounds in $G_{M, M'}$
[Fraïssé '50, Ehrenfeucht '60]

Example $P = \{\text{connected graphs}\}$. Given n , find $M \in P, M' \notin P$ where Duplicator survives n rounds





Ehrenfeucht-Fraïssé games

Lemma

If for every n there are M, M' such that
 $M \in P$, $M' \notin P$, and M, M' n -equivalent
then P is *not* definable in FO

Theorem M, M' n -equivalent iff Duplicator survives n rounds in $G_{M, M'}$
[Fraïssé '50, Ehrenfeucht '60]

Example $P = \{\text{connected graphs}\}$. Given n , find $M \in P, M' \notin P$ where Duplicator survives n rounds





Ehrenfeucht-Fraïssé games

Lemma

If for every n there are M, M' such that
 $M \in P$, $M' \notin P$, and M, M' n -equivalent
then P is *not* definable in FO

Theorem M, M' n -equivalent iff Duplicator survives n rounds in $G_{M, M'}$
[Fraïssé '50, Ehrenfeucht '60]

Example $P = \{\text{connected graphs}\}$. Given n , find $M \in P, M' \notin P$ where Duplicator survives n rounds





Ehrenfeucht-Fraïssé games

Lemma

If for every n there are M, M' such that
 $M \in P$, $M' \notin P$, and M, M' n -equivalent
then P is *not* definable in FO

Theorem M, M' n -equivalent iff Duplicator survives n rounds in $G_{M, M'}$
[Fraïssé '50, Ehrenfeucht '60]

Example $P = \{\text{connected graphs}\}$. Given n , find $M \in P, M' \notin P$ where Duplicator survives n rounds





Ehrenfeucht-Fraïssé games

Lemma

If for every n there are M, M' such that $M \in P$, $M' \notin P$, and M, M' n -equivalent
then P is *not* definable in FO

Theorem M, M' n -equivalent iff Duplicator survives n rounds in $G_{M, M'}$
[Fraïssé '50, Ehrenfeucht '60]

Example $P = \{\text{connected graphs}\}$. Given n , find $M \in P, M' \notin P$ where Duplicator survives n rounds





Ehrenfeucht-Fraïssé games

Lemma

If for every n there are M, M' such that $M \in P$, $M' \notin P$, and M, M' n -equivalent
then P is *not* definable in FO

Theorem M, M' n -equivalent iff Duplicator survives n rounds in $G_{M, M'}$
[Fraïssé '50, Ehrenfeucht '60]

Example $P = \{\text{connected graphs}\}$. Given n , find $M \in P, M' \notin P$ where Duplicator survives n rounds





Ehrenfeucht-Fraïssé games

Lemma

If for every n there are M, M' such that $M \in P$, $M' \notin P$, and M, M' n -equivalent
then P is *not* definable in FO

Theorem M, M' n -equivalent iff Duplicator survives n rounds in $G_{M, M'}$
[Fraïssé '50, Ehrenfeucht '60]

Example $P = \{\text{connected graphs}\}$. Given n , find $M \in P, M' \notin P$ where Duplicator survives n rounds





Ehrenfeucht-Fraïssé games

Lemma

If for every n there are M, M' such that $M \in P$, $M' \notin P$, and M, M' n -equivalent
then P is *not* definable in FO

Theorem M, M' n -equivalent iff Duplicator survives n rounds in $G_{M, M'}$
[Fraïssé '50, Ehrenfeucht '60]

Example $P = \{\text{connected graphs}\}$. Given n , find $M \in P, M' \notin P$ where Duplicator survives n rounds



Ehrenfeucht-Fraïssé games

Lemma If for every n there are M, M' such that $M \in P$, $M' \notin P$, and M, M' n -equivalent then P is *not* definable in FO

Theorem M, M' n -equivalent iff Duplicator survives n rounds in $G_{M, M'}$
[Fraïssé '50, Ehrenfeucht '60]

Example $P = \{\text{even cardinality}\}$. Given n , find $M \in P, M' \notin P$ where Duplicator survives n rounds



$$2^n$$




$$2^n + 1$$

Ehrenfeucht-Fraïssé games


Lemma If for every n there are M, M' such that $M \in P$, $M' \notin P$, and M, M' n -equivalent then P is *not* definable in FO

Theorem M, M' n -equivalent iff Duplicator survives n rounds in $G_{M, M'}$
[Fraïssé '50, Ehrenfeucht '60]

Example $P = \{\text{even cardinality}\}$. Given n , find $M \in P, M' \notin P$ where Duplicator survives n rounds



2^n



$2^n + 1$



Ehrenfeucht-Fraïssé games

Lemma

If for every n there are M, M' such that
 $M \in P$, $M' \notin P$, and M, M' n -equivalent
then P is *not* definable in FO

Theorem M, M' n -equivalent iff Duplicator survives n rounds in $G_{M, M'}$
[Fraïssé '50, Ehrenfeucht '60]

Example $P = \{\text{even cardinality}\}$. Given n , find $M \in P, M' \notin P$ where Duplicator survives n rounds





Ehrenfeucht-Fraïssé games

Lemma If for every n there are M, M' such that $M \in P$, $M' \notin P$, and M, M' n -equivalent then P is *not* definable in FO

Theorem M, M' n -equivalent iff Duplicator survives n rounds in $G_{M, M'}$
[Fraïssé '50, Ehrenfeucht '60]

Example $P = \{\text{even cardinality}\}$. Given n , find $M \in P, M' \notin P$ where Duplicator survives n rounds





Ehrenfeucht-Fraïssé games

Lemma If for every n there are M, M' such that $M \in P$, $M' \notin P$, and M, M' n -equivalent then P is *not* definable in FO

Theorem M, M' n -equivalent iff Duplicator survives n rounds in $G_{M, M'}$
[Fraïssé '50, Ehrenfeucht '60]

Example $P = \{\text{even cardinality}\}$. Given n , find $M \in P, M' \notin P$ where Duplicator survives n rounds





Ehrenfeucht-Fraïssé games

Lemma

If for every n there are M, M' such that
 $M \in P$, $M' \notin P$, and M, M' n -equivalent
then P is *not* definable in FO

Theorem M, M' n -equivalent iff Duplicator survives n rounds in $G_{M, M'}$
[Fraïssé '50, Ehrenfeucht '60]

Example $P = \{\text{even cardinality}\}$. Given n , find $M \in P, M' \notin P$ where Duplicator survives n rounds





Ehrenfeucht-Fraïssé games

Lemma

If for every n there are M, M' such that
 $M \in P$, $M' \notin P$, and M, M' n -equivalent
then P is *not* definable in FO

Theorem M, M' n -equivalent iff Duplicator survives n rounds in $G_{M, M'}$
[Fraïssé '50, Ehrenfeucht '60]

Example $P = \{\text{even cardinality}\}$. Given n , find $M \in P, M' \notin P$ where Duplicator survives n rounds





Ehrenfeucht-Fraïssé games

Lemma

If for every n there are M, M' such that
 $M \in P$, $M' \notin P$, and M, M' n -equivalent
then P is *not* definable in FO

Theorem M, M' n -equivalent iff Duplicator survives n rounds in $G_{M, M'}$
[Fraïssé '50, Ehrenfeucht '60]

Example $P = \{\text{even cardinality}\}$. Given n , find $M \in P, M' \notin P$ where Duplicator survives n rounds



Rule of thumb

If Spoiler plays “close” to previous pebbles,
then Duplicator responds *isomorphically within the corresponding neighbourhoods*
otherwise Duplicator plays “far” but has freedom of choice

Ehrenfeucht-Fraïssé games

Ehrenfeucht-Fraïssé games

Several properties can be proved to be *not* definable in FO:

- connectivity
- parity (i.e. even / odd)
- 2-colorability
- finiteness
- acyclicity

...

Ehrenfeucht-Fraïssé games

Several properties can be proved to be *not* definable in FO:

- connectivity
- parity (i.e. even / odd)
- 2-colorability

- finiteness
- acyclicity

Your turn now!

Ehrenfeucht-Fraïssé games — soundness

Theorem M, M' *n*-equivalent iff Duplicator survives *n* rounds in $G_{M,M'}$
[Fraïssé '50, Ehrenfeucht '60]

Ehrenfeucht-Fraïssé games — soundness

Theorem M, M' *n*-equivalent iff Duplicator survives *n* rounds in $G_{M,M'}$
[Fraïssé '50, Ehrenfeucht '60]

Proof (if direction — from Duplicator's strategy to *n*-equivalence)

Consider ϕ in NNF and with quantifier rank *n*
Suppose $M \models \phi$ and Duplicator survives *n* rounds in $G_{M,M'}$
Need to prove that $M' \models \phi$

Ehrenfeucht-Fraïssé games — soundness

Theorem M, M' *n*-equivalent iff Duplicator survives *n* rounds in $G_{M,M'}$
[Fraïssé '50, Ehrenfeucht '60]

Proof (if direction — from Duplicator's strategy to *n*-equivalence)

Eve wins the *evaluation game* $G_{M,\phi}$

Consider ϕ in NNF and with quantifier rank *n*

Suppose $M \models \phi$ and Duplicator survives *n* rounds in $G_{M,M'}$

Need to prove that $M' \models \phi$

Eve wins the *evaluation game* $G_{M',\phi}$

Ehrenfeucht-Fraïssé games — soundness

Theorem M, M' *n*-equivalent iff Duplicator survives *n* rounds in $G_{M,M'}$
[Fraïssé '50, Ehrenfeucht '60]

Proof (if direction — from Duplicator's strategy to *n*-equivalence)

Eve wins the *evaluation game* $G_{M,\phi}$

Consider ϕ in NNF and with quantifier rank *n*

Suppose $M \models \phi$ and Duplicator survives *n* rounds in $G_{M,M'}$

Need to prove that $M' \models \phi$

Eve wins the *evaluation game* $G_{M',\phi}$

4 cases based on subformula:

- disjunction
- conjunction
- existential quantification
- universal quantification

Ehrenfeucht-Fraïssé games — soundness

Theorem M, M' *n*-equivalent iff Duplicator survives *n* rounds in $G_{M,M'}$
[Fraïssé '50, Ehrenfeucht '60]

Proof (if direction — from Duplicator's strategy to *n*-equivalence)

Eve wins the *evaluation game* $G_{M,\phi}$

Consider ϕ in NNF and with quantifier rank *n*

Suppose $M \models \phi$ and Duplicator survives *n* rounds in $G_{M,M'}$

Need to prove that $M' \models \phi$

Eve wins the *evaluation game* $G_{M',\phi}$

4 cases based on subformula:

- disjunction
- conjunction
- existential quantification
- universal quantification



which disjunct
Eve should I pick ?

evaluation game $G_{M',\phi}$

Ehrenfeucht-Fraïssé games — soundness

Theorem M, M' *n*-equivalent iff Duplicator survives *n* rounds in $G_{M,M'}$
[Fraïssé '50, Ehrenfeucht '60]

Proof (if direction — from Duplicator's strategy to *n*-equivalence)

Eve wins the *evaluation game* $G_{M,\phi}$

Consider ϕ in NNF and with quantifier rank *n*



Suppose $M \models \phi$ and Duplicator survives *n* rounds in $G_{M,M'}$

Need to prove that $M' \models \phi$

Eve wins the *evaluation game* $G_{M',\phi}$

4 cases based on subformula:

- disjunction
- conjunction
- existential quantification
- universal quantification



Ehrenfeucht-Fraïssé games — soundness

Theorem M, M' *n*-equivalent iff Duplicator survives *n* rounds in $G_{M,M'}$
[Fraïssé '50, Ehrenfeucht '60]

Proof (if direction — from Duplicator's strategy to *n*-equivalence)

Eve wins the *evaluation game* $G_{M,\phi}$

Consider ϕ in NNF and with quantifier rank *n*



Suppose $M \models \phi$ and Duplicator survives *n* rounds in $G_{M,M'}$

Need to prove that $M' \models \phi$

Eve wins the *evaluation game* $G_{M',\phi}$

4 cases based on subformula:

- disjunction
- conjunction
- existential quantification
- universal quantification



Ehrenfeucht-Fraïssé games

Theorem M, M' *n*-equivalent iff Duplicator survives *n* rounds in $G_{M,M'}$
[Fraïssé '50, Ehrenfeucht '60]

Proof (if direction — from Duplicator's strategy to *n*-equivalence)

Eve wins the *evaluation game* $G_{M,\phi}$

Consider ϕ in NNF and with quantifier rank *n*

Suppose $M \models \phi$ and Duplicator survives *n* rounds in $G_{M,M'}$

Need to prove that $M' \models \phi$

Eve wins the *evaluation game* $G_{M',\phi}$

4 cases based on subformula:

- disjunction
- conjunction
- existential quantification
- universal quantification



Ehrenfeucht-Fraïssé games

Theorem M, M' *n*-equivalent iff Duplicator survives *n* rounds in $G_{M,M'}$
[Fraïssé '50, Ehrenfeucht '60]

Proof (if direction — from Duplicator's strategy to *n*-equivalence)

Eve wins the *evaluation game* $G_{M,\phi}$

Consider ϕ in NNF and with quantifier rank *n*

Suppose $M \models \phi$ and Duplicator survives *n* rounds in $G_{M,M'}$

Need to prove that $M' \models \phi$

Eve wins the *evaluation game* $G_{M',\phi}$

4 cases based on subformula:

- disjunction
- conjunction
- existential quantification
- universal quantification

evaluation game $G_{M,\phi}$



Adam picked a conjunct
evaluation game $G_{M',\phi}$

Ehrenfeucht-Fraïssé games

Theorem M, M' *n*-equivalent iff Duplicator survives *n* rounds in $G_{M,M'}$
[Fraïssé '50, Ehrenfeucht '60]

Proof (if direction — from Duplicator's strategy to *n*-equivalence)

Eve wins the *evaluation game* $G_{M,\phi}$

Consider ϕ in NNF and with quantifier rank *n*



Suppose $M \models \phi$ and Duplicator survives *n* rounds in $G_{M,M'}$

Need to prove that $M' \models \phi$

Eve wins the *evaluation game* $G_{M',\phi}$

4 cases based on subformula:

- disjunction
- conjunction
- existential quantification
- universal quantification



Ehrenfeucht-Fraïssé games

Theorem M, M' *n*-equivalent iff Duplicator survives *n* rounds in $G_{M,M'}$
[Fraïssé '50, Ehrenfeucht '60]

Proof (if direction — from Duplicator's strategy to *n*-equivalence)

True wins the *evaluation game* $G_{M,\phi}$

Consider ϕ in NNF and with quantifier rank *n*

Suppose $M \models \phi$ and Duplicator survives *n* rounds in $G_{M,M'}$

Need to prove that $M' \models \phi$

True wins the *evaluation game* $G_{M',\phi}$

4 cases based on subformula:

- disjunction
- conjunction
- existential quantification
- universal quantification



Ehrenfeucht-Fraïssé games

Theorem M, M' *n*-equivalent iff Duplicator survives *n* rounds in $G_{M,M'}$
[Fraïssé '50, Ehrenfeucht '60]

Proof (if direction — from Duplicator's strategy to *n*-equivalence)

True wins the *evaluation game* $G_{M,\phi}$

Consider ϕ in NNF and with quantifier rank *n*


Suppose $M \models \phi$ and Duplicator survives *n* rounds in $G_{M,M'}$

Need to prove that $M' \models \phi$

True wins the *evaluation game* $G_{M',\phi}$

4 cases based on subformula:

- disjunction
- conjunction
- existential quantification
- universal quantification



Ehrenfeucht-Fraïssé games

Theorem M, M' *n*-equivalent iff Duplicator survives *n* rounds in $G_{M,M'}$
[Fraïssé '50, Ehrenfeucht '60]

Proof (if direction — from Duplicator's strategy to *n*-equivalence)

True wins the *evaluation game* $G_{M,\phi}$

Consider ϕ in NNF and with quantifier rank *n*



Suppose $M \models \phi$ and Duplicator survives *n* rounds in $G_{M,M'}$

Need to prove that $M' \models \phi$

True wins the *evaluation game* $G_{M',\phi}$

4 cases based on subformula:

- disjunction
- conjunction
- existential quantification
- universal quantification



Ehrenfeucht-Fraïssé games

Theorem M, M' *n*-equivalent iff Duplicator survives *n* rounds in $G_{M,M'}$
[Fraïssé '50, Ehrenfeucht '60]

Proof (if direction — from Duplicator's strategy to *n*-equivalence)

True wins the evaluation game $G_{M,\phi}$

Consider ϕ in NNF and with quantifier rank *n*



Suppose $M \models \phi$ and Duplicator survives *n* rounds in $G_{M,M'}$

Need to prove that $M' \models \phi$

True wins the evaluation game $G_{M',\phi}$

4 cases based on subformula:

- disjunction
- conjunction
- existential quantification
- universal quantification



Ehrenfeucht-Fraïssé games

Theorem M, M' *n*-equivalent iff Duplicator survives *n* rounds in $G_{M,M'}$
[Fraïssé '50, Ehrenfeucht '60]

Proof (if direction — from Duplicator's strategy to *n*-equivalence)


True wins the evaluation game $G_{M,\phi}$

Consider ϕ in NNF and with quantifier rank *n*

Suppose $M \models \phi$ and Duplicator survives *n* rounds in $G_{M,M'}$


Need to prove that $M' \models \phi$

True wins the evaluation game $G_{M',\phi}$



4 cases based on subformula:

- disjunction
- conjunction
- existential quantification
- universal quantification



Ehrenfeucht-Fraïssé games

Theorem M, M' *n*-equivalent iff Duplicator survives *n* rounds in $G_{M,M'}$
[Fraïssé '50, Ehrenfeucht '60]

Proof (if direction — from Duplicator's strategy to *n*-equivalence)


True wins the evaluation game $G_{M,\phi}$

Consider ϕ in NNF and with quantifier rank *n*

Suppose $M \models \phi$ and Duplicator survives *n* rounds in $G_{M,M'}$

Need to prove that $M' \models \phi$

True wins the evaluation game $G_{M',\phi}$



4 cases based on subformula:

- disjunction
- conjunction
- existential quantification
- universal quantification

Ehrenfeucht-Fraïssé games

Theorem M, M' *n*-equivalent iff Duplicator survives *n* rounds in $G_{M,M'}$
[Fraïssé '50, Ehrenfeucht '60]

Proof (if direction — from Duplicator's strategy to *n*-equivalence)

True wins the *evaluation game* $G_{M,\phi}$

Consider ϕ in NNF and with quantifier rank *n*

Suppose $M \models \phi$ and Duplicator survives *n* rounds in $G_{M,M'}$


Need to prove that $M' \models \phi$

True wins the *evaluation game* $G_{M',\phi}$

4 cases based on subformula:

- disjunction
- conjunction
- existential quantification
- universal quantification

EF game $G_{M,M'}$



Ehrenfeucht-Fraïssé games

Theorem M, M' *n*-equivalent iff Duplicator survives *n* rounds in $G_{M,M'}$
[Fraïssé '50, Ehrenfeucht '60]

Proof (if direction — from Duplicator's strategy to *n*-equivalence)

True wins the *evaluation game* $G_{M,\phi}$

Consider ϕ in NNF and with quantifier rank *n*

Suppose $M \models \phi$ and Duplicator survives *n* rounds in $G_{M,M'}$



Need to prove that $M' \models \phi$

True wins the *evaluation game* $G_{M',\phi}$

4 cases based on subformula:

- disjunction
- conjunction
- existential quantification
- universal quantification

EF game $G_{M,M'}$



Ehrenfeucht-Fraïssé games

Theorem M, M' *n*-equivalent iff Duplicator survives *n* rounds in $G_{M,M'}$
[Fraïssé '50, Ehrenfeucht '60]

Proof (if direction — from Duplicator's strategy to *n*-equivalence)

True wins the *evaluation game* $G_{M,\phi}$

Consider ϕ in NNF and with quantifier rank *n*

Suppose $M \models \phi$ and Duplicator survives *n* rounds in $G_{M,M'}$

Need to prove that $M' \models \phi$

True wins the *evaluation game* $G_{M',\phi}$

4 cases based on subformula:

- disjunction
- conjunction
- existential quantification
- universal quantification



Ehrenfeucht-Fraïssé games

Theorem M, M' *n*-equivalent iff Duplicator survives *n* rounds in $G_{M,M'}$
[Fraïssé '50, Ehrenfeucht '60]

Proof (if direction — from Duplicator's strategy to *n*-equivalence)

True wins the *evaluation game* $G_{M,\phi}$

Consider ϕ in NNF and with quantifier rank *n*


Suppose $M \models \phi$ and Duplicator survives *n* rounds in $G_{M,M'}$

Need to prove that $M' \models \phi$

True wins the *evaluation game* $G_{M',\phi}$

4 cases based on subformula:

- disjunction
- conjunction
- existential quantification
- universal quantification



Ehrenfeucht-Fraïssé games

Theorem M, M' *n*-equivalent iff Duplicator survives *n* rounds in $G_{M,M'}$
[Fraïssé '50, Ehrenfeucht '60]

Proof (if direction — from Duplicator's strategy to *n*-equivalence)

True wins the evaluation game $G_{M,\phi}$

Consider ϕ in NNF and with quantifier rank *n*


Suppose $M \models \phi$ and Duplicator survives *n* rounds in $G_{M,M'}$

Need to prove that $M' \models \phi$

True wins the evaluation game $G_{M',\phi}$

4 cases based on subformula:

- disjunction
- conjunction
- existential quantification
- universal quantification



Ehrenfeucht-Fraïssé games — completeness

Theorem M, M' *n*-equivalent iff Duplicator survives *n* rounds in $G_{M,M'}$
[Fraïssé '50, Ehrenfeucht '60]

Ehrenfeucht-Fraïssé games — completeness

Theorem M, M' *n*-equivalent iff Duplicator survives *n* rounds in $G_{M,M'}$
[Fraïssé '50, Ehrenfeucht '60]

Proof (only if direction — from *n*-equivalence to Duplicator's strategy)

Ehrenfeucht-Fraïssé games — completeness

Theorem M, M' *n-equivalent* iff Duplicator survives *n* rounds in $G_{M,M'}$
[Fraïssé '50, Ehrenfeucht '60]

Proof (only if direction — from *n*-equivalence to Duplicator's strategy)

Suppose M, M' are *n-equivalent* (i.e. *for every* ϕ of q.r. *n*, $M \models \phi$ iff $M' \models \phi$)
Need to construct a **strategy** for Duplicator ...

Ehrenfeucht-Fraïssé games — completeness

Theorem M, M' *n-equivalent* iff Duplicator survives *n* rounds in $G_{M,M'}$
[Fraïssé '50, Ehrenfeucht '60]

Proof (only if direction — from *n*-equivalence to Duplicator's strategy)

Suppose M, M' are *n-equivalent* (i.e. *for every* ϕ of q.r. *n*, $M \models \phi$ iff $M' \models \phi$)
Need to construct a **strategy** for Duplicator ...

... To exploit the hypothesis, we'd better have a formula ϕ
 ϕ needs to be *strong enough to cover all cases for Duplicator*

Ehrenfeucht-Fraïssé games — completeness


Theorem M, M' *n-equivalent* iff Duplicator survives *n* rounds in $G_{M,M'}$
[Fraïssé '50, Ehrenfeucht '60]

Proof (only if direction — from *n*-equivalence to Duplicator's strategy)

Suppose M, M' are *n-equivalent* (i.e. *for every* ϕ of q.r. *n*, $M \models \phi$ iff $M' \models \phi$)
Need to construct a **strategy** for Duplicator ...

... To exploit the hypothesis, we'd better have a formula ϕ
 ϕ needs to be *strong enough to cover all cases for Duplicator*

We need a
HUGE



Hintikka formulas

Level- n Hintikka formula of M = strongest formula (up to logical equivalence) of quantifier rank n that holds on M

$$\phi_M^n$$

≈ “FO theory of M relativised to q.r. n ”

Hintikka formulas

Level- n Hintikka formula of M = strongest formula (up to logical equivalence) of quantifier rank n that holds on M

$$\phi_M^n$$

≈ “FO theory of M relativised to q.r. n ”

Constructed inductively on n :

$$\phi_M^0 = \bigwedge_{\substack{\alpha \text{ atomic} \\ M \vDash \alpha}} \alpha \quad \wedge \quad \bigwedge_{\substack{\alpha \text{ atomic} \\ M \not\vDash \alpha}} \neg \alpha$$

Hintikka formulas

Level- n Hintikka formula of M = strongest formula (up to logical equivalence) of quantifier rank n that holds on M

$$\phi_M^n$$

≈ “FO theory of M relativised to q.r. n ”

Constructed inductively on n :

$$\phi_M^0 = \bigwedge_{\substack{\alpha \text{ atomic} \\ M \vDash \alpha}} \alpha \quad \wedge \quad \bigwedge_{\substack{\alpha \text{ atomic} \\ M \not\vDash \alpha}} \neg \alpha$$

$$\phi_M^n = \bigwedge_{\substack{u \in U^M \\ M_u = M[x:=u]}} \exists x \phi_{M_u}^{n-1} \quad \wedge \quad \bigvee_{\substack{u \in U^M \\ M_u = M[x:=u]}} \forall x \phi_{M_u}^{n-1}$$

Ehrenfeucht-Fraïssé games — completeness

$$\phi_M^n = \bigwedge_{\substack{u \in U^M \\ M_u = M[x:=u]}} \exists x \phi_{M_u}^{n-1} \wedge \bigvee_{\substack{u \in U^M \\ M_u = M[x:=u]}} \forall x \phi_{M_u}^{n-1}$$

Ehrenfeucht-Fraïssé games — completeness

We give a strategy
for Duplicator

under the hypothesis
 M, M' *n*-equivalent

$$\phi_M^n = \bigwedge_{u \in U^M} \exists x \phi_{M_u}^{n-1} \wedge \forall x \bigvee_{u \in U^M} \phi_{M_u}^{n-1}$$

$M_u = M[x := u]$

Ehrenfeucht-Fraïssé games — completeness

We give a strategy
for Duplicator

under the hypothesis
 M, M' *n*-equivalent

$$\phi_M^n = \bigwedge_{u \in U^M} \exists x \phi_{M_u}^{n-1} \wedge \forall x \bigvee_{u \in U^M} \phi_{M_u}^{n-1}$$

$M_u = M[x := u]$



EF game $G_{M,M'}$

Spoiler places pebble $u \in U^M$


Ehrenfeucht-Fraïssé games — completeness

We give a strategy
for Duplicator


under the hypothesis
 M, M' *n*-equivalent

$$\phi_M^n = \bigwedge_{u \in U^M} \exists x \phi_{M_u}^{n-1} \wedge \forall x \bigvee_{u \in U^M} \phi_{M_u}^{n-1}$$

$M_u = M[x := u]$



Adam picks left conjunct $\wedge \exists x \dots$
then picks conjunct $\exists x \phi_{M_u}^{n-1}$



Spoiler places pebble $u \in U^M$

Ehrenfeucht-Fraïssé games — completeness

We give a strategy
for Duplicator



under the hypothesis
 M, M' *n*-equivalent

$$\phi_M^n = \bigwedge_{u \in U^M} \exists x \phi_{M_u}^{n-1} \wedge \forall x \bigvee_{u \in U^M} \phi_{M_u}^{n-1}$$

$M_u = M[x := u]$

Facts:

- $M \models \phi_M^n$
and Eve has a *standard* winning strategy in $G_{M,\phi}$



Ehrenfeucht-Fraïssé games – completeness


We give a strategy
for Duplicator

under the hypothesis
 M, M' *n*-equivalent

$$\phi_M^n = \bigwedge_{u \in U^M} \exists x \phi_{M_u}^{n-1} \wedge \forall x \bigvee_{u \in U^M} \phi_{M_u}^{n-1}$$

Facts:

- $M \models \phi_M^n$
and Eve has a
standard winning
strategy in $G_{M,\phi}$



Spoiler places pebble $u \in U^M$

EF game $G_{M,M'}$

Ehrenfeucht-Fraïssé games — completeness

We give a strategy
for Duplicator


under the hypothesis
 M, M' *n*-equivalent

$$\phi_M^n = \bigwedge_{u \in U^M} \exists x \phi_{M_u}^{n-1} \wedge \forall x \bigvee_{u \in U^M} \phi_{M_u}^{n-1}$$

$M_u = M[x := u]$

Facts:

- $M \models \phi_M^n$
and Eve has a *standard* winning strategy in $G_{M,\phi}$



Ehrenfeucht-Fraïssé games — completeness

We give a strategy
for Duplicator

under the hypothesis
 M, M' *n*-equivalent

$$\phi_M^n = \bigwedge_{u \in U^M} \exists x \phi_{M_u}^{n-1} \wedge \forall x \bigvee_{u \in U^M} \phi_{M_u}^{n-1}$$

$M_u = M[x := u]$



Facts:

- $M \models \phi_M^n$

and Eve has a
standard winning
strategy in $G_{M,\phi}$

- $M' \models \phi_M^n$

so Eve has
also a winning
strategy in $G_{M',\phi}$



Ehrenfeucht-Fraïssé games — completeness

We give a strategy
for Duplicator

under the hypothesis
 M, M' *n*-equivalent


$$\phi_M^n = \bigwedge_{u \in U^M} \exists x \phi_{M_u}^{n-1} \wedge \forall x \bigvee_{u \in U^M} \phi_{M_u}^{n-1}$$

$M_u = M[x := u]$

Facts:

- $M \models \phi_M^n$
and Eve has a *standard* winning strategy in $G_{M,\phi}$

- $M' \models \phi_M^n$
so Eve has also a winning strategy in $G_{M',\phi}$



Ehrenfeucht-Fraïssé games — completeness

We give a strategy for Duplicator


under the hypothesis
 M, M' *n*-equivalent

$$\phi_M^n = \bigwedge_{u \in U^M} \exists x \phi_{M_u}^{n-1} \wedge \forall x \bigvee_{u \in U^M} \phi_{M_u}^{n-1}$$

$M_u = M[x := u]$

Facts:

- $M \models \phi_M^n$ and Eve has a *standard* winning strategy in $G_{M,\phi}$
- $M' \models \phi_M^n$ so Eve has also a winning strategy in $G_{M',\phi}$



Ehrenfeucht-Fraïssé games — completeness

We give a strategy
for Duplicator

under the hypothesis
 M, M' *n*-equivalent

$$\phi_M^n = \bigwedge_{u \in U^M} \exists x \phi_{M_u}^{n-1} \wedge \forall x \bigvee_{u \in U^M} \phi_{M_u}^{n-1}$$

$M_u = M[x := u]$

evaluation game $G_{M,\phi}$

evaluation game $G_{M',\phi}$

Facts:

- $M \models \phi_M^n$
and Eve has a *standard* winning strategy in $G_{M,\phi}$

- $M' \models \phi_M^n$
so Eve has also a winning strategy in $G_{M',\phi}$



EF game $G_{M,M'}$

Spoiler places pebble $v \in U^{M'}$

Ehrenfeucht-Fraïssé games — completeness

We give a strategy
for Duplicator

under the hypothesis
 M, M' *n*-equivalent


Facts:

- $M \models \phi_M^n$
and Eve has a *standard* winning strategy in $G_{M,\phi}$

- $M' \models \phi_M^n$
so Eve has also a winning strategy in $G_{M',\phi}$

$$\phi_M^n = \bigwedge_{u \in U^M} \exists x \phi_{M_u}^{n-1} \wedge \forall x \bigvee_{u \in U^M} \phi_{M_u}^{n-1}$$

$M_u = M[x:=u]$



Ehrenfeucht-Fraïssé games — completeness

We give a strategy
for Duplicator

under the hypothesis
 M, M' *n*-equivalent

Facts:

- $M \models \phi_M^n$
and Eve has a *standard* winning strategy in $G_{M,\phi}$

- $M' \models \phi_M^n$
so Eve has also a winning strategy in $G_{M',\phi}$

$$\phi_M^n = \bigwedge_{u \in U^M} \exists x \phi_{M_u}^{n-1} \wedge \forall x \bigvee_{u \in U^M} \phi_{M_u}^{n-1}$$

$M_u = M[x:=u]$

evaluation game $G_{M,\phi}$

evaluation game $G_{M',\phi}$



Adam picks right conjunct $\forall x \bigvee \dots$
then binds x to v



Eve picks *some* disjunct $\phi_{M_u}^{n-1}$
and continue the game

EF game $G_{M,M'}$



Spoiler places pebble $v \in U^{M'}$

Ehrenfeucht-Fraïssé games — completeness

We give a strategy
for Duplicator

under the hypothesis
 M, M' *n*-equivalent


Facts:

- $M \models \phi_M^n$
and Eve has a *standard* winning strategy in $G_{M,\phi}$

- $M' \models \phi_M^n$
so Eve has also a winning strategy in $G_{M',\phi}$

$$\phi_M^n = \bigwedge_{u \in U^M} \exists x \phi_{M_u}^{n-1} \wedge \forall x \bigvee_{u \in U^M} \phi_{M_u}^{n-1}$$

$M_u = M[x:=u]$



Ehrenfeucht-Fraïssé games — completeness

We give a strategy
for Duplicator

under the hypothesis
 M, M' *n*-equivalent

Facts:

- $M \models \phi_M^n$



and Eve has a
standard winning
strategy in $G_{M,\phi}$

- $M' \models \phi_M^n$

so Eve has
also a winning
strategy in $G_{M',\phi}$

$$\phi_M^n = \bigwedge_{u \in U^M} \exists x \phi_{M_u}^{n-1} \wedge \forall x \bigvee_{u \in U^M} \phi_{M_u}^{n-1}$$

$M_u = M[x:=u]$



Ehrenfeucht-Fraïssé games — completeness

We give a strategy for Duplicator

under the hypothesis
 M, M' *n*-equivalent


Facts:

- $M \models \phi_M^n$
 and Eve has a *standard* winning strategy in $G_{M,\phi}$

- $M' \models \phi_M^n$
 so Eve has also a winning strategy in $G_{M',\phi}$

$$\phi_M^n = \bigwedge_{u \in U^M} \exists x \phi_{M_u}^{n-1} \wedge \forall x \bigvee_{u \in U^M} \phi_{M_u}^{n-1}$$

$M_u = M[x:=u]$



Ehrenfeucht-Fraïssé games – a few more things

Theorem M, M' n -equivalent iff Duplicator survives n rounds in $G_{M,M'}$
 iff ϕ_M^n and $\phi_{M'}^n$ are logically equivalent

Ehrenfeucht-Fraïssé games — a few more things

Theorem M, M' n -equivalent iff Duplicator survives n rounds in $G_{M,M'}$

iff ϕ_M^n and $\phi_{M'}^n$ are logically equivalent

S₀,

1. ϕ_M^n can be used as a representant of the n -equivalence class of M
 2. For every ϕ' of q.r. n , $\phi' \in FO[M]$ iff ϕ' is a logical consequence of ϕ_M^n

Another use of Ehrenfeucht-Fraïssé games — 0/1 Law

Theorem (0/1 Law)

[Glebskii et al. '69, Fagin '76]

Every FO formula ϕ is
either almost surely true ($P_\infty[\phi] = 1$)
or almost surely false ($P_\infty[\phi] = 0$)

Another use of Ehrenfeucht-Fraïssé games — 0/1 Law

Theorem (0/1 Law)

[Glebskii et al. '69, Fagin '76]


Every FO formula ϕ is
either almost surely true ($P_\infty[\phi] = 1$)
or almost surely false ($P_\infty[\phi] = 0$)

Proof

Let n = quantifier rank of ϕ

$$\delta_n = \forall x_1, \dots, x_n \forall y_1, \dots, y_n \exists z \bigwedge_{i,j} x_i \neq y_j \wedge E(x_i, z) \wedge \neg E(y_j, z)$$

(extension formula)



Another use of Ehrenfeucht-Fraïssé games — 0/1 Law

Theorem (0/1 Law)

[Glebskii et al. '69, Fagin '76]


Every FO formula ϕ is
either almost surely true ($P_\infty[\phi] = 1$)
or almost surely false ($P_\infty[\phi] = 0$)

Proof

Let n = quantifier rank of ϕ

$$\delta_n = \forall x_1, \dots, x_n \ \forall y_1, \dots, y_n \ \exists z \ \bigwedge_{i,j} x_i \neq y_j \wedge E(x_i, z) \wedge \neg E(y_j, z)$$

(extension formula)



Fact 1: If $M \models \delta_n \wedge M' \models \delta_n$ then
Duplicator survives n rounds on $G_{M,M'}$

Another use of Ehrenfeucht-Fraïssé games — 0/1 Law

Theorem (0/1 Law)

[Glebskii et al. '69, Fagin '76]


Every FO formula ϕ is
either almost surely true ($P_\infty[\phi] = 1$)
or almost surely false ($P_\infty[\phi] = 0$)

Proof

Let n = quantifier rank of ϕ

$$\delta_n = \forall x_1, \dots, x_n \ \forall y_1, \dots, y_n \ \exists z \ \bigwedge_{i,j} x_i \neq y_j \wedge E(x_i, z) \wedge \neg E(y_j, z)$$

(extension formula)



Fact 1: If $M \models \delta_n \wedge M' \models \delta_n$ then
Duplicator survives n rounds on $G_{M,M'}$

Fact 2: $P_\infty[\delta_n] = 1$
(δ_n is almost surely true)

Another use of Ehrenfeucht-Fraïssé games — 0/1 Law

Theorem (0/1 Law)

[Glebskii et al. '69, Fagin '76]


Every FO formula ϕ is
either almost surely true ($P_\infty[\phi] = 1$)
or almost surely false ($P_\infty[\phi] = 0$)

Proof

Let n = quantifier rank of ϕ

$$\delta_n = \forall x_1, \dots, x_n \forall y_1, \dots, y_n \exists z \bigwedge_{i,j} x_i \neq y_j \wedge E(x_i, z) \wedge \neg E(y_j, z)$$

(extension formula)



Fact 1: If $M \models \delta_n \wedge M' \models \delta_n$ then
Duplicator survives n rounds on $G_{M,M'}$

Fact 2: $P_\infty[\delta_n] = 1$
(δ_n is almost surely true)

- a) There is $M \models \delta_n \wedge \phi \Rightarrow$ (by Fact 1) for every M' if $M' \models \delta_n$ then $M' \models \phi$
- Thus, $P_\infty[\delta_n] \leq P_\infty[\phi]$
- \Rightarrow (by Fact 2) $P_\infty[\delta_n] = 1$, hence $P_\infty[\phi] = 1$
- 2 cases
- 

Another use of Ehrenfeucht-Fraïssé games — 0/1 Law

Theorem (0/1 Law)

[Glebskii et al. '69, Fagin '76]


Every FO formula ϕ is
either almost surely true ($P_\infty[\phi] = 1$)
or almost surely false ($P_\infty[\phi] = 0$)

Proof

Let n = quantifier rank of ϕ

$$\delta_n = \forall x_1, \dots, x_n \forall y_1, \dots, y_n \exists z \bigwedge_{i,j} x_i \neq y_j \wedge E(x_i, z) \wedge \neg E(y_j, z)$$

(extension formula)



Fact 1: If $M \models \delta_n \wedge M' \models \delta_n$ then
Duplicator survives n rounds on $G_{M,M'}$

Fact 2: $P_\infty[\delta_n] = 1$
(δ_n is almost surely true)

- 2 cases
- a) There is $M \models \delta_n \wedge \phi \Rightarrow$ (by Fact 1) for every M' if $M' \models \delta_n$ then $M' \models \phi$
Thus, $P_\infty[\delta_n] \leq P_\infty[\phi]$
 \Rightarrow (by Fact 2) $P_\infty[\delta_n] = 1$, hence $P_\infty[\phi] = 1$
 - b) There is no $M \models \delta_n \wedge \phi \Rightarrow$ (by Fact 2) there is $M \models \delta_n$
 $\Rightarrow M \models \delta_n \wedge \neg\phi \Rightarrow$ (by case a) $P_\infty[\neg\phi] = 1$

Yes another use of games: synthesis (evaluation games in disguise!)



A. Church

Yes another use of games: synthesis (evaluation games in disguise!)

“Given a *requirement* which a *circuit* is to satisfy, we may suppose the requirement expressed in some suitable logistic system which is an extension of restricted recursive arithmetic. The *synthesis problem* is then to find recursion equivalences representing a circuit that satisfies the given requirement (or alternatively, to determine that there is no such circuit).”



A. Church

Yes another use of games: synthesis (evaluation games in disguise!)

Recall again the plain reachability problem encoded in QBF

k bits

0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1
0	0	0	0	1	1	1	1	1	1	1	1	0	0	0	0	0
0	0	1	1	1	1	0	0	0	1	1	1	1	0	0	0	0
0	1	1	0	0	1	1	0	0	1	1	0	0	1	1	1	0

$$\exists \bar{p}_1 \dots \exists \bar{p}_n \ \phi_{\text{path}}(\bar{p}_1, \dots, \bar{p}_n)$$

Yes another use of games: synthesis (evaluation games in disguise!)

Recall again the plain reachability problem encoded in QBF

0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1
0	0	0	0	1	1	1	1	1	1	1	1	0	0	0	0	0
0	0	1	1	1	1	0	0	0	1	1	1	1	0	0	0	0
0	1	1	0	0	1	1	0	0	1	1	0	0	1	1	1	0

$$\exists \bar{p}_1 \dots \exists \bar{p}_n \ \phi_{\text{path}}(\bar{p}_1, \dots, \bar{p}_n)$$

Now, suppose first 2 bits (\bar{q}_i) are controlled by **Environment**
last 2 bits (\bar{p}_i) are controlled by **Circuit**

Yes another use of games: synthesis (evaluation games in disguise!)

Recall again the plain reachability problem encoded in QBF

0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1
0	0	0	0	1	1	1	1	1	1	1	1	0	0	0	0	0
0	0	1	1	1	1	0	0	0	0	1	1	1	1	0	0	0
0	1	1	0	0	1	1	0	0	1	1	0	0	1	1	0	0

$$\exists \bar{p}_1 \dots \exists \bar{p}_n \ \phi_{\text{path}}(\bar{p}_1, \dots, \bar{p}_n)$$

Now, suppose first 2 bits (\bar{q}_i) are controlled by **Environment**
last 2 bits (\bar{p}_i) are controlled by **Circuit**

Question: can **Circuit** always reach goal, no matter how **Environment** behaves?
If so, can we synthesise a strategy for **Circuit**?

Yes another use of games: synthesis (evaluation games in disguise!)

Recall again the plain reachability problem encoded in QBF

0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1
0	0	0	0	1	1	1	1	1	1	1	1	0	0	0	0	0
0	0	1	1	1	1	0	0	0	0	1	1	1	1	0	0	0
0	1	1	0	0	1	1	0	0	1	1	0	0	1	1	0	0

$$\forall \bar{q}_1 \exists \bar{p}_1 \dots \forall \bar{q}_n \exists \bar{p}_n \ \phi_{\text{path}}(\bar{q}_1, \bar{p}_1, \dots, \bar{q}_n, \bar{p}_n)$$

Now, suppose first 2 bits (\bar{q}_i) are controlled by **Environment**
last 2 bits (\bar{p}_i) are controlled by **Circuit**

Question: can **Circuit** always reach goal, no matter how **Environment** behaves?
If so, can we synthesise a strategy for **Circuit**?

Things to remember



Things to remember

- EF games are a powerful tool (sound & complete) to study definability in FO
technique: 1) given property P and $n \in \mathbb{N}$
 2) find two models $M \in P, M' \notin P$ (which may depend on n !)
 3) show that Duplicator has strategy to survive n rounds in $G_{M,M'}$
- EF games can also be easily adapted to other logics and problems



What next?

More models: infinite words, infinite trees

More power: MSO = Monadic Second-order logic

More tools: automata

An appetiser — FO logic over words

Fix $\Sigma = \{A, B, C, \dots\}$ set of unary relational symbols

An appetiser — FO logic over words



Fix $\Sigma = \{A, B, C, \dots\}$ set of unary relational symbols

An appetiser — FO logic over words



Fix $\Sigma = \{A, B, C, \dots\}$ set of unary relational symbols

Consider models of the form $M = (\{0, 1, \dots, n\}, \leq, A^M, B^M, C^M, \dots)$



Sets partitioning $\{0, 1, \dots, n\}$

An appetiser — FO logic over words



Fix $\Sigma = \{A, B, C, \dots\}$ set of unary relational symbols

Consider models of the form $M = (\{0, 1, \dots, n\}, \leq, A^M, B^M, C^M, \dots)$



Sets partitioning $\{0, 1, \dots, n\}$

Every such M can be identified with a word $w_M \in \Sigma^*$

An appetiser — FO logic over words

Fix $\Sigma = \{A, B, C, \dots\}$ set of unary relational symbols

Consider models of the form $M = (\{0, 1, \dots, n\}, \leq, A^M, B^M, C^M, \dots)$



Sets partitioning $\{0, 1, \dots, n\}$

Every such M can be identified with a word $w_M \in \Sigma^*$

An appetiser — FO logic over words

Fix $\Sigma = \{A, B, C, \dots\}$ set of unary relational symbols

Consider models of the form $M = (\{0, 1, \dots, n\}, \leq, A^M, B^M, C^M, \dots)$



Sets partitioning $\{0, 1, \dots, n\}$

Every such M can be identified with a word $w_M \in \Sigma^*$

So,

1. FO formulas of signature $\{\leq, A, B, C, \dots\}$ can be evaluated on words over Σ
2. Every such formula ϕ defines a language $L_\phi = \{ w_M \in \Sigma^* \mid M \models \phi \}$

An appetiser — FO logic over words

So,

1. FO formulas of signature $\{\leq, A, B, C, \dots\}$ can be evaluated on words over Σ
2. Every such formula ϕ defines a language $L_\phi = \{ w_M \in \Sigma^* \mid M \models \phi \}$

An appetiser — FO logic over words

So,

1. FO formulas of signature $\{\leq, A, B, C, \dots\}$ can be evaluated on words over Σ
2. Every such formula ϕ defines a language $L_\phi = \{ w_M \in \Sigma^* \mid M \models \phi \}$

Examples

- $\phi = \exists x \forall y (y \leq x \rightarrow A(y)) \wedge (y > x \rightarrow B(y))$ defines $L_\phi = A^*B^*$

An appetiser — FO logic over words

So,

1. FO formulas of signature $\{\leq, A, B, C, \dots\}$ can be evaluated on words over Σ
2. Every such formula ϕ defines a language $L_\phi = \{ w_M \in \Sigma^* \mid M \models \phi \}$

Examples

- $\phi = \exists x \forall y (y \leq x \rightarrow A(y)) \wedge (y > x \rightarrow B(y))$ defines $L_\phi = A^*B^*$
- $\phi = \forall x (A(x) \rightarrow B(x+1)) \wedge (B(x) \rightarrow B(x-1))$ defines $L_\phi = (AB)^*$

An appetiser — FO logic over words

So,

1. FO formulas of signature $\{\leq, A, B, C, \dots\}$ can be evaluated on words over Σ
2. Every such formula ϕ defines a language $L_\phi = \{ w_M \in \Sigma^* \mid M \models \phi \}$

Examples

- $\phi = \exists x \forall y (y \leq x \rightarrow A(y)) \wedge (y > x \rightarrow B(y))$ defines $L_\phi = A^*B^*$
- $\phi = \forall x (A(x) \rightarrow B(x+1)) \wedge (B(x) \rightarrow B(x-1))$ defines $L_\phi = (AB)^*$
- Can you define in FO the language $L = A^* B A^*$? And $L = (AA)^*$?