

# First-Order logic (FO)

# First-Order logic (FO)



# First-Order logic (FO)





# FO = First-Order logic

Vocabulary	<u>Relational symbols:</u>	$\Sigma = \{R, S, T, \dots\}$	(aka <u>signature</u> )
	<u>Variables:</u>	$x, y, \dots, x_1, x_2, \dots$	
	<u>Quantifiers:</u>	$\exists, \forall$	
	Boolean connectives:	$\vee, \wedge, \neg, \rightarrow, \leftrightarrow$	

# FO = First-Order logic

Vocabulary	<u>Relational symbols:</u>	$\Sigma = \{R, S, T, \dots\}$	(aka <u>signature</u> )
	<u>Variables:</u>	$x, y, \dots, x_1, x_2, \dots$	
	<u>Quantifiers:</u>	$\exists, \forall$	
	Boolean connectives:	$\vee, \wedge, \neg, \rightarrow, \leftrightarrow$	
Syntax	$\phi : \quad R(x_1, \dots, x_k) \quad   \quad \dots \quad   \quad \phi \vee \phi \quad   \quad \phi \wedge \phi \quad   \quad \neg \phi \quad   \quad \phi \rightarrow \phi \quad   \quad \phi \leftrightarrow \phi$ $\quad \quad \exists x \phi \quad   \quad \forall x \phi \quad   \quad \dots$		

# FO = First-Order logic

Vocabulary	<u>Relational symbols:</u>	$\Sigma = \{R, S, T, \dots\}$	(aka <u>signature</u> )
	<u>Variables:</u>	$x, y, \dots, x_1, x_2, \dots$	
	<u>Quantifiers:</u>	$\exists, \forall$	
	Boolean connectives:	$\vee, \wedge, \neg, \rightarrow, \leftrightarrow$	

Syntax	$\phi :$	$R(x_1, \dots, x_k)$		$\dots$		$\phi \vee \phi$		$\phi \wedge \phi$		$\neg \phi$		$\phi \rightarrow \phi$		$\phi \leftrightarrow \phi$
		$\exists x \phi$		$\forall x \phi$		$\dots$								

Semantics	Now a model consists of a <u>universe</u>	$U^M$
		+ some <u>mappings</u>
		$R \mapsto R^M \subseteq U^M \times \dots \times U^M$ $x \mapsto x^M \in U^M$

# FO = First-Order logic

Vocabulary	<u>Relational symbols:</u>	$\Sigma = \{R, S, T, \dots\}$	(aka <u>signature</u> )
	<u>Variables:</u>	$x, y, \dots, x_1, x_2, \dots$	
	<u>Quantifiers:</u>	$\exists, \forall$	
	Boolean connectives:	$\vee, \wedge, \neg, \rightarrow, \leftrightarrow$	

Syntax	$\phi :$	$R(x_1, \dots, x_k)$		$\dots$		$\phi \vee \phi$		$\phi \wedge \phi$		$\neg \phi$		$\phi \rightarrow \phi$		$\phi \leftrightarrow \phi$
		$\exists x \phi$		$\forall x \phi$		$\dots$								

Semantics	Now a model consists of a <u>universe</u>		$U^M$
	+ some <u>mappings</u>		$R \mapsto R^M \subseteq U^M \times \dots \times U^M$
			$x \mapsto x^M \in U^M$

$M \models \phi_1 \vee \phi_2$       iff     $M \models \phi_1$     or     $M \models \phi_2$   
...

$M \models R(x_1, \dots, x_k)$     iff     $(x_1^M, \dots, x_k^M) \in R^M$

$M \models \exists x \phi$                 iff     $M[x:=u] \models \phi$  for *some*  $u \in U^M$

$M \models \forall x \phi$                 iff     $M[x:=u] \models \phi$  for *every*  $u \in U^M$

# Examples

Syntax  $\phi :$   $R(x_1, \dots, x_k)$  |  $\dots$  |  $\phi \vee \phi$  |  $\phi \wedge \phi$  |  $\neg \phi$  |  $\phi \rightarrow \phi$  |  $\phi \leftrightarrow \phi$   
 $\exists x \phi$  |  $\forall x \phi$  |  $\dots$

“All humans are mortal. Socrates is human. So Socrates is mortal.”

$$\phi(y) = ( (\forall x A(x) \rightarrow B(x)) \& A(y) ) \rightarrow B(y)$$



# Examples

Syntax  $\phi$  :  $R(x_1, \dots, x_k)$  |  $\dots$  |  $\phi \vee \phi$  |  $\phi \wedge \phi$  |  $\neg \phi$  |  $\phi \rightarrow \phi$  |  $\phi \leftrightarrow \phi$   
 $\exists x \phi$  |  $\forall x \phi$  |  $\dots$

“All humans are mortal. Socrates is human. So Socrates is mortal.”

$$\phi(y) = ( (\forall x A(x) \rightarrow B(x)) \ \& \ A(y) ) \rightarrow B(y)$$

M :  $U^M = \{\text{Socrates, Plato, Cyclop, Jupiter}\}$   
 $A^M = \{\text{Socrates, Plato}\}$   
 $B^M = \{\text{Socrates, Plato, Cyclop}\}$   
 $y^M = \text{Socrates}$

# Examples

Syntax  $\phi :$   $R(x_1, \dots, x_k)$  |  $\dots$  |  $\phi \vee \phi$  |  $\phi \wedge \phi$  |  $\neg \phi$  |  $\phi \rightarrow \phi$  |  $\phi \leftrightarrow \phi$   
 $\exists x \phi$  |  $\forall x \phi$  |  $\dots$

“There is a node in the graph that is isolated from all other nodes.”

$$\phi = \exists x \forall y \neg(x=y) \rightarrow \neg E(x,y)$$

# Examples

Syntax  $\phi :$   $R(x_1, \dots, x_k)$  |  $\dots$  |  $\phi \vee \phi$  |  $\phi \wedge \phi$  |  $\neg \phi$  |  $\phi \rightarrow \phi$  |  $\phi \leftrightarrow \phi$   
 $\exists x \phi$  |  $\forall x \phi$  |  $\dots$

“There is a node in the graph that is isolated from all other nodes.”

$$\phi = \exists x \forall y \neg(x=y) \rightarrow \neg E(x,y)$$

$M :$   $U^M = \{\text{nodes of a graph}\}$   
 $E^M = \{\text{edges of a graph}\}$

# Examples

Syntax

$\phi : R(x_1, \dots, x_k) \mid \dots \mid \phi \vee \phi \mid \phi \wedge \phi \mid \neg \phi \mid \phi \rightarrow \phi \mid \phi \leftrightarrow \phi$   
 $\exists x \phi \mid \forall x \phi \mid \dots$

# Examples

Syntax  $\phi$  :  $R(x_1, \dots, x_k)$  |  $\dots$  |  $\phi \vee \phi$  |  $\phi \wedge \phi$  |  $\neg \phi$  |  $\phi \rightarrow \phi$  |  $\phi \leftrightarrow \phi$   
 $\exists x \phi$  |  $\forall x \phi$  |  $\dots$

“There’s a man such that when he runs, everybody runs.”

$$\phi = \exists x R(x) \rightarrow \forall y R(y)$$

# Examples

Syntax  $\phi$  :  $R(x_1, \dots, x_k)$  |  $\dots$  |  $\phi \vee \phi$  |  $\phi \wedge \phi$  |  $\neg \phi$  |  $\phi \rightarrow \phi$  |  $\phi \leftrightarrow \phi$   
 $\exists x \phi$  |  $\forall x \phi$  |  $\dots$

“There’s a man such that when he runs, everybody runs.”

$$\phi = \exists x R(x) \rightarrow \forall y R(y)$$

$M$  :  $U^M = \{\text{Ben, Han, Leia, Luke}\}$   
 $R^M = \{\text{Ben, Han}\}$



# Examples

Syntax  $\phi$  :  $R(x_1, \dots, x_k)$  |  $\dots$  |  $\phi \vee \phi$  |  $\phi \wedge \phi$  |  $\neg \phi$  |  $\phi \rightarrow \phi$  |  $\phi \leftrightarrow \phi$   
 $\exists x \phi$  |  $\forall x \phi$  |  $\dots$

“There’s a man such that when he runs, everybody runs.”

$$\phi = \exists x R(x) \rightarrow \forall y R(y)$$

$M$  :  $U^M = \{\text{Ben, Han, Leia, Luke}\}$   
 $R^M = \{\text{Ben, Han}\}$

$M'$  :  $U^{M'} = \{\text{Ben, Han, Leia, Luke}\}$   
 $R^{M'} = \{\text{Ben, Han, Leia, Luke}\}$

# Examples

- “R is a function”

$$\phi = \forall x \exists y R(x,y) \wedge \forall z R(x,z) \rightarrow y=z$$

in this case, one can use the shorthand

$$\text{“}R(x)=\text{...” for } \exists y R(x,y) \wedge \forall z R(x,z) \rightarrow z=\text{...”}$$

# Examples

- “R is a function”

$$\phi = \forall x \exists y R(x,y) \wedge \forall z R(x,z) \rightarrow y=z$$

in this case, one can use the shorthand

$$\text{“}R(x)=\text{...” for } \exists y R(x,y) \wedge \forall z R(x,z) \rightarrow z=\text{...”}$$

- “+ is commutative”

$$\phi = \forall x \forall y x+y = y+x$$

note: + is a ternary relational symbol, so “ $x+y=z$ ” is shorthand for “ $+(x,y,z)$ ”

# Examples

- “R is a function”

$$\phi = \forall x \exists y R(x,y) \wedge \forall z R(x,z) \rightarrow y=z$$

in this case, one can use the shorthand

$$\text{“}R(x)=\text{...” for } \exists y R(x,y) \wedge \forall z R(x,z) \rightarrow z=...$$

- “+ is commutative”

$$\phi = \forall x \forall y x+y = y+x$$

note: + is a ternary relational symbol, so “ $x+y=z$ ” is shorthand for “ $+(x,y,z)$ ”

- “+ admits *zero* and *inverses*”

$$\phi = \exists x_0 \forall y x_0+y = y \wedge \forall y \exists z y+z = x_0$$

# Exercices

- “f is continuous”

$$\phi = \forall x \forall \varepsilon \exists \delta \forall y \quad ||x-y|| < \delta \rightarrow ||f(x) - f(y)|| < \varepsilon$$

- “f is uniformly continuous”

$$\phi = \forall \varepsilon \exists \delta \forall x \forall y \quad ||x-y|| < \delta \rightarrow ||f(x) - f(y)|| < \varepsilon$$

# Exercices

- “f is continuous”

$$\phi = \forall x \forall \varepsilon \exists \delta \forall y \ ||x-y|| < \delta \rightarrow ||f(x) - f(y)|| < \varepsilon$$

- “f is uniformly continuous”

$$\phi = \forall \varepsilon \exists \delta \forall x \forall y \ ||x-y|| < \delta \rightarrow ||f(x) - f(y)|| < \varepsilon$$

What is an appropriate signature for the above formulas?



# Exercices

- “f is continuous”  $\phi = \forall x \forall \varepsilon \exists \delta \forall y \ ||x-y|| < \delta \rightarrow ||f(x) - f(y)|| < \varepsilon$
- “f is uniformly continuous”  $\phi = \forall \varepsilon \exists \delta \forall x \forall y \ ||x-y|| < \delta \rightarrow ||f(x) - f(y)|| < \varepsilon$

What is an appropriate signature for the above formulas?

Are the formulas equivalent? Is one a consequence of another? Can you prove it?

(hint:  $\exists x \forall y \alpha \rightarrow \forall y \exists x \alpha$  assuming universe is non-empty)

# Exercices

Choose appropriate universes and signatures, and define these properties in FO:

1. “There are infinitely many Prime numbers”  $\phi = \dots$

2. “In the tree,  $z$  is the least common ancestor of  $x$  and  $y$ ”  $\phi(x,y,z) = \dots$

3. “Polynomial  $p$  evaluates to  $y$  on  $x$ ” (for fixed  $p$ )  $\phi_p(x,y) = \dots$

4. “The graph is strongly connected”  $\phi = \dots$

5. “In the infinite sequence of  $a$ ’s and  $b$ ’s, every  $a$  is followed by  $b$ ”  $\phi = \dots$

# Normal forms

Prenex [+CNF/DNF]

as for QBF, i.e.  $\phi = Q_{x_1} \dots Q_{x_n} \alpha(x_1, \dots, x_n)$

NNF (Negation Normal Form)

$\phi :$   $\exists x \phi \mid \forall x \phi \mid \phi \vee \phi \mid \phi \wedge \phi \mid \alpha$   
 $\alpha :$   $R(x_1, \dots, x_k) \mid \neg R(x_1, \dots, x_k)$

# Normal forms

Prenex [+CNF/DNF]

as for QBF, i.e.  $\phi = Q_{x_1} \dots Q_{x_n} \alpha(x_1, \dots, x_n)$

NNF (Negation Normal Form)

$\phi : \quad \exists x \phi \mid \forall x \phi \mid \phi \vee \phi \mid \phi \wedge \phi \mid \alpha$   
 $\alpha : \quad R(x_1, \dots, x_k) \mid \neg R(x_1, \dots, x_k)$

**Lemma**      Given  $\phi$  ( $\leftrightarrow$ -free), one can compute in polynomial time an *equivalent* formula  $\phi^*$  in NNF

**Proof**      As for propositional logic, push negations inside:

$$\neg \forall \phi \rightsquigarrow \exists \neg \phi$$

$$\neg \exists \phi \rightsquigarrow \forall \neg \phi$$

$$\neg(\phi_1 \wedge \phi_2) \rightsquigarrow \neg \phi_1 \vee \neg \phi_2$$

$$\neg(\phi_1 \vee \phi_2) \rightsquigarrow \neg \phi_1 \wedge \neg \phi_2$$

## Model-checking problem

input: formula  $\phi$  + *finite* model  $M$   
output: yes iff  $M \models \phi$

## Satisfiability problem

input: formula  $\phi$   
output: yes iff  $M \models \phi$  for *some*  $M$

(recall:  $\phi$  valid iff  $\neg\phi$  is not satisfiable  
 $\phi, \phi'$  equivalent iff  $\phi \leftrightarrow \phi'$  is valid)



## Model-checking problem

input: formula  $\phi$  + *finite* model  $M$   
output: yes iff  $M \models \phi$

💀 **UNDECIDABLE** 💀

## Satisfiability problem

input: formula  $\phi$   
output: yes iff  $M \models \phi$  for *some*  $M$

(recall:  $\phi$  valid iff  $\neg\phi$  is not satisfiable  
 $\phi, \phi'$  equivalent iff  $\phi \leftrightarrow \phi'$  is valid)



# Algorithms — model-checking

Model-check( $\varphi$ ,  $M$ )

```
if  $\varphi = R(x_1, \dots, x_k)$  then
  if  $(x_1^M, \dots, x_k^M) \in R^M$  then
    return true
  else
    return false
else if  $\varphi = \varphi_1 \vee \varphi_2$  then
  return Model-check( $\varphi_1$ ,  $M$ ) OR
    Model-check( $\varphi_2$ ,  $M$ )
else if ...
...
else if  $\varphi = \exists x \varphi'$  then
  for  $u \in U^M$  do
    if Model-check( $\varphi'$ ,  $M[x:=u]$ ) then
      return true
  return false
else if  $\varphi = \forall x \varphi'$  then
  for  $u \in U^M$  do
    if NOT Model-check( $\varphi'$ ,  $M[x:=u]$ ) then
      return false
  return true
```

# Algorithms — satisfiability

**Theorem** [Trakhtenbrot '50]


Satisfiability of FO is **undecidable**

# Algorithms — satisfiability

**Theorem** [Trakhtenbrot '50]

Satisfiability of FO is **undecidable**

**Proof** by reduction from Domino (aka Tiling) problem...




# Algorithms — satisfiability

Theorem [Trakhtenbrot '50]

Satisfiability of FO is **undecidable**

Proof by reduction from Domino (aka Tiling) problem...



Reduction from **P** to **P'**:

(think of “**P** easier than **P'**”)




Algorithm **A** that solves **P** by using  
an oracle that returns solutions to **P'**

e.g. many-one reduction: for all  $x$   $P(x)$  iff  $P'(A(x))$

# The (undecidable) Domino problem

## Domino


**Input:** 4-sided dominos:




# The (undecidable) Domino problem

## Domino

**Input:** 4-sided dominos:



**Output:** Is it possible to form a white-bordered rectangle? (of any size)







# The (undecidable) Domino problem

## Domino

**Input:** 4-sided dominos:



**Output:** Is it possible to form a white-bordered rectangle? (of any size)




**Rules:** sides must match,  
you can't rotate the dominos, but you can 'clone' them.

# The (undecidable) Domino problem

## Domino - Why is it undecidable?




It can encode *halting* computations of Turing machines:




# The (undecidable) Domino problem

## Domino - Why is it undecidable?

It can encode *halting* computations of Turing machines:




(head is elsewhere,  
symbol is not modified)




# The (undecidable) Domino problem

## Domino - Why is it undecidable?


It can encode *halting* computations of Turing machines:



(head is elsewhere,  
symbol is not modified)






(head is here, symbol is  
rewritten, head moves right)





# The (undecidable) Domino problem

## Domino - Why is it undecidable?


It can encode *halting* computations of Turing machines:




(head is elsewhere,  
symbol is not modified)



(head is here, symbol is  
rewritten, head moves right)




(head is here, symbol is  
rewritten, head moves left)




# The (undecidable) Domino problem

## Domino - Why is it undecidable?


It can encode *halting* computations of Turing machines:




(head is elsewhere,  
symbol is not modified)




(head is here, symbol is  
rewritten, head moves right)



(head is here, symbol is  
rewritten, head moves left)




(initial configuration)




# The (undecidable) Domino problem

## Domino - Why is it undecidable?


It can encode *halting* computations of Turing machines:




(head is elsewhere,  
symbol is not modified)




(head is here, symbol is  
rewritten, head moves right)



(head is here, symbol is  
rewritten, head moves left)




(initial configuration)



(halting configuration)

...




# Domino reduces to Sat-FO (domino has a solution iff $\phi$ satisfiable)

1. There is a grid:  $H(,)$  and  $V(,)$  are relations representing bijections such that...




# Domino reduces to Sat-FO (domino has a solution iff $\phi$ satisfiable)

1. There is a grid:  $H(,)$  and  $V(,)$  are relations representing bijections such that...




# Domino reduces to Sat-FO (domino has a solution iff $\phi$ satisfiable)

1. There is a grid:  $H(,)$  and  $V(,)$  are relations representing bijections such that...




# Domino reduces to Sat-FO (domino has a solution iff $\phi$ satisfiable)

1. There is a **grid**:  $H(,)$  and  $V(,)$  are relations representing bijections such that...



# Domino reduces to Sat-FO (domino has a solution iff $\phi$ satisfiable)

1. There is a grid:  $H(,)$  and  $V(,)$  are relations representing bijections such that...



2. Assign one domino to each node:  
a unary relation


$D_{\text{domino}}(x)$

for each domino



# Domino reduces to Sat-FO (domino has a solution iff $\phi$ satisfiable)

1. There is a grid:  $H(,)$  and  $V(,)$  are relations representing bijections such that...



2. Assign one domino to each node:  
a unary relation

$D_{\text{domino}}(x)$

for each domino




3. Match the sides  $\forall x \forall y$

if  $H(x,y)$ , then  $D_a(x) \wedge D_b(y)$

for some dominos  $a,b$  that 'match'  
horizontally (Idem vertically)

# Domino reduces to Sat-FO (domino has a solution iff $\phi$ satisfiable)

1. There is a grid:  $H(,)$  and  $V(,)$  are relations representing bijections such that...



2. Assign one domino to each node:  
a unary relation

$D_{\boxed{\begin{smallmatrix} \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \end{smallmatrix}}} (x)$

for each domino



3. Match the sides  $\forall x \forall y$

if  $H(x,y)$ , then  $D_a(x) \wedge D_b(y)$

for some dominos  $a,b$  that 'match'  
horizontally (Idem vertically)

4. Borders are white.

# Recap + quiz

- Model-checking for FO (does  $M \models \phi$ ?) is **PSPACE**-complete
- Satisfiability for FO (does  $M \models \phi$  for some  $M$ ?) is **undecidable**

# Recap + quiz

- Model-checking for FO (does  $M \models \phi$ ?) is **PSPACE**-complete
- Satisfiability for FO (does  $M \models \phi$  for some  $M$ ?) is **undecidable**

What about

- Validity for FO? (Problem def.: does  $M \models \phi$  for every  $M$ ?)
- Equivalence for FO? (Problem def.: is it true that, for every  $M$ ,  
 $M \models \phi$  iff  $M \models \phi'$  ?)



# Recap + quiz

- Model-checking for FO (does  $M \models \phi$ ?) is **PSPACE**-complete
- Satisfiability for FO (does  $M \models \phi$  for some  $M$ ?) is **undecidable**

What about

- Validity for FO? (Problem def.: does  $M \models \phi$  for every  $M$ ?)
- Equivalence for FO? (Problem def.: is it true that, for every  $M$ ,  
 $M \models \phi$  iff  $M \models \phi'$ ?)

Can you recall the complexity of analogous problems for

- Propositional logic?
- QBF?

Logical theory of a model  $M$  = set of all formulas  $\phi$  that hold on  $M$

Logical theory of a model  $M$  = set of all formulas  $\phi$  that hold on  $M$

$\text{FO}[U^M, R^M, S^M, \dots]$  denotes the FO theory of  $M = (U^M, R^M, S^M, \dots)$

Logical theory of a model  $M$  = set of all formulas  $\phi$  that hold on  $M$

$\text{FO}[U^M, R^M, S^M, \dots]$  denotes the FO theory of  $M = (U^M, R^M, S^M, \dots)$

## Example

$\text{FO}[\mathbb{N}, <] = \{ \exists x (x=x), \forall x \exists y x < y, \exists y \forall x \neg(x < y), \forall x \forall y x=y \vee x < y \vee y < x, \dots \}$

Logical theory of a model  $M$  = set of all formulas  $\phi$  that hold on  $M$

$\text{FO}[U^M, R^M, S^M, \dots]$  denotes the FO theory of  $M = (U^M, R^M, S^M, \dots)$

## Example

$\text{FO}[\mathbb{N}, <] = \{ \exists x (x=x), \forall x \exists y x < y, \exists y \forall x \neg(x < y), \forall x \forall y x=y \vee x < y \vee y < x, \dots \}$

(notation abuse: relation = is often present, but not explicitly listed  
any symbol  $R$  is often identified with its relation  $R^M$ )

$\text{FO}[\mathbb{N}, +, \cdot]$  = Peano arithmetic

$\text{FO}[\mathbb{R}, +, \cdot]$  = Arithmetic theory of real numbers

$\text{FO}[\mathbb{Z}, +]$  = Presburger arithmetic

$\text{FO}[\mathbb{N}^2, \leq_1, \leq_2]$  = First-order theory of the unlabelled grid

$\text{FO}[\{0,1\}, =]$   $\approx$  {Valid QBFs}

$\text{FO}[V_R, E_R]$  = First-order theory of “random” graph

$\text{FO}[C_M, T_M]$  = First-order theory of the transition graph of a Turing machine  $M$

$\text{FO}[\mathbb{N}, +, \cdot]$  = Peano arithmetic

$\text{FO}[\mathbb{R}, +, \cdot]$  = Arithmetic theory of real numbers

$\text{FO}[\mathbb{Z}, +]$  = Presburger arithmetic

$\text{FO}[\mathbb{N}^2, \leq_1, \leq_2]$  = First-order theory of the unlabelled grid

$\text{FO}[\{0,1\}, =]$   $\approx$  {Valid QBFs}

$\text{FO}[V_R, E_R]$  = First-order theory of “random” graph

$\text{FO}[C_M, T_M]$  = First-order theory of the transition graph of a Turing machine M

How do I  
compare  
them?



# Logical reductions

Reduction from  $P$  to  $P'$ :

Algorithm  $A$  that solves  $P$  by using  
an oracle that returns solutions to  $P'$

e.g.      for all  $x$      $P(x)$  iff  $P'(A(x))$



# Logical reductions

Reduction from  $P$  to  $P'$ :

Algorithm  $A$  that solves  $P$  by using  
an oracle that returns solutions to  $P'$

e.g.      for all  $x$      $P(x)$  iff  $P'(A(x))$

Take  $P = \text{FO}[M] = \{\phi \mid M \models \phi\}$   
 $P' = \text{FO}[M'] = \{\phi' \mid M' \models \phi'\}$

for all  $\phi$      $M \models \phi$  iff  $M' \models A(\phi)$

described by a logical  
interpretation of  $M$  into  $M'$



# Logical reductions

Reduction from  $P$  to  $P'$ :

Algorithm  $A$  that solves  $P$  by using an oracle that returns solutions to  $P'$

e.g.      for all  $x$      $P(x)$  iff  $P'(A(x))$

Take  $P = \text{FO}[M] = \{\phi \mid M \models \phi\}$   
 $P' = \text{FO}[M'] = \{\phi' \mid M' \models \phi'\}$

for all  $\phi$      $M \models \phi$  iff  $M' \models A(\phi)$

described by a logical  
interpretation of  $M$  into  $M'$



FO interpretation of  $M$  into  $M'$ :      a mapping  $\alpha : R \mapsto \alpha_R$  such that

$M[\bar{u}] \models R(\bar{x})$  iff  $M'[\bar{x} := \bar{u}] \models \alpha_R(\bar{x})$

# Logical reductions

FO interpretation of  $M$  into  $M'$ : a mapping  $\alpha: R \mapsto \alpha_R$  such that

$$M[\bar{u}] \models R(\bar{x}) \text{ iff } M'[\bar{x} := \bar{u}] \models \alpha_R(\bar{x})$$

# Logical reductions

FO interpretation of  $M$  into  $M'$ : a mapping  $\alpha: R \mapsto \alpha_R$  such that

$$M[\bar{u}] \models R(\bar{x}) \text{ iff } M'[\bar{x} := \bar{u}] \models \alpha_R(\bar{x})$$

## Examples

- interpretation of  $M = (\mathbb{N}, \leq)$  into  $M' = (\mathbb{N}, +)$

$$\alpha_{\leq}(x, y) = \exists z \ y = x + z$$

# Logical reductions

F0 interpretation of  $M$  into  $M'$ : a mapping  $\alpha: R \mapsto \alpha_R$  such that

$$M[\bar{u}] \models R(\bar{x}) \text{ iff } M'[\bar{x} := \bar{u}] \models \alpha_R(\bar{x})$$

## Examples

- interpretation of  $M = (\mathbb{N}, \leq)$  into  $M' = (\mathbb{N}, +)$

$$\alpha_{\leq}(x, y) = \exists z \ y = x + z$$

- interpretation of  $M = (\{0,1\}^*, \leq_{\text{inorder}})$  into  $M' = (\{0,1\}^*, 0, 1, \cdot)$   
 $\approx (\mathbb{Q}, \leq)$

$$\alpha_{\leq_{\text{inorder}}}(x, y) = \exists x', y', z \ (x = z \cdot 0 \cdot x' \wedge y = z \cdot 1 \cdot y') \vee \\ (x = y \cdot 0 \cdot x') \vee (y = x \cdot 1 \cdot x')$$

# Logical reductions

In fact, an FO interpretation of  $M$  into  $M'$  is more complex (and powerful)

- definitions of relations:  $\alpha_R(\bar{x})$  such that  $R^M = \{ \bar{u} \mid M'[\bar{x} := \bar{u}] \models \alpha_R(\bar{x}) \}$

(e.g. to interpret  $(\mathbb{N}, \leq)$  into  $(\mathbb{N}, +)$ )

# Logical reductions

In fact, an FO interpretation of  $M$  into  $M'$  is more complex (and powerful)

- definitions of relations:  $\alpha_R(\bar{x})$  such that  $R^M = \{ \bar{u} \mid M'[\bar{x} := \bar{u}] \models \alpha_R(\bar{x}) \}$

(e.g. to interpret  $(\mathbb{N}, \leq)$  into  $(\mathbb{N}, +)$ )

- definition of universe:  $\alpha_U(x)$  such that  $U^M = \{ u \mid M'[x := u] \models \alpha_U(x) \}$

(e.g. to interpret  $(\mathbb{N}, \leq)$  into  $(\mathbb{Z}, \leq, 0)$ )

# Logical reductions

In fact, an FO interpretation of  $M$  into  $M'$  is more complex (and powerful)

- definitions of relations:  $\alpha_R(\bar{x})$  such that  $R^M = \{ \bar{u} \mid M'[\bar{x} := \bar{u}] \models \alpha_R(\bar{x}) \}$

(e.g. to interpret  $(\mathbb{N}, \leq)$  into  $(\mathbb{N}, +)$ )

- definition of universe:  $\alpha_U(x)$  such that  $U^M = \{ u \mid M'[x := u] \models \alpha_U(x) \}$

(e.g. to interpret  $(\mathbb{N}, \leq)$  into  $(\mathbb{Z}, \leq, 0)$ )

- k-dimensionality: elements of  $U^M$  can be *k-tuples* of elements of  $U^{M'}$

(e.g. to interpret  $(\mathbb{C}, +, \cdot)$  into  $(\mathbb{R}, +, \cdot)$ )



# Logical reductions

In fact, an FO interpretation of  $M$  into  $M'$  is more complex (and powerful)

- definitions of relations:  $\alpha_R(\bar{x})$  such that  $R^M = \{ \bar{u} \mid M'[\bar{x} := \bar{u}] \models \alpha_R(\bar{x}) \}$

(e.g. to interpret  $(\mathbb{N}, \leq)$  into  $(\mathbb{N}, +)$ )

- definition of universe:  $\alpha_U(x)$  such that  $U^M = \{ u \mid M'[x := u] \models \alpha_U(x) \}$

(e.g. to interpret  $(\mathbb{N}, \leq)$  into  $(\mathbb{Z}, \leq, 0)$ )

- k-dimensionality: elements of  $U^M$  can be *k-tuples* of elements of  $U^{M'}$

(e.g. to interpret  $(\mathbb{C}, +, \cdot)$  into  $(\mathbb{R}, +, \cdot)$ )

- quotient:  $\alpha_=(\bar{x}, \bar{y})$  such that  $M[\dots] \models (\bar{x} = \bar{y})$  iff  $M'[\dots] \models \alpha_=(\bar{x}, \bar{y})$

(e.g. to interpret  $(\mathbb{Q}, +, \cdot)$  into  $(\mathbb{Z}, +, \cdot)$ )

# Logical reductions

Given  $M'$  and an FO interpretation  $\alpha = (\alpha_U, \alpha_-, \alpha_R, \alpha_S, \dots)$   
the interpreted model is  $\alpha(M') = (U^M, R^M, S^M, \dots)$  where

- $U^M = \{ [\bar{u}]_{\approx} \mid M'[\bar{x} := \bar{u}] \models \alpha_U(\bar{x}) \}$
- $\bar{u} \approx \bar{v}$  iff  $M'[\bar{x} := \bar{u}, \bar{y} := \bar{v}] \models \alpha_-(\bar{x}, \bar{y})$
- $R^M = \{ ([\bar{u}_1]_{\approx}, \dots, [\bar{u}_k]_{\approx}) \mid M'[\bar{x}_1 := \bar{u}_1, \dots, \bar{x}_k := \bar{u}_k] \models \alpha_R(\bar{x}_1, \dots, \bar{x}_k) \}$   
(needs to be well-defined, namely,  $\approx$  needs to be a congruence w.r.t. every relation R)
- ...

# Logical reductions

Given  $M'$  and an FO interpretation  $\alpha = (\alpha_U, \alpha_-, \alpha_R, \alpha_S, \dots)$   
the interpreted model is  $\alpha(M') = (U^M, R^M, S^M, \dots)$  where

- $U^M = \{ [\bar{u}]_{\approx} \mid M'[\bar{x} := \bar{u}] \models \alpha_U(\bar{x}) \}$
- $\bar{u} \approx \bar{v}$  iff  $M'[\bar{x} := \bar{u}, \bar{y} := \bar{v}] \models \alpha_-(\bar{x}, \bar{y})$
- $R^M = \{ ([\bar{u}_1]_{\approx}, \dots, [\bar{u}_k]_{\approx}) \mid M'[\bar{x}_1 := \bar{u}_1, \dots, \bar{x}_k := \bar{u}_k] \models \alpha_R(\bar{x}_1, \dots, \bar{x}_k) \}$   
(needs to be well-defined, namely,  $\approx$  needs to be a congruence w.r.t. every relation R)
- ...

**Theorem** If  $\alpha = (\alpha_U, \alpha_-, \alpha_R, \alpha_S, \dots)$  is an FO interpretation of  $M$  into  $M'$   
then  $\text{FO}[M]$  *reduces to*  $\text{FO}[M']$ , namely, there is an algorithm  $A_\alpha$

for all  $\phi$       $M \models \phi$  iff  $M' \models A_\alpha(\phi)$

# Some fancy FO theories

$\text{FO}[\mathbb{N}, +, \cdot]$  = Peano arithmetic

$\text{FO}[\mathbb{R}, +, \cdot]$  = Arithmetic theory of real numbers

$\text{FO}[\mathbb{Z}, +]$  = Presburger arithmetic

$\text{FO}[\mathbb{N}^2, \leq_1, \leq_2]$  = First-order theory of the unlabelled grid

$\text{FO}[\{0,1\}, =]$   $\approx$  {Valid QBFs}

$\text{FO}[V_R, E_R]$  = First-order theory of “random” graph

$\text{FO}[C_M, T_M]$  = First-order theory of the transition graph of a Turing machine  $M$

# $\text{FO}[\mathbb{N}, +, \cdot]$ — Peano arithmetic

## Theorem

Peano arithmetic is undecidable

(one cannot check whether  $(\mathbb{N}, +, \cdot) \models \phi$  for a given  $\phi$ )

# $\text{FO}[\mathbb{N}, +, \cdot]$ — Peano arithmetic

## Theorem

Peano arithmetic is undecidable

(one cannot check whether  $(\mathbb{N}, +, \cdot) \models \phi$  for a given  $\phi$ )

**Proof** by reduction from undecidable Hilbert's 10th problem... [Matiyasevic '70]

### Hilbert's 10th

Given a polynomial  $p(x, y, z, \dots)$

tell whether  $p(x, y, z, \dots) = 0$  for *some integers*  $x, y, z$

# $\text{FO}[\mathbb{N}, +, \cdot]$ — Peano arithmetic

## Theorem

Peano arithmetic is undecidable

(one cannot check whether  $(\mathbb{N}, +, \cdot) \models \phi$  for a given  $\phi$ )

**Proof** by reduction from undecidable Hilbert's 10th problem... [Matiyasevic '70]

### Hilbert's 10th

Given a polynomial  $p(x, y, z, \dots)$   
tell whether  $p(x, y, z, \dots) = 0$  for *some integers*  $x, y, z$

1. Given polynomial  $p(x, y, z, \dots)$ , inductively construct  $\phi_p(x, y, z, \dots, t)$  such that
$$(\mathbb{Z}, +, \cdot, x, y, z, \dots, t) \models \phi_p \text{ iff } p(x, y, z) = t$$
2. Interpret  $(\mathbb{Z}, +, \cdot, 0)$  into  $(\mathbb{N}, +, \cdot)$

# Some fancy FO theories

$\text{FO}[\mathbb{N}, +, \cdot]$  = Peano arithmetic

 **UNDECIDABLE**   
(reduction from H's 10th)

$\text{FO}[\mathbb{R}, +, \cdot]$  = Arithmetic theory of real numbers

$\text{FO}[\mathbb{Z}, +]$  = Presburger arithmetic

$\text{FO}[\mathbb{N}^2, \leq_1, \leq_2]$  = First-order theory of the unlabelled grid

$\text{FO}[\{0,1\}, =]$   $\approx$  {Valid QBFs}

$\text{FO}[V_R, E_R]$  = First-order theory of “random” graph

$\text{FO}[C_M, T_M]$  = First-order theory of the transition graph of a Turing machine  $M$



# $\text{FO}[\mathbb{R}, +, \cdot]$ — Arithmetic theory of real numbers

## Theorem

[Tarski '51]

Every FO formula  $\phi$  over  $(\mathbb{R}, +, \cdot)$  can be effectively transformed into an equivalent quantifier-free formula  $\phi^*$

# $\text{FO}[\mathbb{R}, +, \cdot]$ — Arithmetic theory of real numbers

**Theorem**  
[Tarski '51]

Every FO formula  $\phi$  over  $(\mathbb{R}, +, \cdot)$  can be effectively transformed into an equivalent quantifier-free formula  $\phi^*$

**Corollary**

Given  $\phi$ , one can decide whether  $(\mathbb{R}, +, \cdot) \models \phi$

# $\text{FO}[\mathbb{R}, +, \cdot]$ — Arithmetic theory of real numbers

**Theorem** [Tarski '51] Every FO formula  $\phi$  over  $(\mathbb{R}, +, \cdot)$  can be effectively transformed into an equivalent quantifier-free formula  $\phi^*$

**Corollary** Given  $\phi$ , one can decide whether  $(\mathbb{R}, +, \cdot) \models \phi$

Algebraic geometry

Continuous & discrete dynamical systems


Programs verification

Computer graphics

Robotics

Coding theory & Cryptography

Grammars & Transducers



# Some fancy FO theories

$\text{FO}[\mathbb{N}, +, \cdot]$  = Peano arithmetic

 **UNDECIDABLE**   
(reduction from H's 10th)

$\text{FO}[\mathbb{R}, +, \cdot]$  = Arithmetic theory of real numbers

 **DECIDABLE**   
(quantifier elimination)

$\text{FO}[\mathbb{Z}, +]$  = Presburger arithmetic

$\text{FO}[\mathbb{N}^2, \leq_1, \leq_2]$  = First-order theory of the unlabelled grid

$\text{FO}[\{0,1\}, =]$   $\approx$  {Valid QBFs}

$\text{FO}[V_R, E_R]$  = First-order theory of “random” graph

$\text{FO}[C_M, T_M]$  = First-order theory of the transition graph of a Turing machine  $M$

# FO[ $\mathbb{Z}, +$ ] — Presburger arithmetic

**Theorem** [Presburger '29] Every FO formula  $\phi$  over  $(\mathbb{Z}, +, 0, 1, \leq, |)$  can be effectively transformed into an equivalent quantifier-free formula  $\phi^*$

**Corollary** Given  $\phi$  over  $(\mathbb{Z}, +)$ , one can decide whether  $(\mathbb{Z}, +) \models \phi$

# FO[ $\mathbb{Z}, +$ ] — Presburger arithmetic

**Theorem** [Presburger '29] Every FO formula  $\phi$  over  $(\mathbb{Z}, +, 0, 1, \leq, |)$  can be effectively transformed into an equivalent quantifier-free formula  $\phi^*$

**Corollary** Given  $\phi$  over  $(\mathbb{Z}, +)$ , one can decide whether  $(\mathbb{Z}, +) \models \phi$

## Proof idea

Show how to remove an innermost quantifier  $Qz$  from  $\phi = \dots Qz \alpha(\dots, z)$

# FO[ $\mathbb{Z}, +$ ] — Presburger arithmetic

**Theorem** [Presburger '29] Every FO formula  $\phi$  over  $(\mathbb{Z}, +, 0, 1, \leq, |)$  can be effectively transformed into an equivalent quantifier-free formula  $\phi^*$

**Corollary** Given  $\phi$  over  $(\mathbb{Z}, +)$ , one can decide whether  $(\mathbb{Z}, +) \models \phi$

## Proof idea

Show how to remove an innermost quantifier  $Qz$  from  $\phi = \dots Qz \alpha(\dots, z)$

Assume:

- $Qz = \exists z$  (if not, treat  $\forall z$  as  $\neg \exists z \neg$ )
- $\alpha$  is  $\vee$ -free (if not, commute  $\exists$  and  $\vee$ )

# FO[ $\mathbb{Z}, +$ ] — Presburger arithmetic

**Theorem** [Presburger '29] Every FO formula  $\phi$  over  $(\mathbb{Z}, +, 0, 1, \leq, |)$  can be effectively transformed into an equivalent quantifier-free formula  $\phi^*$

**Corollary** Given  $\phi$  over  $(\mathbb{Z}, +)$ , one can decide whether  $(\mathbb{Z}, +) \models \phi$

## Proof idea

Show how to remove an innermost quantifier  $Qz$  from  $\phi = \dots Qz \alpha(\dots, z)$

Assume:

- $Qz = \exists z$  (if not, treat  $\forall z$  as  $\neg \exists z \neg$ )
- $\alpha$  is  $\vee$ -free (if not, commute  $\exists$  and  $\vee$ )

**Example**  $\exists z \alpha(x, y, z) = \exists z (2x + 4y - 3z \leq 7) \wedge (3x - y + 2z \leq -4)$



# FO[ $\mathbb{Z}, +$ ] — Presburger arithmetic

**Theorem** [Presburger '29] Every FO formula  $\phi$  over  $(\mathbb{Z}, +, 0, 1, \leq, |)$  can be effectively transformed into an equivalent quantifier-free formula  $\phi^*$

## Proof idea

Show how to remove an innermost quantifier  $Qz$  from  $\phi = \dots Qz \alpha(\dots, z)$

Assume:

- $Qz = \exists z$  (if not, treat  $\forall z$  as  $\neg \exists z \neg$ )
- $\alpha$  is  $\vee$ -free (if not, commute  $\exists$  and  $\vee$ )

**Example**  $\exists z \alpha(x, y, z) = \exists z (2x + 4y - 3z \leq 7) \wedge (3x - y + 2z \leq -4)$

# FO[ $\mathbb{Z}, +$ ] — Presburger arithmetic

**Theorem** [Presburger '29] Every FO formula  $\phi$  over  $(\mathbb{Z}, +, 0, 1, \leq, |)$  can be effectively transformed into an equivalent quantifier-free formula  $\phi^*$

## Proof idea

Show how to remove an innermost quantifier  $Qz$  from  $\phi = \dots Qz \alpha(\dots, z)$

Assume:

- $Qz = \exists z$  (if not, treat  $\forall z$  as  $\neg \exists z \neg$ )
- $\alpha$  is  $\vee$ -free (if not, commute  $\exists$  and  $\vee$ )

**Example**  $\exists z \alpha(x, y, z) = \exists z (2x + 4y - 3z \leq 7) \wedge (3x - y + 2z \leq -4)$   
 $\exists z (2x + 4y - 3z \leq 7) \wedge (3x - y + 2z \leq -4)$

# FO[ $\mathbb{Z}, +$ ] — Presburger arithmetic

**Theorem** [Presburger '29] Every FO formula  $\phi$  over  $(\mathbb{Z}, +, 0, 1, \leq, |)$  can be effectively transformed into an equivalent quantifier-free formula  $\phi^*$

## Proof idea

Show how to remove an innermost quantifier  $Qz$  from  $\phi = \dots Qz \alpha(\dots, z)$

Assume:

- $Qz = \exists z$  (if not, treat  $\forall z$  as  $\neg \exists z \neg$ )
- $\alpha$  is  $\vee$ -free (if not, commute  $\exists$  and  $\vee$ )

**Example**  $\exists z \alpha(x, y, z) = \exists z (2x + 4y - 3z \leq 7) \wedge (3x - y + 2z \leq -4)$   
 $\exists z (2x + 4y - 7 \leq 3z) \wedge (2z \leq -3x + y - 4)$

# FO[ $\mathbb{Z}, +$ ] — Presburger arithmetic

**Theorem** [Presburger '29] Every FO formula  $\phi$  over  $(\mathbb{Z}, +, 0, 1, \leq, |)$  can be effectively transformed into an equivalent quantifier-free formula  $\phi^*$

## Proof idea

Show how to remove an innermost quantifier  $Qz$  from  $\phi = \dots Qz \alpha(\dots, z)$

Assume:

- $Qz = \exists z$  (if not, treat  $\forall z$  as  $\neg \exists z \neg$ )
- $\alpha$  is  $\vee$ -free (if not, commute  $\exists$  and  $\vee$ )

## Example

$$\begin{aligned}\exists z \alpha(x, y, z) &= \exists z (2x + 4y - 3z \leq 7) \wedge (3x - y + 2z \leq -4) \\ &= \exists z (2x + 4y - 7 \leq 3z) \wedge (2z \leq -3x + y - 4) \\ &= \exists z 2 \cdot (2x + 4y - 7 \leq 3z) \wedge (2z \leq -3x + y - 4) \cdot 3\end{aligned}$$

# FO[ $\mathbb{Z}, +$ ] — Presburger arithmetic

**Theorem** [Presburger '29] Every FO formula  $\phi$  over  $(\mathbb{Z}, +, 0, 1, \leq, |)$  can be effectively transformed into an equivalent quantifier-free formula  $\phi^*$

## Proof idea

Show how to remove an innermost quantifier  $Qz$  from  $\phi = \dots Qz \alpha(\dots, z)$

Assume:

- $Qz = \exists z$  (if not, treat  $\forall z$  as  $\neg \exists z \neg$ )
- $\alpha$  is  $\vee$ -free (if not, commute  $\exists$  and  $\vee$ )

**Example**

$$\begin{aligned}\exists z \alpha(x, y, z) &= \exists z (2x + 4y - 3z \leq 7) \wedge (3x - y + 2z \leq -4) \\ &= \exists z (2x + 4y - 7 \leq 3z) \wedge (2z \leq -3x + y - 4) \\ &= \exists z (4x + 8y - 14 \leq 6z) \wedge (6z \leq -9x + 3y - 12)\end{aligned}$$

# FO[ $\mathbb{Z}, +$ ] — Presburger arithmetic

**Theorem** [Presburger '29] Every FO formula  $\phi$  over  $(\mathbb{Z}, +, 0, 1, \leq, |)$  can be effectively transformed into an equivalent quantifier-free formula  $\phi^*$

## Proof idea

Show how to remove an innermost quantifier  $Qz$  from  $\phi = \dots Qz \alpha(\dots, z)$

Assume:

- $Qz = \exists z$  (if not, treat  $\forall z$  as  $\neg \exists z \neg$ )
- $\alpha$  is  $\vee$ -free (if not, commute  $\exists$  and  $\vee$ )

## Example

temporarily assume formulas  
are over the reals or the rationals...

$$\exists z \alpha(x, y, z) = \exists z (2x + 4y - 3z \leq 7) \wedge (3x - y + 2z \leq -4)$$

$$\exists z (2x + 4y - 7 \leq 3z) \wedge (2z \leq -3x + y - 4)$$

$$\exists z (4x + 8y - 14 \leq 6z) \wedge (6z \leq -9x + 3y - 12)$$

# FO[ $\mathbb{Z}, +$ ] — Presburger arithmetic

**Theorem** [Presburger '29] Every FO formula  $\phi$  over  $(\mathbb{Z}, +, 0, 1, \leq, |)$  can be effectively transformed into an equivalent quantifier-free formula  $\phi^*$

## Proof idea

Show how to remove an innermost quantifier  $Qz$  from  $\phi = \dots Qz \alpha(\dots, z)$

Assume:

- $Qz = \exists z$  (if not, treat  $\forall z$  as  $\neg \exists z \neg$ )
- $\alpha$  is  $\vee$ -free (if not, commute  $\exists$  and  $\vee$ )

## Example

temporarily assume formulas  
are over the reals or the rationals...

$$\exists z \alpha(x, y, z) = \exists z (2x + 4y - 3z \leq 7) \wedge (3x - y + 2z \leq -4)$$

$$\exists z (2x + 4y - 7 \leq 3z) \wedge (2z \leq -3x + y - 4)$$

$$\exists z (4x + 8y - 14 \leq 6z) \wedge (6z \leq -9x + 3y - 12)$$

$$\exists z (4x + 8y - 14 \leq 6z) \wedge (6z \leq -9x + 3y - 12)$$

# FO[ $\mathbb{Z}, +$ ] — Presburger arithmetic

**Theorem** [Presburger '29] Every FO formula  $\phi$  over  $(\mathbb{Z}, +, 0, 1, \leq, |)$  can be effectively transformed into an equivalent quantifier-free formula  $\phi^*$

## Proof idea

Show how to remove an innermost quantifier  $Qz$  from  $\phi = \dots Qz \alpha(\dots, z)$

Assume:

- $Qz = \exists z$  (if not, treat  $\forall z$  as  $\neg \exists z \neg$ )
- $\alpha$  is  $\vee$ -free (if not, commute  $\exists$  and  $\vee$ )

## Example

temporarily assume formulas  
are over the reals or the rationals...

$$\exists z \alpha(x, y, z) = \exists z (2x + 4y - 3z \leq 7) \wedge (3x - y + 2z \leq -4)$$

$$\exists z (2x + 4y - 7 \leq 3z) \wedge (2z \leq -3x + y - 4)$$

$$\exists z (4x + 8y - 14 \leq 6z) \wedge (6z \leq -9x + 3y - 12)$$

$$4x + 8y - 14 \leq -9x + 3y - 12$$



# FO[ $\mathbb{Z}, +$ ] — Presburger arithmetic

**Theorem** [Presburger '29] Every FO formula  $\phi$  over  $(\mathbb{Z}, +, 0, 1, \leq, |)$  can be effectively transformed into an equivalent quantifier-free formula  $\phi^*$

## Proof idea

Show how to remove an innermost quantifier  $Qz$  from  $\phi = \dots Qz \alpha(\dots, z)$

Assume:

- $Qz = \exists z$  (if not, treat  $\forall z$  as  $\neg \exists z \neg$ )
- $\alpha$  is  $\vee$ -free (if not, commute  $\exists$  and  $\vee$ )

**Example**  $\exists z \alpha(x, y, z) = \exists z (2x + 4y - 3z \leq 7) \wedge (3x - y + 2z \leq -4)$

temporarily assume formulas  
are over the reals or the rationals...

$$\exists z (2x + 4y - 7 \leq 3z) \wedge (2z \leq -3x + y - 4)$$
$$\exists z (4x + 8y - 14 \leq 6z) \wedge (6z \leq -9x + 3y - 12)$$
$$4x + 8y - 14 \leq -9x + 3y - 12$$
$$4x + 8y - 14 \leq -9x + 3y - 12$$

# FO[ $\mathbb{Z}, +$ ] — Presburger arithmetic

**Theorem** [Presburger '29] Every FO formula  $\phi$  over  $(\mathbb{Z}, +, 0, 1, \leq, |)$  can be effectively transformed into an equivalent quantifier-free formula  $\phi^*$

## Proof idea

Show how to remove an innermost quantifier  $Qz$  from  $\phi = \dots Qz \alpha(\dots, z)$

Assume:

- $Qz = \exists z$  (if not, treat  $\forall z$  as  $\neg \exists z \neg$ )
- $\alpha$  is  $\vee$ -free (if not, commute  $\exists$  and  $\vee$ )

## Example

temporarily assume formulas  
are over the reals or the rationals...

$$\exists z \alpha(x, y, z) = \exists z (2x + 4y - 3z \leq 7) \wedge (3x - y + 2z \leq -4)$$

$$\exists z (2x + 4y - 7 \leq 3z) \wedge (2z \leq -3x + y - 4)$$

$$\exists z (4x + 8y - 14 \leq 6z) \wedge (6z \leq -9x + 3y - 12)$$

$$4x + 8y - 14 \leq -9x + 3y - 12$$

$$(4-9)x + (8-3)y - (14-12) \leq 0$$

# FO[ $\mathbb{Z}, +$ ] — Presburger arithmetic

**Theorem** [Presburger '29] Every FO formula  $\phi$  over  $(\mathbb{Z}, +, 0, 1, \leq, |)$  can be effectively transformed into an equivalent quantifier-free formula  $\phi^*$

## Proof idea

Show how to remove an innermost quantifier  $Qz$  from  $\phi = \dots Qz \alpha(\dots, z)$

Assume:

- $Qz = \exists z$  (if not, treat  $\forall z$  as  $\neg \exists z \neg$ )
- $\alpha$  is  $\vee$ -free (if not, commute  $\exists$  and  $\vee$ )

## Example

temporarily assume formulas  
are over the reals or the rationals...

$$\exists z \alpha(x, y, z) = \exists z (2x + 4y - 3z \leq 7) \wedge (3x - y + 2z \leq -4)$$

$$\exists z (2x + 4y - 7 \leq 3z) \wedge (2z \leq -3x + y - 4)$$

$$\exists z (4x + 8y - 14 \leq 6z) \wedge (6z \leq -9x + 3y - 12)$$

$$4x + 8y - 14 \leq -9x + 3y - 12$$

$$(-5)x + (5)y - (2) \leq 0$$

# FO[ $\mathbb{Z}, +$ ] — Presburger arithmetic

**Theorem** [Presburger '29] Every FO formula  $\phi$  over  $(\mathbb{Z}, +, 0, 1, \leq, |)$  can be effectively transformed into an equivalent quantifier-free formula  $\phi^*$

## Proof idea

Show how to remove an innermost quantifier  $Qz$  from  $\phi = \dots Qz \alpha(\dots, z)$

Assume:

- $Qz = \exists z$  (if not, treat  $\forall z$  as  $\neg \exists z \neg$ )
- $\alpha$  is  $\vee$ -free (if not, commute  $\exists$  and  $\vee$ )

## Example

temporarily assume formulas  
are over the reals or the rationals...

$$\exists z \alpha(x, y, z) = \exists z (2x + 4y - 3z \leq 7) \wedge (3x - y + 2z \leq -4)$$

$$\exists z (2x + 4y - 7 \leq 3z) \wedge (2z \leq -3x + y - 4)$$

$$\exists z (4x + 8y - 14 \leq 6z) \wedge (6z \leq -9x + 3y - 12)$$

$$4x + 8y - 14 \leq -9x + 3y - 12$$

$$-5x + 5y - 2 \leq 0$$

# FO[ $\mathbb{Z}, +$ ] — Presburger arithmetic

**Theorem** [Presburger '29] Every FO formula  $\phi$  over  $(\mathbb{Z}, +, 0, 1, \leq, |)$  can be effectively transformed into an equivalent quantifier-free formula  $\phi^*$

## Proof idea

Show how to remove an innermost quantifier  $Qz$  from  $\phi = \dots Qz \alpha(\dots, z)$

Assume:

- $Qz = \exists z$  (if not, treat  $\forall z$  as  $\neg \exists z \neg$ )
- $\alpha$  is  $\vee$ -free (if not, commute  $\exists$  and  $\vee$ )

## Example

$$\exists z \alpha(x, y, z) = \exists z (2x + 4y - 3z \leq 7) \wedge (3x - y + 2z \leq -4)$$

$$\exists z (2x + 4y - 7 \leq 3z) \wedge (2z \leq -3x + y - 4)$$

$$\exists z (4x + 8y - 14 \leq 6z) \wedge (6z \leq -9x + 3y - 12)$$

$$4x + 8y - 14 \leq -9x + 3y - 12$$

$$-5x + 5y - 2 \leq 0$$

~~temporarily assume formulas  
are over the reals or the rationals...~~

# FO[ $\mathbb{Z}, +$ ] — Presburger arithmetic

**Theorem** [Presburger '29] Every FO formula  $\phi$  over  $(\mathbb{Z}, +, 0, 1, \leq, |)$  can be effectively transformed into an equivalent quantifier-free formula  $\phi^*$

## Proof idea

Show how to remove an innermost quantifier  $Qz$  from  $\phi = \dots Qz \alpha(\dots, z)$

Assume:

- $Qz = \exists z$  (if not, treat  $\forall z$  as  $\neg \exists z \neg$ )
- $\alpha$  is  $\vee$ -free (if not, commute  $\exists$  and  $\vee$ )

## Example

$$\exists z \alpha(x, y, z) = \exists z (2x + 4y - 3z \leq 7) \wedge (3x - y + 2z \leq -4)$$

$$\exists z (2x + 4y - 7 \leq 3z) \wedge (2z \leq -3x + y - 4)$$

$$\exists z (4x + 8y - 14 \leq 6z) \wedge (6z \leq -9x + 3y - 12)$$

$$4x + 8y - 14 + m \leq -9x + 3y - 12$$

$$-5x + 5y - 2 + m \leq 0$$

~~temporarily assume formulas  
are over the reals or the rationals...~~

# FO[ $\mathbb{Z}, +$ ] — Presburger arithmetic

**Theorem** [Presburger '29] Every FO formula  $\phi$  over  $(\mathbb{Z}, +, 0, 1, \leq, |)$  can be effectively transformed into an equivalent quantifier-free formula  $\phi^*$

## Proof idea

Show how to remove an innermost quantifier  $Qz$  from  $\phi = \dots Qz \alpha(\dots, z)$

Assume:

- $Qz = \exists z$  (if not, treat  $\forall z$  as  $\neg \exists z \neg$ )
- $\alpha$  is  $\vee$ -free (if not, commute  $\exists$  and  $\vee$ )

## Example

$$\exists z \alpha(x, y, z) = \exists z (2x + 4y - 3z \leq 7) \wedge (3x - y + 2z \leq -4)$$

~~temporarily assume formulas  
are over the reals or the rationals...~~

$$\exists z (2x + 4y - 7 \leq 3z) \wedge (2z \leq -3x + y - 4)$$

$$\exists z (4x + 8y - 14 \leq 6z) \wedge (6z \leq -9x + 3y - 12)$$

$$6 \mid 4x + 8y - 14 + m \wedge 4x + 8y - 14 + m \leq -9x + 3y - 12$$

$$6 \mid 4x + 8y - 14 + m \wedge -5x + 5y - 2 + m \leq 0$$

# FO[ $\mathbb{Z}, +$ ] — Presburger arithmetic

**Theorem** [Presburger '29] Every FO formula  $\phi$  over  $(\mathbb{Z}, +, 0, 1, \leq, |)$  can be effectively transformed into an equivalent quantifier-free formula  $\phi^*$

## Proof idea

Show how to remove an innermost quantifier  $Qz$  from  $\phi = \dots Qz \alpha(\dots, z)$

Assume:

- $Qz = \exists z$  (if not, treat  $\forall z$  as  $\neg \exists z \neg$ )
- $\alpha$  is  $\vee$ -free (if not, commute  $\exists$  and  $\vee$ )

## Example

$$\exists z \alpha(x, y, z) = \exists z (2x + 4y - 3z \leq 7) \wedge (3x - y + 2z \leq -4)$$

~~temporarily assume formulas  
are over the reals or the rationals...~~

$$\exists z (2x + 4y - 7 \leq 3z) \wedge (2z \leq -3x + y - 4)$$

$$\exists z (4x + 8y - 14 \leq 6z) \wedge (6z \leq -9x + 3y - 12)$$

$$\forall_{m=0,\dots,5} \quad 6 \mid 4x + 8y - 14 + m \wedge 4x + 8y - 14 + m \leq -9x + 3y - 12$$

$$\forall_{m=0,\dots,5} \quad 6 \mid 4x + 8y - 14 + m \wedge -5x + 5y - 2 + m \leq 0$$



# Some fancy FO theories

$\text{FO}[\mathbb{N}, +, \cdot]$  = Peano arithmetic

 **UNDECIDABLE**   
(reduction from H's 10th)

$\text{FO}[\mathbb{R}, +, \cdot]$  = Arithmetic theory of real numbers

 **DECIDABLE**   
(quantifier elimination)

$\text{FO}[\mathbb{Z}, +]$  = Presburger arithmetic

 **DECIDABLE**   
(quantifier elimination)

$\text{FO}[\mathbb{N}^2, \leq_1, \leq_2]$  = First-order theory of the unlabelled grid

$\text{FO}[\{0,1\}, =]$   $\approx$  {Valid QBFs}

$\text{FO}[V_R, E_R]$  = First-order theory of “random” graph

$\text{FO}[C_M, T_M]$  = First-order theory of the transition graph of a Turing machine  $M$

# Some fancy FO theories

$\text{FO}[\mathbb{N}, +, \cdot]$  = Peano arithmetic

 **UNDECIDABLE**   
(reduction from H's 10th)

$\text{FO}[\mathbb{R}, +, \cdot]$  = Arithmetic theory of real numbers

 **DECIDABLE**   
(quantifier elimination)

$\text{FO}[\mathbb{Z}, +]$  = Presburger arithmetic

 **DECIDABLE**   
(quantifier elimination)

$\text{FO}[\mathbb{N}^2, \leq_1, \leq_2]$  = First-order theory of the unlabelled grid

 **DECIDABLE**   
(interpreted in the former)

$\text{FO}[\{0,1\}, =]$   $\approx$  {Valid QBFs}

$\text{FO}[V_R, E_R]$  = First-order theory of “random” graph

$\text{FO}[C_M, T_M]$  = First-order theory of the transition graph of a Turing machine  $M$

# Some fancy FO theories

$\text{FO}[\mathbb{N}, +, \cdot]$  = Peano arithmetic

 **UNDECIDABLE**   
(reduction from H's 10th)

$\text{FO}[\mathbb{R}, +, \cdot]$  = Arithmetic theory of real numbers

 **DECIDABLE**   
(quantifier elimination)

$\text{FO}[\mathbb{Z}, +]$  = Presburger arithmetic

 **DECIDABLE**   
(quantifier elimination)

$\text{FO}[\mathbb{N}^2, \leq_1, \leq_2]$  = First-order theory of the unlabelled grid

 **DECIDABLE**   
(interpreted in the former)

$\text{FO}[\{0,1\}, =]$   $\approx$  {Valid QBFs}

$\text{FO}[V_R, E_R]$  = First-order theory of “random” graph

$\text{FO}[C_M, T_M]$  = First-order theory of the transition graph of a Turing machine M

**Lemma**      Given any **QBF**  $\phi$  without free variables,  
one can construct an **FO formula**  $\phi^*$  such that

$$\models \phi \quad \text{iff} \quad (\{0,1\}, =) \models \phi^*$$

# FO[{0,1}, =] — The FO theory of Boolean algebra

**Lemma**      Given any **QBF**  $\phi$  without free variables,  
one can construct an **FO formula**  $\phi^*$  such that

$$\models \phi \quad \text{iff} \quad (\{0,1\}, =) \models \phi^*$$

**Proof**

define  $\phi^* = \exists t \phi[x / (x=t)]$  (for all bound variables  $x$ )

# FO[{0,1}, =] — The FO theory of Boolean algebra

**Lemma**      Given any QBF  $\phi$  without free variables,  
one can construct an FO formula  $\phi^*$  such that

$$\models \phi \quad \text{iff} \quad (\{0,1\}, =) \models \phi^*$$

**Proof**

define  $\phi^* = \exists t \phi[x / (x=t)]$  (for all bound variables  $x$ )

**Corollary**      FO[{0,1}, =] encodes the set of valid QBF formulas

# Some fancy FO theories

$\text{FO}[\mathbb{N}, +, \cdot]$  = Peano arithmetic

 **UNDECIDABLE**   
(reduction from H's 10th)

$\text{FO}[\mathbb{R}, +, \cdot]$  = Arithmetic theory of real numbers

 **DECIDABLE**   
(quantifier elimination)

$\text{FO}[\mathbb{Z}, +]$  = Presburger arithmetic

 **DECIDABLE**   
(quantifier elimination)

$\text{FO}[\mathbb{N}^2, \leq_1, \leq_2]$  = First-order theory of the unlabelled grid

 **DECIDABLE**   
(interpreted in the former)

$\text{FO}[\{0,1\}, =]$   $\approx$  {Valid QBFs}

EASY

$\text{FO}[V_R, E_R]$  = First-order theory of “random” graph

$\text{FO}[C_M, T_M]$  = First-order theory of the transition graph of a Turing machine  $M$

# Some fancy FO theories

$\text{FO}[\mathbb{N}, +, \cdot]$  = Peano arithmetic

 **UNDECIDABLE**   
(reduction from H's 10th)

$\text{FO}[\mathbb{R}, +, \cdot]$  = Arithmetic theory of real numbers

 **DECIDABLE**   
(quantifier elimination)

$\text{FO}[\mathbb{Z}, +]$  = Presburger arithmetic

 **DECIDABLE**   
(quantifier elimination)

$\text{FO}[\mathbb{N}^2, \leq_1, \leq_2]$  = First-order theory of the unlabelled grid

 **DECIDABLE**   
(interpreted in the former)

$\text{FO}[\{0,1\}, =]$   $\approx$  {Valid QBFs}

EASY

$\text{FO}[V_R, E_R]$  = First-order theory of “random” graph

$\text{FO}[C_M, T_M]$  = First-order theory of the transition graph of a Turing machine M



# FO[ $V_R, E_R$ ] — The FO theory of the “random” graph

A different perspective and a coarser view on expressiveness...

What percentage of finite graphs verify a given FO sentence?



# Probability of a formula

$P_n[\phi]$  = probability that  $\phi$  holds on a random finite graph with  $n$  nodes

# Probability of a formula

$P_n[\phi]$  = probability that  $\phi$  holds on a random finite graph with  $n$  nodes

$$P_\infty[\phi] = \lim_{n \rightarrow \infty} P_n[\phi]$$

# Probability of a formula

$P_n[\phi]$  = probability that  $\phi$  holds on a random finite graph with  $n$  nodes

$$P_\infty[\phi] = \lim_{n \rightarrow \infty} P_n[\phi]$$

**Example** For  $\phi$  = “the graph is complete”,

$$\text{we have } P_n[\phi] = \frac{1}{2^{n(n-1)}}$$

$$\text{and hence } P_\infty[\phi] = 0$$

# Probability of a formula

## Theorem (0/1 Law)

[Glebskii et al. '69, Fagin '76]

Every FO formula  $\phi$  is

either almost surely true ( $P_\infty[\phi] = 1$ )

or almost surely false ( $P_\infty[\phi] = 0$ )

# Probability of a formula

## Theorem (0/1 Law)

[Glebskii et al. '69, Fagin '76]

Every FO formula  $\phi$  is

either almost surely true ( $P_\infty[\phi] = 1$ )

or almost surely false ( $P_\infty[\phi] = 0$ )

## Examples

- $\phi = \text{"there is a triangle"}$

$$P_\infty[\phi] = 1$$

# Probability of a formula

## Theorem (0/1 Law)

[Glebskii et al. '69, Fagin '76]

Every FO formula  $\phi$  is

either almost surely true ( $P_\infty[\phi] = 1$ )

or almost surely false ( $P_\infty[\phi] = 0$ )

## Examples

- $\phi =$  “there is a triangle”
- $\phi =$  “there no 5-clique”

$$P_\infty[\phi] = 1$$

$$P_\infty[\phi] = 0$$

# Probability of a formula

## Theorem (0/1 Law)

[Glebskii et al. '69, Fagin '76]

Every FO formula  $\phi$  is

either almost surely true ( $P_\infty[\phi] = 1$ )

or almost surely false ( $P_\infty[\phi] = 0$ )

## Examples

- $\phi =$  “there is a triangle”
- $\phi =$  “there no 5-clique”
- $\phi =$  “even number of edges”
- $\phi =$  “even number of nodes”

$$P_\infty[\phi] = 1$$

$$P_\infty[\phi] = 0$$

**Your turn!**



# Probability of a formula

## Theorem (0/1 Law)

[Glebskii et al. '69, Fagin '76]

Every FO formula  $\phi$  is

either almost surely true ( $P_\infty[\phi] = 1$ )

or almost surely false ( $P_\infty[\phi] = 0$ )

## Examples

- $\phi =$  “there is a triangle”
- $\phi =$  “there no 5-clique”
- $\phi =$  “even number of edges”
- $\phi =$  “even number of nodes”

$$P_\infty[\phi] = 1$$

$$P_\infty[\phi] = 0$$

$$P_\infty[\phi] = 1/2$$

**Your turn!**

$P_\infty[\phi]$  not even defined

# Probability of a formula

## Theorem (0/1 Law)

[Glebskii et al. '69, Fagin '76]

Every FO formula  $\phi$  is

either almost surely true ( $P_\infty[\phi] = 1$ )

or almost surely false ( $P_\infty[\phi] = 0$ )

## Examples

- $\phi =$  “there is a triangle”
- $\phi =$  “there no 5-clique”
- $\phi =$  “even number of edges”
- $\phi =$  “even number of nodes”
- $\phi =$  “more edges than nodes”

$$P_\infty[\phi] = 1$$

$$P_\infty[\phi] = 0$$

$$P_\infty[\phi] = 1/2$$

$P_\infty[\phi]$  not even defined

$$P_\infty[\phi] = 1$$


( yet not FO-definable... )

**Your turn!**

# The “random” infinite graph

Every FO formula  $\phi$  is either almost surely true or almost surely false,  
*and this depends on whether  $(V_R, E_R) \models \phi$*

The “random” graph  
 $(V_R, E_R)$






# The “random” infinite graph

Every FO formula  $\phi$  is either almost surely true or almost surely false,  
*and this depends on whether*  $(V_R, E_R) \models \phi$

The “random” graph  
 $(V_R, E_R)$




each pair of nodes  $i, j$   
is connected with  
probability  $1/2$

# The “random” infinite graph

Every FO formula  $\phi$  is either almost surely true or almost surely false,  
*and this depends on whether*  $(V_R, E_R) \models \phi$

The “random” graph  
 $(V_R, E_R)$






# The “random” infinite graph

Every FO formula  $\phi$  is either almost surely true or almost surely false,  
*and this depends on whether*  $(V_R, E_R) \models \phi$

The “random” graph  
 $(V_R, E_R)$



each pair of nodes  $i, j$   
is connected with  
probability  $1/2$

each pair of nodes  $i, j$   
is connected if  
 $i$ -th bit of  $j$  is 1

the unique  
graph that  
satisfies  
 $\delta_k$  for all  $k$

# Probability of a formula - application


**Theorem** [Grandjean '83]

One can decide in **PSPACE** whether  $\phi$  is almost surely true on finite graphs

# Probability of a formula - application

**Theorem** [Grandjean '83]

One can decide in **PSPACE** whether  $\phi$  is almost surely true on finite graphs






# Probability of a formula - application

**Theorem** [Grandjean '83]


One can decide in **PSPACE** whether  $\phi$  is almost surely true on finite graphs



# Probability of a formula - application


**Theorem [Grandjean '83]**

One can decide in **PSPACE** whether  $\phi$  is almost surely true on finite graphs



**Model-checking on large graphs/databases**


Don't bother checking the formula,  
either it's *almost surely true* or *almost surely false*!



# Probability of a formula - application

**Theorem** [Grandjean '83]

One can decide in **PSPACE** whether  $\phi$  is almost surely true on finite graphs




Disclaimer:

0/1 Law only  
applies applies to  
unconstrained graphs

Model-checking on large graphs/databases

Don't bother checking the formula,  
either it's *almost surely true* or *almost surely false*!



# Some fancy FO theories

$\text{FO}[\mathbb{N}, +, \cdot]$  = Peano arithmetic

 **UNDECIDABLE**   
(reduction from H's 10th)

$\text{FO}[\mathbb{R}, +, \cdot]$  = Arithmetic theory of real numbers

 **DECIDABLE**   
(quantifier elimination)

$\text{FO}[\mathbb{Z}, +]$  = Presburger arithmetic

 **DECIDABLE**   
(quantifier elimination)

$\text{FO}[\mathbb{N}^2, \leq_1, \leq_2]$  = First-order theory of the unlabelled grid

 **DECIDABLE**   
(interpreted in the former)

$\text{FO}[\{0,1\}, =]$   $\approx$  {Valid QBFs}

EASY

$\text{FO}[V_R, E_R]$  = First-order theory of “random” graph

 **DECIDABLE**   
(0/1 Law)

$\text{FO}[C_M, T_M]$  = First-order theory of the transition graph of a Turing machine M

# Some fancy FO theories

$\text{FO}[\mathbb{N}, +, \cdot]$  = Peano arithmetic

 **UNDECIDABLE**   
(reduction from H's 10th)

$\text{FO}[\mathbb{R}, +, \cdot]$  = Arithmetic theory of real numbers

 **DECIDABLE**   
(quantifier elimination)

$\text{FO}[\mathbb{Z}, +]$  = Presburger arithmetic

 **DECIDABLE**   
(quantifier elimination)

$\text{FO}[\mathbb{N}^2, \leq_1, \leq_2]$  = First-order theory of the unlabelled grid

 **DECIDABLE**   
(interpreted in the former)

$\text{FO}[\{0,1\}, =]$   $\approx$  {Valid QBFs}

EASY

$\text{FO}[V_R, E_R]$  = First-order theory of “random” graph

 **DECIDABLE**   
(0/1 Law)

$\text{FO}[C_M, T_M]$  = First-order theory of the transition graph of a Turing machine M

 **DECIDABLE**   
(automatic structure)

# Things to remember





# Things to remember

- FO is cool and quite expressive
- Model-checking is decidable (in **PSPACE**) when the universe is finite  
Satisfiability, validity, equivalence are all undecidable (reduction from Domino)
- For infinite universes, one can fix a model and study its FO theory  
Some FO theories are decidable, some are not
- Some FO theories can be reduced to others via FO interpretations

