

Multistate models

Bendix Carstensen Steno Diabetes Center Copenhagen
Herlev, Denmark
<http://BendixCarstensen.com>

SPE, Tartu, Estonia,

June 2023

<http://BendixCarstensen.com/SPE>

Multistate models

Bendix Carstensen, Martyn Plummer

Multistate models

SPE, Tartu, Estonia,

June 2023

<http://BendixCarstensen.com/SPE>

ms-Markov

Common assumptions in survival analysis

1. Subjects are **either** “healthy” **or** “diseased”, with no intermediate state.

Common assumptions in survival analysis

1. Subjects are **either** “healthy” **or** “diseased”, with no intermediate state.
2. The disease is **irreversible**, or requires intervention to be cured.

Common assumptions in survival analysis

1. Subjects are **either** “healthy” **or** “diseased”, with no intermediate state.
2. The disease is **irreversible**, or requires intervention to be cured.
3. The time of disease incidence is known **exactly**.

Common assumptions in survival analysis

1. Subjects are **either** “healthy” **or** “diseased”, with no intermediate state.
2. The disease is **irreversible**, or requires intervention to be cured.
3. The time of disease incidence is known **exactly**.
4. The disease is **accurately** diagnosed.

Common assumptions in survival analysis

1. Subjects are **either** “healthy” **or** “diseased”, with no intermediate state.
2. The disease is **irreversible**, or requires intervention to be cured.
3. The time of disease incidence is known **exactly**.
4. The disease is **accurately** diagnosed.

Common assumptions in survival analysis

1. Subjects are **either** “healthy” **or** “diseased”, with no intermediate state.
2. The disease is **irreversible**, or requires intervention to be cured.
3. The time of disease incidence is known **exactly**.
4. The disease is **accurately** diagnosed.

These assumptions are true for **death** and many **chronic diseases**.

Common assumptions in survival analysis

1. Subjects are **either** “healthy” **or** “diseased”, with no intermediate state.
2. The disease is **irreversible**, or requires intervention to be cured.
3. The time of disease incidence is known **exactly**.
4. The disease is **accurately** diagnosed.

These assumptions are true for **death** and many **chronic diseases**.

A question of definition:

- consider occurrence of **recording of** a given disease

A model for cervical cancer

Invasive squamous cell cancer of the cervix is preceded by cervical intraepithelial neoplasia (CIN)



A model for cervical cancer

Invasive squamous cell cancer of the cervix is preceded by cervical intraepithelial neoplasia (CIN)



Purpose of a screening programme is to detect and treat CIN — status of persons obtained at screening dates

A model for cervical cancer

Invasive squamous cell cancer of the cervix is preceded by cervical intraepithelial neoplasia (CIN)



Purpose of a screening programme is to detect and treat CIN — status of persons obtained at screening dates

Aim of the modeling the **transition rates** between states, is to be able predict how population moves between states

A model for cervical cancer

Invasive squamous cell cancer of the cervix is preceded by cervical intraepithelial neoplasia (CIN)



Purpose of a screening programme is to detect and treat CIN — status of persons obtained at screening dates

Aim of the modeling the transition rates between **states**, is to be able predict how population moves between **states**

A model for cervical cancer

Invasive squamous cell cancer of the cervix is preceded by cervical intraepithelial neoplasia (CIN)



Purpose of a screening programme is to detect and treat CIN — status of persons obtained at screening dates

Aim of the modeling the transition rates between states, is to be able predict how population moves between states

- Transition rates between states

A model for cervical cancer

Invasive squamous cell cancer of the cervix is preceded by cervical intraepithelial neoplasia (CIN)



Purpose of a screening programme is to detect and treat CIN — status of persons obtained at screening dates

Aim of the modeling the transition rates between states, is to be able predict how population moves between states

- ▶ Transition rates between states
- ▶ Probability of state occupancy

Markov models for multistate processes

The natural generalization of Poisson regression to multiple disease states:

Markov models for multistate processes

The natural generalization of Poisson regression to multiple disease states:

- ▶ transition between states depends **only** on current state

Markov models for multistate processes

The natural generalization of Poisson regression to multiple disease states:

- ▶ transition between states depends **only** on current state
- ▶ — this is the **Markov** property

Markov models for multistate processes

The natural generalization of Poisson regression to multiple disease states:

- ▶ transition between states depends **only** on current state
- ▶ — this is the **Markov** property
- ▶ \Rightarrow transition rates are constant over time

Markov models for multistate processes

The natural generalization of Poisson regression to multiple disease states:

- ▶ transition between states depends **only** on current state
- ▶ — this is the **Markov** property
- ▶ \Rightarrow transition rates are constant over time
- ▶ (time-fixed) covariates may influence transition rates

Markov models for multistate processes

The natural generalization of Poisson regression to multiple disease states:

- ▶ transition between states depends **only** on current state
- ▶ — this is the **Markov** property
- ▶ \Rightarrow transition rates are constant over time
- ▶ (time-fixed) covariates may influence transition rates
- ▶ the formal Markov property is **very** restrictive

Markov models for multistate processes

The natural generalization of Poisson regression to multiple disease states:

- ▶ transition between states depends **only** on current state
- ▶ — this is the **Markov** property
- ▶ \Rightarrow transition rates are constant over time
- ▶ (time-fixed) covariates may influence transition rates
- ▶ the formal Markov property is **very** restrictive
- ▶ in the clinical literature “Markov model” is often used about any type of multistate model

Components of a multistate (Markov) model

- ▶ Define the disease states

Components of a multistate (Markov) model

- ▶ Define the disease states
- ▶ Define which transitions between states are allowed

Components of a multistate (Markov) model

- ▶ Define the disease states
- ▶ Define which transitions between states are allowed
- ▶ Select covariates influencing transition rates (may be different between transitions)

Components of a multistate (Markov) model

- ▶ Define the disease states
- ▶ Define which transitions between states are allowed
- ▶ Select covariates influencing transition rates (may be different between transitions)
- ▶ Not a trivial task — do we want *e.g.*


Components of a multistate (Markov) model

- ▶ Define the disease states
- ▶ Define which transitions between states are allowed
- ▶ Select covariates influencing transition rates (may be different between transitions)
- ▶ Not a trivial task — do we want *e.g.*
 - ▶ cause of death (CVD, Cancer, Other)

Components of a multistate (Markov) model

- ▶ Define the disease states
- ▶ Define which transitions between states are allowed
- ▶ Select covariates influencing transition rates (may be different between transitions)
- ▶ Not a trivial task — do we want *e.g.*
 - ▶ cause of death (CVD, Cancer, Other)
 - ▶ disease status at death (prev.CVD, prev.Can, neither)

A more complicated multistate model



Likelihood for a multistate model

- ▶ The likelihood of the model depends on the probability of being in state j at time t_1 , given that you were in state i at time t_0 .

Likelihood for a multistate model

- ▶ The likelihood of the model depends on the probability of being in state j at time t_1 , given that you were in state i at time t_0 .
- ▶ Assume transition rates constant in small time intervals

Likelihood for a multistate model

- ▶ The likelihood of the model depends on the probability of being in state j at time t_1 , given that you were in state i at time t_0 .
- ▶ Assume transition rates constant in small time intervals
- ▶ \Rightarrow each interval for a person contributes term(s) to the likelihood

Likelihood for a multistate model

- ▶ The likelihood of the model depends on the probability of being in state j at time t_1 , given that you were in state i at time t_0 .
- ▶ Assume transition rates constant in small time intervals
- ▶ \Rightarrow each interval for a person contributes term(s) to the likelihood
- ▶ one term for each possible transition between states

Likelihood for a multistate model

- ▶ The likelihood of the model depends on the probability of being in state j at time t_1 , given that you were in state i at time t_0 .
- ▶ Assume transition rates constant in small time intervals
- ▶ \Rightarrow each interval for a person contributes term(s) to the likelihood
- ▶ one term for each possible transition between states
- ▶ the total likelihood for person p in intervals i is a product of these terms, $d_{pi} \log(\lambda_{pi}) - \lambda_{pi} y_{pi}$

Likelihood for a multistate model

- ▶ The likelihood of the model depends on the probability of being in state j at time t_1 , given that you were in state i at time t_0 .
- ▶ Assume transition rates constant in small time intervals
- ▶ \Rightarrow each interval for a person contributes term(s) to the likelihood
- ▶ one term for each possible transition between states
- ▶ the total likelihood for person p in intervals i is a product of these terms, $d_{pi} \log(\lambda_{pi}) - \lambda_{pi} y_{pi}$
- ▶ \Rightarrow each term has the form of the likelihood for a Poisson variate d with mean λy

Likelihood for a multistate model

- ▶ each term has the form of the likelihood for a Poisson variate d with mean λy

Likelihood for a multistate model

- ▶ each term has the form of the likelihood for a Poisson variate d with mean λy
- ▶ terms are **not** independent, but the total likelihood is a product; hence of the same form as the likelihood from independent Poisson variates

Likelihood for a multistate model

- ▶ each term has the form of the likelihood for a Poisson variate d with mean λy
- ▶ terms are **not** independent, but the total likelihood is a product; hence of the same form as the likelihood from independent Poisson variates
- ▶ but observations from intervals from one person are neither Poisson nor independent

Realms of multistate modeling

- ▶ intensities — dimension time^{-1}

Realms of multistate modeling

- ▶ intensities — dimension time^{-1}
- ▶ state probabilities — dimensionless, time^0
integral of intensities w.r.t. to time

Realms of multistate modeling

- ▶ intensities — dimension time^{-1}
- ▶ state probabilities — dimensionless, time^0
integral of intensities w.r.t. to time
- ▶ sojourn times — dimension time^1
integral of state probabilities w.r.t. to time

Realms of multistate modeling

- ▶ intensities — dimension time^{-1}
- ▶ state probabilities — dimensionless, time^0
 integral of intensities w.r.t. to time
- ▶ sojourn times — dimension time^1
 integral of state probabilities w.r.t. to time

Classes of multistate models

- ▶ Markov model: transition between states depends **only** on current state \Rightarrow transition rates are constant
time-homogeneous Markov model

Classes of multistate models

- ▶ Markov model: transition between states depends **only** on current state \Rightarrow transition rates are constant
time-homogeneous Markov model
- ▶ If transition rates depend on the **same timescale** only we have a time-**inhomogeneous** Markov model

Classes of multistate models

- ▶ Markov model: transition between states depends **only** on current state \Rightarrow transition rates are constant
time-homogeneous Markov model
- ▶ If transition rates depend on the **same timescale** only we have a time-**inhomogeneous** Markov model
- ▶ If transition rates depend on the time since entry to the current state we have a **semi**-Markov model

Classes of multistate models

- ▶ Markov model: transition between states depends **only** on current state \Rightarrow transition rates are constant
time-homogeneous Markov model
- ▶ If transition rates depend on the **same timescale** only we have a time-**inhomogeneous** Markov model
- ▶ If transition rates depend on the time since entry to the current state we have a **semi**-Markov model
- ▶ If transition rates depend on several timescales we have a general multistate model (there is no name for this)

Classes of multistate models

- ▶ Markov model: transition between states depends **only** on current state \Rightarrow transition rates are constant
time-homogeneous Markov model
- ▶ If transition rates depend on the **same timescale** only we have a time-inhomogeneous Markov model
- ▶ If transition rates depend on the time since entry to the current state we have a semi-Markov model
- ▶ If transition rates depend on several timescales we have a general multistate model (there is no name for this)

Computing state probabilities from intensities in multistate models

- ▶ time-homogeneous Markov model:
closed-form formulae exist

Computing state probabilities from intensities in multistate models

- ▶ time-homogeneous Markov model:
closed-form formulae exist
- ▶ time-**inh**omogeneous Markov model:
closed-form formulae exist (a bit more complicated)

Computing state probabilities from intensities in multistate models

- ▶ time-homogeneous Markov model:
closed-form formulae exist
- ▶ time-**inh**omogeneous Markov model:
closed-form formulae exist (a bit more complicated)
- ▶ **semi**-Markov model:
no closed form formulae exist

Computing state probabilities from intensities in multistate models

- ▶ time-homogeneous Markov model:
closed-form formulae exist
- ▶ time-**inh**omogeneous Markov model:
closed-form formulae exist (a bit more complicated)
- ▶ **semi**-Markov model:
no closed form formulae exist
- ▶ general multistate model:
no closed form formulae exist

Computing state probabilities from intensities in multistate models

- ▶ time-homogeneous Markov model:
closed-form formulae exist
- ▶ time-**inh**omogeneous Markov model:
closed-form formulae exist (a bit more complicated)
- ▶ **semi**-Markov model:
no closed form formulae exist
- ▶ general multistate model:
no closed form formulae exist

Computing state probabilities from intensities in multistate models

- ▶ time-homogeneous Markov model:
closed-form formulae exist
- ▶ time-**inh**omogeneous Markov model:
closed-form formulae exist (a bit more complicated)
- ▶ **semi**-Markov model:
no closed form formulae exist
- ▶ general multistate model:
no closed form formulae exist

No formulae means that any inference on state probabilities and sojourn times must be based on **simulation** from the model.

Multistate models with Lexis

Bendix Carstensen

Multistate models

SPE, Tartu, Estonia,

June 2023

<http://BendixCarstensen.com/SPE>

ms-Lexis

Example: Renal failure data from Steno

Hovind P, Tarnow L, Rossing P, Carstensen B, and Parving H-H: Improved survival in patients obtaining remission of nephrotic range albuminuria in diabetic nephropathy. *Kidney Int.*, 66(3):1180–1186, 2004.

Example: Renal failure data from Steno

Hovind P, Tarnow L, Rossing P, Carstensen B, and Parving H-H: Improved survival in patients obtaining remission of nephrotic range albuminuria in diabetic nephropathy. *Kidney Int.*, 66(3):1180–1186, 2004.

- ▶ Endpoint of interest: Death or end stage renal disease (ESRD), i.e. dialysis or kidney transplant.

Example: Renal failure data from Steno

Hovind P, Tarnow L, Rossing P, Carstensen B, and Parving H-H: Improved survival in patients obtaining remission of nephrotic range albuminuria in diabetic nephropathy. *Kidney Int.*, 66(3):1180–1186, 2004.



- ▶ Endpoint of interest: Death or end stage renal disease (ESRD), i.e. dialysis or kidney transplant.
- ▶ 96 patients entering at nephrotic range albuminuria (NRA), i.e. $\text{U-alb} > 300\text{mg/day}$.

Example: Renal failure data from Steno

Hovind P, Tarnow L, Rossing P, Carstensen B, and Parving H-H: Improved survival in patients obtaining remission of nephrotic range albuminuria in diabetic nephropathy. *Kidney Int.*, 66(3):1180–1186, 2004.

- ▶ Endpoint of interest: Death or end stage renal disease (ESRD), i.e. dialysis or kidney transplant.
- ▶ 96 patients entering at nephrotic range albuminuria (NRA), i.e. $\text{U-alb} > 300\text{mg/day}$.
- ▶ Is remission from this condition (i.e return to $\text{U-alb} < 300\text{mg/day}$) predictive of the prognosis?

		Remission	
		Total	
			Yes No
No. patients		125	32 93
No. events		77	8 69
Follow-up time (years)		1084.7	259.9 824.8
Cox-model:			
Timescale:	Time since nephrotic range albuminuria (NRA)		
Entry:	2.5 years of GFR-measurements after NRA		
Outcome:	ESRD or Death		
Estimates:	RR	95% c.i.	p
Fixed covariates:			
Sex (F vs. M):	0.92	(0.53,1.57)	0.740
Age at NRA (per 10 years):	1.42	(1.08,1.87)	0.011
Time-dependent covariate:			
Obtained remission:	0.28	(0.13,0.59)	0.001



Features of the analysis

- ▶ Remission is included as a time-dependent variable.

```
renal[1:5,]  
id      dob      doe      dor      dox  event  
17 1967.944 1996.013      NA 1997.094      2  
26 1959.306 1989.535 1989.814 1996.136      1  
27 1962.014 1987.846      NA 1993.239      3  
33 1950.747 1995.243 1995.717 2003.993      0  
42 1961.296 1987.884 1996.650 2003.955      0
```

Note patient 26, 33 and 42 obtain remission.

Features of the analysis

- ▶ Remission is included as a time-dependent variable.
- ▶ Age at entry is included as a fixed variable.

```
renal[1:5,]  
id      dob      doe      dor      dox event  
17 1967.944 1996.013      NA 1997.094      2  
26 1959.306 1989.535 1989.814 1996.136      1  
27 1962.014 1987.846      NA 1993.239      3  
33 1950.747 1995.243 1995.717 2003.993      0  
42 1961.296 1987.884 1996.650 2003.955      0
```

Note patient 26, 33 and 42 obtain remission.

```

> Lr <- Lexis(entry = list(per = doe,
+                           age = doe-dob,
+                           tfi = 0),
+             exit = list(per = dox),
+             exit.status = event>0,
+             states = c("NRA", "ESRD"),
+             data = renal)
> summary(Lr)


```

Transitions:


To

From	NRA	ESRD	Records:	Events:	Risk time:	Persons:
NRA	48	77	125	77	1084.67	125

```
> boxes(Lr, boxpos = list(x = c(25, 75),  
+                           y = c(75, 25)),  
+       scale.R = 100, show.BE = TRUE )
```



Illness-death model



λ : reversion rate.

μ_{NRA} : mortality/ESRD rate **before** reversion.

μ_{rem} : mortality/ESRD rate **after** reversion.

Cutting follow-up at remission: cutLexis


```
> Lc <- cutLexis(Lr, cut = Lr$dor,  
+               timescale = "per",  
+               new.state = "Rem",  
+               precursor.states = "NRA")  
> summary(Lc)
```

Transitions:

	To						
From	NRA	Rem	ESRD	Records:	Events:	Risk time:	Persons:
NRA	24	29	69	122	98	824.77	122
Rem	0	24	8	32	8	259.90	32
Sum	24	53	77	154	106	1084.67	125

Showing states and FU: boxes.Lexis

```
> boxes(Lc, boxpos = list(x = c(15, 85, 50),  
+                          y = c(85, 85, 20)),  
+       scale.R = 100, show.BE = TRUE)
```



Cutting follow up at events: cutLexis

```
> Lc <- cutLexis( Lr, cut = Lr$dor,  
+               timescale = "per",  
+               new.state = "Rem",  
+               precursor.states = "NRA",  
+               split.states = TRUE )  
> summary( Lc )
```

Transitions:

To

From	NRA	Rem	ESRD	ESRD(Rem)	Records:	Events:	Risk time:	Persons:
NRA	24	29	69	0	122	98	824.77	122
Rem	0	24	0	8	32	8	259.90	32
Sum	24	53	69	8	154	106	1084.67	125

Cutting follow up at events: cutLexis

```
> Lc <- cutLexis( Lr, cut = Lr$dor,  
+               timescale = "per",  
+               new.state = "Rem",  
+               precursor.states = "NRA",  
+               split.states = TRUE )  
> summary( Lc )
```

Transitions:

To

From	NRA	Rem	ESRD	ESRD(Rem)	Records:	Events:	Risk time:	Persons:
NRA	24	29	69	0	122	98	824.77	122
Rem	0	24	0	8	32	8	259.90	32
Sum	24	53	69	8	154	106	1084.67	125

Cutting follow up at events: cutLexis

```
> Lc <- cutLexis( Lr, cut = Lr$dor,  
+               timescale = "per",  
+               new.state = "Rem",  
+               precursor.states = "NRA",  
+               split.states = TRUE )  
> summary( Lc )
```


Transitions:

To

From	NRA	Rem	ESRD	ESRD(Rem)	Records:	Events:	Risk time:	Persons:
NRA	24	29	69	0	122	98	824.77	122
Rem	0	24	0	8	32	8	259.90	32
Sum	24	53	69	8	154	106	1084.67	125

Showing states and FU: boxes.Lexis

```
> boxes(Lc, boxpos = list(x = c(15, 85, 15, 85),  
+                          y = c(85, 85, 20, 20)),  
+       scale.R = 100)
```



Likelihood for a general MS-model

- ▶ Product of likelihoods for each transition
— each one as for a survival model

Likelihood for a general MS-model

- ▶ Product of likelihoods for each transition
— each one as for a survival model
- ▶ **Risk time** is the risk time in the “From” state

Likelihood for a general MS-model


- ▶ Product of likelihoods for each transition
— each one as for a survival model
- ▶ **Risk time** is the risk time in the “From” state
- ▶ **Events** are transitions to the “To” state

Likelihood for a general MS-model

- ▶ Product of likelihoods for each transition
— each one as for a survival model
- ▶ **Risk time** is the risk time in the “From” state
- ▶ **Events** are transitions to the “To” state
- ▶ All other transitions out of “From” are treated as **censorings**


Likelihood for a general MS-model

- ▶ Product of likelihoods for each transition
— each one as for a survival model
- ▶ **Risk time** is the risk time in the “From” state
- ▶ **Events** are transitions to the “To” state
- ▶ All other transitions out of “From” are treated as **censorings**
- ▶ Possible to fit models separately for each transition



Cox-analysis with remission as time-dependent covariate:


- Ignores λ , the remission rate.



Cox-analysis with remission as time-dependent covariate:

- ▶ Ignores λ , the remission rate.
- ▶ Assumes μ_{NRA} and μ_{rem} use the same timescale.

Model for all transitions




Cox-model:

- Different timescales for transitions possible

Poisson-model:

Model for all transitions




Cox-model:

- ▶ Different timescales for transitions possible
- ▶ ... only one per transition

Poisson-model:

Model for all transitions




Cox-model:

- ▶ Different timescales for transitions possible
- ▶ ... only one per transition
- ▶ No explicit representation of estimated rates.

Poisson-model:

Model for all transitions




Cox-model:

- ▶ Different timescales for transitions possible
- ▶ ... only one per transition
- ▶ No explicit representation of estimated rates.

Poisson-model:

- ▶ Timescales can be different

Model for all transitions




Cox-model:

- ▶ Different timescales for transitions possible
- ▶ ... only one per transition
- ▶ No explicit representation of estimated rates.

Poisson-model:

- ▶ Timescales can be different
- ▶ Multiple timescales can be accommodated simultaneously

Model for all transitions



Cox-model:

- ▶ Different timescales for transitions possible
- ▶ ... only one per transition
- ▶ No explicit representation of estimated rates.

Poisson-model:

- ▶ Timescales can be different
- ▶ Multiple timescales can be accommodated simultaneously
- ▶ Explicit representation of all transition rates

Calculating state probabilities

P {Remission **before** time t }

$$= \int_0^t \lambda(u) \exp \left(- \int_0^u \lambda(s) + \mu_{\text{NRA}} \, ds \right) \, du$$

P {Being in remission **at** time t }

$$= \int_0^t \lambda(u) \exp \left(- \int_0^u \lambda(s) + \mu_{\text{NRA}}(s) \, ds \right) \times \\ \exp \left(- \int_u^t \mu_{\text{rem}}(s) \, ds \right) \, du$$

Note μ_{rem} could also depend on u , time since obtained remission.

Sketch of programming, assuming that λ (`lambda`), μ_{NRA} (`mu.nra`) and μ_{rem} (`mu.rem`) are known at any age (stored in vectors)

```
c.rem      <- cumsum(lambda)
c.mort.nra <- cumsum(mu.nra)
c.mort.rem <- cumsum(mu.rem)
pr1 <- cumsum(lambda * exp(-(c.rem + c.mort.nra)))

intgr(t,s) <-
function(t,s){
  lambda[s] * exp(-(c.rem[s] + c.mort.nra[s])) *
    exp(-(c.mort.rem[t] - c.mort.rem[s]))
}
for(t in 1:100) p2[t] <- sum(intgr(t,1:t))
```

If μ_{rem} also depends on time since remission, then `c.mort.rem` should have an extra argument—technically very complicated