

Poisson and Binary Regression

Janne Pitkäniemi

Finnish Cancer Registry

Tampere university

Statistical Practice in Epidemiology (2026,Tartu)

Elapse of time and Epidemiology

Epidemiology deals with the occurrence of event (disease) in populations observed over time


- ▶ concepts of risk and rate are used to measure the frequency with which the event (disease) cases occur
- ▶ **risk** is defined as $\frac{D}{N}$, where D is the number of people who developed the disease during pre-specified follow-up from 0 to t and N is the number of disease-free population at the beginning of follow-up and
- ▶ **rate** is defined as $\frac{D}{Y}$, where Y is the amount of person-time at risk observed when following disease free subjects from 0 to t.
- ▶ Note: risk increases with t but rate can vary depending on the length of the follow-up period.
- ▶ **Virtually all prospective follow-up studies include loss to follow-up censoring and risk must be estimated using appropriate methods described in this course.**

Points to be covered

- ▶ Incidence rates, rate ratios and rate differences from *follow-up studies* can be computed by fitting *Poisson regression models*.
- ▶ Risk ratios and differences can be computed from binary data by fitting *Logistic regression models*.
- ▶ Both models are special instances of *Generalized linear models*.
- ▶ There are various ways to do these tasks in R.


The Estonian Biobank cohort: survival among the elderly

Follow-up of 60 random individuals aged 75-103 at recruitment, until death (●) or censoring (○) in April 2014 (linkage with the Estonian Causes of Death Registry). (time-scale: calendar time).



The Estonian Biobank cohort: survival among the elderly

Follow-up time for 60 random individuals aged 75-103 at recruitment (time-scale: time in study).



Events, dates and risk time

- ▶ Mortality as the outcome:
 - d: indicator for **status** at exit:
 - 1: death observed
 - 0: censored alive

- ▶ Dates:

doe = date of **E**ntry to follow-up,
dox = date of e**X**it, end of follow-up.

- ▶ Follow-up time (years) computed as:

$$y = (\text{dox} - \text{doe}) / 365.25$$

Crude overall rate computed by hand and model

Total no. cases, person-years & rate (/1000 y):

```
> D <- sum( d ); Y <- sum(y) ; R <- D/(Y/1000)
```

```
> round( c(D=D, Y=Y, R=R), 2)
```

D	Y	R
884.00	11678.24	75.70

R-implementation of the rate estimation with Poisson regression:

A model with offset term

```
> m1 <- glm( D ~ 1, family=poisson,  
  offset=log(Y))
```

```
> coef(m1)  
( Intercept )  
-2.581
```

A model with poisreg—family (Epi package)

```
> glm(cbind(D, Y) ~1, family=poisreg)
```

```
Coefficients :  
( Intercept )  
-2.581
```

From the coefficient we get estimate of the rate $\exp(-2.581) * 1000 = 75.70$

Constant hazard — Poisson model

Let $Y \sim \exp(\lambda)$, then $f(y; \lambda) = \lambda e^{-\lambda y} I(y > 0)$

Constant rate model: $\lambda(y) = \lambda$ and observed data $\{(y_i, \delta_i); i = 1, \dots, n\}$.

The likelihood $L(\lambda) = \prod_{i=1}^n \lambda^{\delta_i} e^{-\lambda y_i}$ and

$$\log(L) = \sum_{i=1}^n [\delta_i \log(\lambda) - \lambda y_i]$$

Solving the *score equations*:

$$\frac{\partial \log L(\lambda)}{\partial \lambda} = \sum \left[\frac{\delta_i}{\lambda} - y_i \right] = \frac{D}{\lambda} - Y = 0 \text{ and } D - \lambda Y = 0$$

→ **maximum likelihood estimator (MLE)** of λ :

$$\hat{\lambda} = \frac{D}{Y} = \frac{\text{number of cases}}{\text{total person-time}} = \text{empirical rate!}$$

offset term — Poisson model

- ▶ Previous model without offset: Intercept $6.784 = \log(884)$
- ▶ We should use an offset if we suspect that the underlying **population sizes (person-years) differ** for each of the observed counts – For example varying person-years by sex, age, treatment group, ...
- ▶ We need a term in the model that "scales" the likelihood, but does not depend on model parameters (include a **term with reg. coef. fixed to 1**) – offset term is $\log(y)$
- ▶ This is all taken care of by family=poisreg – recommend to use

$$\begin{aligned}\log\left(\frac{\mu}{y}\right) &= \beta_0 + \beta_1 x_1 \\ \log(\mu) &= 1 \times \log(y) + \beta_0 + \beta_1 x_1\end{aligned}$$

Comparing rates: The Thorotrast Study

- ▶ Cohort of seriously ill patients in Denmark on whom angiography of brain was performed.
- ▶ Exposure: contrast medium used in angiography,
 1. `thor` = thorotrast (with ^{232}Th), used 1935-50
 2. `ctrl` = other medium (?), used 1946-63
- ▶ Outcome of interest: death

`doe` = date of **E**ntry to follow-up,

`dox` = date of e**X**it, end of follow-up.

- ▶ `data(thoro)` in the `Epi` package.

Tabulating rates: thorotrast vs. control

Tabulating cases, person-years & rates by group

```
> stat.table( contrast ,  
+             list ( N = count(),  
+                   D = sum(d),  
+                   Y = sum(y),  
+                   rate = ratio(d,y,1000) ) )
```

contrast	N	D	Y	rate
ctrl	1236	797.00	30517.56	26.12
thor	807	748.00	19243.85	38.87

Rate ratio estimation with Poisson regression

- ▶ Include contrast as the explanatory variable (factor).
- ▶ Insert person years in units that you want rates in

```
> m2 <- glm( cbind(d,y/1000) ~ contrast,family = poisreg(link="log") )  
> round( summary(m2)$coef, 4)[, 1:2]
```

	Estimate	Std. Error
(Intercept)	3.2626	0.0354
contrast thor	0.3977	0.0509

- ▶ Rate ratio and CI?

Call function `ci.exp()` in `Epi`

```
> round( ci.exp( m2 ), 3 )  
              exp(Est.)  2.5%  97.5%  
(Intercept)    26.116 24.364 27.994  
contrast thor     1.488  1.347  1.644
```

Rates in groups with Poisson regression

- ▶ Include contrast as the explanatory variable (factor).
- ▶ Remove the intercept (-1)
- ▶ Insert person-years in units that you want rates in

```
> m3 <- glm( cbind(d,y/1000) ~ factor(contrast)-1,family = poisreg)
> round( summary(m3)$coef, 4)[, 1:2]
```

		Estimate	Std. Error
contrast	ctrl	3.2626	0.0354
contrast	thor	3.6602	0.0366

```
> round( ci.exp( m3 ), 3 )
```

		exp(Est.)	2.5%	97.5%
contrast	ctrl	26.116	24.364	27.994
contrast	thor	38.870	36.181	41.757

Rate difference estimation with Poisson regression

- The approach with d/y enables additive rate models too:

```
> contrast<-c(0,1)
> m5 <-glm(cbind(d,y/1000) ~contrast,
            family=poisreg(link="identity") )
> round( ci.exp(m5,Exp=F), 3 )
```

	Estimate	2.5%	97.5%
(Intercept)	26.116	24.303	27.929
contrast thor	12.753	9.430	16.077

Binary data: Treatment success Y/N

85 diabetes-patients with foot-wounds:

- ▶ Dalterapin (Dal)
- ▶ Placebo (Pl)

Treatment/Placebo given to diabetes patients, the design is prospective and outcome is measured better(Y)/worse(N). Is the probability of outcome more than 15% – yes, then use the risk difference or risk ratio (RR)

	Treatment group	
	Dalterapin	Placebo
Better	29	20
Worse	14	22
Total	43	42

$$\hat{p}_{\text{Dal}} = \frac{29}{43} = 67\% \quad \hat{p}_{\text{Pl}} = \frac{20}{42} = 47\%$$

Binary data: Crosstabulation analysis of 2x2 table

```
> library(Epi)
> dlt <- rbind( c(29,14), c(20,22) )
> colnames( dlt ) <- c("Better","Worse")
> rownames( dlt ) <- c("Dal","Pl")
> kable(twoby2( dlt ), "latex")
```

2 by 2 table analysis:

	Better	Worse	P(Better)	95% conf. interval
Dal	29	14	0.6744	0.5226 0.7967
Pl	20	22	0.4762	0.3316 0.6249

95% conf. interval

Relative Risk: 1.4163 0.9694 2.0692

Sample Odds Ratio: 2.2786 0.9456 5.4907

Conditional MLE Odds Ratio: 2.2560 0.8675 6.0405

Probability difference: 0.1982 -0.0110 0.3850

Exact P-value: 0.0808

Asymptotic P-value: 0.0665

Binary regression - Estimation of risk ratio (Relative risk)

```
> library(Epi)
> library(xtable)
> dlt <- data.frame(rbind( c(29,14),c(20,22) ))
> colnames( dlt ) <- c("Better","Worse")
> dlt$trt <- c(1,0)
> b2<-glm(cbind(Better,Worse)~trt,
+         family=binomial(link="log"),
+         data=dlt)
> xtable(round( ci.exp( b2 ), digits=6 ))
```

	exp(Est.)	2.5%	97.5%
(Intercept)	0.48	0.35	0.65
trt	1.42	0.97	2.07

Diabetics with Dalterapin treatment are 1.4 times likely to get better than those treated with placebo

Binary regression - Estimation of risk difference

```
> library(Epi)
> library(xtable)
> dlt <- data.frame(rbind( c(29,14),c(20,22) ))
> colnames( dlt ) <- c("Better","Worse")
> dlt$trt <- c(1,0)
> b2<-glm(cbind(Better,Worse)~trt,
+         family=binomial(link="identity"),
+         data=dlt)
> xtable(round( ci.exp( b2,Exp=F ), digits=6 ))
```

	Estimate	2.5%	97.5%
(Intercept)	0.48	0.33	0.63
trt	0.20	-0.01	0.40

Twenty percent more of the Diabetics with Dalterapin treatment are getting better compared to Diabetics treated with placebo

Conclusion: What did we learn?

- ▶ Rates, their ratio and difference can be analysed by Poisson regression
- ▶ In Poisson models the response can be either:
 - ▶ case indicator d with `offset = log(y)`, or
 - ▶ case and person-years `cbind(d,y)` with `poisreg-family` (Epi-package)
- ▶ Both may be fitted on either grouped data, or individual records.
- ▶ Binary outcome can be modeled with binary regression.