FIBRED TABULATORS

S.P MTHIMKHLDJGJFDSLULU

Abstract. This paper defines tabulators in strict double fibration (i.e., double fibrations of double categories with tabulators). Additionally, we give a Beck-Chevalley Condition for fibred tabulators. Moreover, we show that fibred tabulators are characterized by a preservation property.


Introduction

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi.

1 Fibrations

In this section we recall some basic facts about fibrations and their examples.

Definition 1.1. Let $P\colon \mathcal{E} \to \mathcal{B}$ be a functor. A morphism $f\colon X \to Y$ in $\mathcal{E}$ is *cartesian* over $u\colon I \to J$ i.e., $P(f) = u$ if for any $v\colon K \to I$ in $\mathcal{B}$ and $h\colon Z \to Y$ over $u \circ v$ there is a unique morphism $g\colon Z \to X$ over v with $h = f \circ g$.



A morphism $k\colon X \to Y$ is called *vertical* if it lies over an identity in $\mathcal{B}$. For an object $J \in \mathcal{B}$, the fibre of P at J denoted by $P(J)$ is the subcategory of $\mathcal{E}$ whose morphisms k lie over id_J.

Definition 1.2. A functor $P\colon \mathcal{E} \to \mathcal{B}$ is called *fibred category* or a *fibration* if for every $Y \in \mathcal{E}$ and $u\colon I \to PY$ there is a cartesian morphism $\bar{u}(E)\colon u^*E \to E$ over u. If P is a fibration, a *cleavage* for P denoted $\varphi(u, E)$ is a choice of cartesian lifts for u at E.

1.1 Examples

Let $\mathcal{C}^2$ denote the slice category of $\mathcal{C}$ over itself

Example 1.3 (Fundamental Fibration)**.** Let $\mathcal{C}$ be a category with pullbacks. Then the functor $\mathrm{cod}\colon \mathcal{C}^2 \to \mathcal{C}$ is a fibration; a commutative square is a cartesian arrow in $\mathcal{C}^2$ precisely when it is a pullback.

Example 1.4. The domain functor $\mathrm{dom}\colon \mathcal{C}^2 \to \mathcal{C}$ is a fibration.

Example 1.5 (Family fibration)**.** Every category $\mathcal{C}$ gives rise to a *family fibration* $\mathcal{F}\colon Fam(\mathcal{C}) \to Set$. A morphism $(u, \{\varphi_i\}_{i\in I})$ is cartesian when every φ_i is an isomorphism. Then $\mathcal{F}$ is a fibration since given $u\colon I \to J$ and $\{Y_j\}_{j\in J}$, $(u, \{1_{Y_{u(i)}}\})\colon \{Y_{u(i)}\}_{i\in I} \to \{Y_j\}_{j\in J}$ is cartesian over u.

Date: October 26, 2026.
MSC2020: Primary 00A05, Secondary 00A66.

1.2 Fibred Finite Limits


In [Str18] fibred finite limits are given in terms of terminal objects and pullbacks. Furthermore, [Jac99] captures the notion of fibred terminal objects in **Fib** using fibred adjunctions.

Definition 1.6. A fibration $P\colon \mathcal{E} \to \mathcal{B}$ has fibred terminal objects if every fibre $P(I)$ has a terminal object 1_I and they are preserved by cartesianness.

In $\mathbf{Fib}/\mathcal{B}$, the category of fibrations over $\mathcal{B}$, a fibration P has fibred terminals if the fibred functor $F\colon P \to id_B$ has a right adjoint. Moreover, this property is characterized by preservation as follows.

Lemma 1.7. *Let $\mathcal{B}$ be a category with a terminal object $\mathbf{1}_\mathcal{B}$. Then $P\colon \mathcal{E} \to \mathcal{B}$ has fibred terminals if and only if $\mathcal{E}$ has a terminal $1_\mathcal{E}$ and $P(\mathbf{1}_\mathcal{E})$ is a terminal in $\mathcal{B}$.*

Proof. Consider the following diagram in **Fib**




□

Futhermore, we have the following theorem

Theorem 1.8. *If $\mathcal{B}$ has a terminal object $\mathbf{1}_\mathcal{B}$ and $P\colon \mathcal{E} \to \mathcal{B}$ is a fibration, the following are equivalent*

(1) P has a terminal object in $\mathbf{Fib}/\mathcal{B}$.
(2) P has a terminal object in $\mathbf{Fib}$.

Proof. By the lifting and factorization of adjoints



Given $!_\mathcal{B} \dashv \mathbf{1}_\mathcal{B}$, it follows that $P \dashv \mathbf{1}_\mathcal{E}$ if and only if $!_\mathcal{E} \dashv \mathbf{1}_\mathcal{E}$ over $!_\mathcal{B} \dashv \mathbf{1}_\mathcal{B}$. □

Theorem 1.9. *Let $\mathcal{B}$ be a category with pullbacks. A fibration $P\colon \mathcal{E} \to \mathcal{B}$ has fibred pullbacks if and only if $\mathcal{E}$ has and P preserves pullbacks.*

References

[Jac99] B. Jacobs. *Categorical Logic and Type Theory*, volume 141 of *Studies in Logic and the Foundations of Mathematics*. North Holland, 1999.
[Str18] Thomas Streicher. Fibered Categories a la Jean Benabou. Preprint, arXiv:1801.02927 [math.CT] (2018), 2018.

5 Melrose Road, Muizenberg, Cape Town, South Africa
Email address: siyam@aims.ac.za