



Math for Data

Dr. S. M.
Moosavi

Data Sets

Solutions

Mathematics for Data Science


Dr. S. M. Moosavi

smohsenmoosavi2009@gmail.com

July 28, 2024



The following slides are arranged (with some modifications) based on the book "*Math for Data Science*" by "**Omar Hijab**".



You can follow me on [Linkedin](#). Also, for course materials such as slides and the related python codes, see this [Github](#) repository.



Outline

Math for Data

Dr. S. M.
Moosavi

Data Sets

Solutions

1 Data Sets

2 Solutions



Outline

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction
Averages and Vector
Spaces
Two Dimensions
Complex Numbers
Mean and Covariance

Solutions

1 Data Sets

2 Solutions



What is a dataset?

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

Definition 1.1

Geometrically, a dataset is a sample of N points x_1, x_2, \dots, x_N in d -dimensional space \mathbb{R}^d . Algebraically, a dataset is an $N \times d$ matrix.

Practically speaking, the following are all representations of datasets:

matrix = CSV file = spreadsheet = SQL table = array = dataframe

Definition 1.2

Each point $x = (t_1, t_2, \dots, t_d)$ in the dataset is a sample or an example, and the components t_1, t_2, \dots, t_d of a sample point x are its features or attributes. As such, d -dimensional space \mathbb{R}^d is feature space.

Definition 1.3

Sometimes one of the features is separated out as the label. In this case, the dataset is a labelled dataset.



Iris dataset

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

The *Iris dataset* contains 150 examples of four features of Iris flowers, and there are three classes of Irises, *Setosa*, *Versicolor* and *Virginica*, with 50 samples from each class.

Samples (instances, observations)						Petal	
	Sepal length	Sepal width	Petal length	Petal width	Class label		
1	5.1	3.5	1.4	0.2	Setosa		
2	4.9	3.0	1.4	0.2	Setosa		
...							
50	6.4	3.5	4.5	1.2	Versicolor		
...							
150	5.9	3.0	5.0	1.8	Virginica		
Features (attributes, measurements, dimensions)					Class labels (targets)	Sepal	



MNIST dataset

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions


Complex Numbers

Mean and Covariance

Solutions

The *MNIST dataset* consists of 60,000 images of hand-written digits. There are 10 classes of images, corresponding to each digit $0, 1, \dots, 9$. We seek to compress the images while preserving as much as possible of the images' characteristics.

Each image is a grayscale 28×28 pixel image. Since $28^2 = 784$, each image is a point in $d = 784$ dimensions. Here there are $N = 60000$ samples and $d = 784$ features.





Exercises

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance


Solutions

Exercise 1.1

Use sklearn to download Iris dataset.

Exercise 1.2

- *From keras read the MNIST dataset.*
- *Let $(\text{train_X}, \text{train_y}), (\text{test_X}, \text{test_y}) = \text{mnist.load_data}()$*
- *Let $\text{pixels} = \text{train_X}[1]$.*
- *Do for loops over i and j in $\text{range}(28)$ and use scatter to plot points at location (i,j) with size given by $\text{pixels}[i,j]$, then show the following image.*





Introduction

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

Suppose we have a population of things (people, tables, numbers, vectors, images, etc.) and we have a sample of size N from this population:

$$\mathbf{1} = [\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N]$$

The total population is the *population* or the *sample space*.

Example 1.1

The sample space consists of all real numbers and we take $N = 5$ samples from

$$\mathbf{1} = [3.95, 3.20, 3.10, 5.55, 6.93]$$

Example 1.2

The sample space consists of all integers and we take $N = 5$ samples from

$$\mathbf{1} = [35, -32, -8, 45, -8]$$



Introduction

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

Example 1.3

The sample space consists of all Python strings and we take $N = 5$ samples from

```
l = ['a2e?', '%#T', '7y5', ' ', 'kkk>><</', '[]*+']
```

Example 1.4

The sample space consists of all HTML colors and we take $N = 5$ samples from

```
1 from random import choice
2 import matplotlib.pyplot as plt
3
4 def hexcolor():
5     return "#" + ''.join([choice('0123456789abcdef') for
6                             _ in range(6)])
7
8 for i in range(5): plt.scatter(i,0, c=hexcolor())
plt.show()
```



Mean

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

Let \mathbf{l} be a list as above. The goal is to compute the sample *average* or *mean* of the list, which is

$$\text{mean} = \text{average} = \frac{x_1 + x_2 + \cdots + x_N}{N}.$$

In the Example (1.1), the average is

$$\frac{3.95 + 3.20 + 3.10 + 5.55 + 6.93}{5} = 4.546.$$

Example 1.5

```
1  import numpy as np
2
3  dataset = np.array([3.95, 3.20, 3.10, 5.55, 6.93])
4  print(np.mean(dataset))
5
6  output: 4.546
```

In the Example (1.2), the average is $\frac{32}{5}$. In the Example (1.3), while we can add strings, we can't divide them by 5, so the average is undefined. Similarly for colors: the average is undefined.



Vector space

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

A sample space or population V is called a *vector space* if, roughly speaking, one can compute means or averages in V . In this case, we call the members of the population "vectors".

Definition 1.4 (Vector space)

Let V be a set. V is a vector space (over \mathbb{R}) if for every $u, v, w \in V$ and $r, s \in \mathbb{R}$:

- 1 *vectors can be added (and the sum $v + w$ is back in V);*
- 2 *vector addition is commutative $v + w = w + v$*
- 3 *vector addition is associative $u + (v + w) = (u + v) + w$;*
- 4 *there is a zero vector $\mathbf{0}$ ($\mathbf{0} + v = v$);*
- 5 *vectors v have negatives (or opposites) $-v$ ($v + (-v) = \mathbf{0}$);*
- 6 *vectors can be multiplied by real numbers (and the product rv is back in V);*
- 7 *multiplication is distributive over addition $(r + s)v = rv + sv$ and $r(u + v) = ru + rv$;*
- 8 *$1v = v$ and $0v = \mathbf{0}$;*
- 9 *$r(sv) = (rs)v$.*



Centered dataset

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance



Solutions

Definition 1.5 (Centered Versus Non-Centered)

If x_1, x_2, \dots, x_N is a dataset of points with mean m and

$$v_1 = x_1 - m, v_2 = x_2 - m, \dots, v_N = x_N - m,$$

then v_1, v_2, \dots, v_N is a centered dataset of vectors where its mean is zero.





Some notes

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces


Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

- When we work with vector spaces, numbers are referred to as *scalars*.
- When we multiply a vector v by a scalar r to get the scaled vector rv , we call it *scalar multiplication*.
- The set of all real numbers \mathbb{R} is a vector space.
- The set of all integers \mathbb{Z} is not a vector space.
- The set of all rational numbers \mathbb{Q} is a vector space over \mathbb{Q} but not over \mathbb{R} .
- The set of all Python strings is not a vector space.
- Usually, we can't take sample means from a population, we instead take the sample mean of a *statistic* associated to the population. A statistic is an assignment of a number $f(\text{item})$ to each item in the population. For example, the human population on Earth is not a vector space (they can't be added), but their heights is a vector space (heights can be added). For the Python strings, a statistic might be the length of the strings. For the HTML colors, a statistic is the HTML code of the color.



In general, a statistic need not be a number. A statistic can be anything that "behaves like a number". For example, $f(\text{item})$ can be a vector or a matrix. More generally, a statistic's values may be anything that lives in a vector space V .



Cartesian plane

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces


Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

The *cartesian plane* \mathbb{R}^2 , also called the 2-dimensional real space is a vector space.



For $\mathbf{v}_1 = (x_1, y_1), \mathbf{v}_2 = (x_2, y_2) \in \mathbb{R}^2$ and $t \in \mathbb{R}$ define

- $\mathbf{v}_1 + \mathbf{v}_2 = (x_1 + x_2, y_1 + y_2)$ (Addition).
- $\mathbf{0} = (0, 0)$ (Zero).
- $t\mathbf{v}_1 = (tx_1, ty_1)$ (Scaling).
- $-\mathbf{v}_1 = (-1)\mathbf{v}_1$ (Negative).
- $\mathbf{v}_1 - \mathbf{v}_2 = \mathbf{v}_1 + (-\mathbf{v}_2) = (x_1 - x_2, y_1 - y_2)$ (Subtraction).



Operations

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction


Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions





2d example

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

Example 1.6

```
1  import numpy as np
2
3  v1 = (1,2)
4  v2 = (3,4)
5  print(v1 + v2 == (1+3,2+4)) # returns False
6
7  v1 = [1,2]
8  v2 = [3,4]
9  print(v1 + v2 == [1+3,2+4]) # returns False
10
11 v1 = np.array([1,2])
12 v2 = np.array([3,4])
13 print(v1 + v2 == np.array([1+3,2+4]))
14 # returns [ True  True]
15 print(3*v1 == np.array([3,6]))
16 # returns [ True  True]
17 print(-v1 == np.array([-1,-2]))
18 # returns [ True  True]
19 print(v1 - v2 == np.array([1-3,2-4]))
20 # returns [ True  True]
```



2d example

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

For the two-dimensional dataset

$$\mathbf{x}_1 = (1, 2), \mathbf{x}_2 = (3, 4), \mathbf{x}_3 = (-2, 11), \mathbf{x}_4 = (0, 66),$$

or, equivalently,

$$\mathbf{x} = \begin{pmatrix} 1 & 2 \\ 3 & 4 \\ -2 & 11 \\ 0 & 66 \end{pmatrix},$$

the average is

$$\frac{(1, 2) + (3, 4) + (-2, 11) + (0, 66)}{4} = (0.5, 20.75).$$

Example 1.7

```
1 import numpy as np
2
3 dataset = np.array([[1,2], [3,4], [-2,11], [0,66]])
4 print(np.mean(dataset, axis=0))
5 # returns [ 0.5 , 20.75]
```



2d example

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers


Mean and Covariance

Solutions

Example 1.8

Generate a 2 dimensional dataset of random points and their mean

```
1 import numpy as np
2 from numpy.random import random as rd
3 import matplotlib.pyplot as plt
4 N = 20
5 dataset = np.array([[rd(), rd()] for _ in range(N)])
6 mean = np.mean(dataset,axis=0)
7 plt.grid()
8 X, Y = dataset[:,0], dataset[:,1]
9 plt.scatter(X,Y)
10 plt.scatter(*mean)
11 plt.annotate('$m$', xy=mean+0.01)
12 plt.show()
```





Magnitude

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

Definition 1.6 (Distance Formula)

If $\mathbf{v}_1 = (x_1, y_1)$ and $\mathbf{v}_2 = (x_2, y_2)$, then the distance between \mathbf{v}_1 and \mathbf{v}_2 is

$$|\mathbf{v}_1 - \mathbf{v}_2| = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}.$$

The distance of $\mathbf{v} = (x, y)$ to the origin $\mathbf{0} = (0, 0)$ is its magnitude or norm or length

$$r = |\mathbf{v}| = |\mathbf{v} - \mathbf{0}| = \sqrt{x^2 + y^2}.$$

Example 1.9

For $\mathbf{v}_1 = (1, 2)$ and $\mathbf{v}_2 = (3, 4)$

$$|\mathbf{v}_1| = \sqrt{1^2 + 2^2} = \sqrt{5} \simeq 2.236,$$

$$|\mathbf{v}_1 - \mathbf{v}_2| = \sqrt{(1 - 3)^2 + (2 - 4)^2} = \sqrt{4 + 4} = \sqrt{8} \simeq 2.828.$$

```

1  import numpy as np
2
3  v1 = np.array([1,2])
4  v2 = np.array([3,4])
5  print(np.linalg.norm(v1)) #returns 2.23606797749979
6  print(np.linalg.norm(v1-v2)) #returns 2.

```



Polar representation

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions


Complex Numbers

Mean and Covariance

Solutions

In terms of r and θ , the *polar representation* of (x, y) is

$$x = r \cos \theta, \quad y = r \sin \theta.$$



The *unit circle* consists of the vectors which are distance 1 from the origin $\mathbf{0}$. When \mathbf{v} is on the unit circle, the magnitude of \mathbf{v} is 1, and we say \mathbf{v} is a *unit vector*. In this case, the line formed by the scalings of \mathbf{v} intersects the unit circle at $\pm \mathbf{v}$.

When \mathbf{v} is a unit vector, then $r = 1$ and $\mathbf{v} = (x, y) = (\cos \theta, \sin \theta)$.



Polar representation

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

By the distance formula, a vector $\mathbf{v} = (x, y)$ is a unit vector when

$$x^2 + y^2 = 1.$$

More generally, any circle with *center* (a, b) and radius r consists of vectors $\mathbf{v} = (x, y)$ satisfying

$$(x - a)^2 + (y - b)^2 = r^2.$$

Let R be a point on the unit circle, and let $t > 0$. The scaled point tR is on the circle with center $(0, 0)$ and radius t . Moreover, if Q is any point, $Q + tR$ is on the circle with center Q and radius t . It is easy to check that $|t\mathbf{v}| = |t||\mathbf{v}|$ for any real number t and vector \mathbf{v} .

From this, if a vector \mathbf{v} is unit and $r > 0$, then $r\mathbf{v}$ has magnitude r . If \mathbf{v} is any vector not equal to the zero vector, then $r = |\mathbf{v}|$ is positive, and

$$\left| \frac{1}{r} \mathbf{v} \right| = \frac{1}{r} |\mathbf{v}| = \frac{1}{r} r = 1$$

so \mathbf{v}/r is a unit vector.



Inner product

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

Definition 1.7

Let $\mathbf{v}_1 = (x_1, y_1)$, $\mathbf{v}_2 = (x_2, y_2) \in \mathbb{R}^2$. The inner product or the dot product of \mathbf{v}_1 and \mathbf{v}_2 is given algebraically as

$$\mathbf{v}_1 \cdot \mathbf{v}_2 = x_1x_2 + y_1y_2.$$

From the geometric view, we have:

Theorem 1.1 (Dot Product Identity)

$$x_1x_2 + y_1y_2 = \mathbf{v}_1 \cdot \mathbf{v}_2 = |\mathbf{v}_1||\mathbf{v}_2| \cos \theta,$$

where θ is the angle between \mathbf{v}_1 and \mathbf{v}_2 .

Exercise 1.3

Prove the "Dot Product Identity", Theorem (1.1).

Hint: Use Pythagoras' theorem for general triangles.



The angle between two vectors

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

In Python, the dot product is given by `numpy.dot` and as a consequence of the dot product identity, we have the code for the angle between two vectors:

$$\theta_{\mathbf{v}_1, \mathbf{v}_2} = \arccos \left(\frac{\mathbf{v}_1 \cdot \mathbf{v}_2}{|\mathbf{v}_1| |\mathbf{v}_2|} \right).$$

Example 1.10

Find the angle between the vectors $\mathbf{v}_1 = (1, 2)$ and $\mathbf{v}_2 = (3, 4)$.

```
1  import numpy as np
2
3  def angle(u,v):
4      a = np.dot(u,v)
5      b = np.dot(u,u)
6      c = np.dot(v,v)
7      theta = np.arccos(a / np.sqrt(b*c))
8      return np.degrees(theta)
9
10 v1 = np.array([1,2])
11 v2 = np.array([3,4])
12 print(angle(v1,v2)) #returns 10.304846468766044 in
                        degree
```



Cauchy-Schwarz Inequality

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

Recall that $-1 \leq \cos \theta \leq 1$. Using the dot product identity, we obtain the important inequality:

Theorem 1.2 (Cauchy-Schwarz Inequality)

If u and v are any two vectors, then

$$-|u||v| \leq u \cdot v \leq |u||v|.$$

Exercise 1.4

Prove the "Cauchy-Schwarz Inequality".



2d linear equations system

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

Consider the homogeneous system

$$\begin{cases} ax + by = 0 \\ cx + dy = 0 \end{cases} \quad (1.1)$$

and let A be the 2×2 matrix

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}. \quad (1.2)$$

$(x, y) = (-b, a)$ is a solution of the first equation in (1.1). If we want this to be a solution of the second equation as well, we must have $cx + dy = ad - bc = 0$.

Definition 1.8 (Determinant)

The determinant of A is

$$\det(A) = \det \begin{pmatrix} a & b \\ c & d \end{pmatrix} = ad - bc.$$



2d linear equations system

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

Theorem 1.3 (Homogeneous System)

*When $\det(A) = 0$, the homogeneous system (1.1) has a nonzero solution, and all solutions are scalar multiples of $(x, y) = (-b, a)$.
When $\det(A) \neq 0$, the only solution is $(x, y) = (0, 0)$.*

For the inhomogeneous case

$$\begin{cases} ax + by = e \\ cx + dy = f \end{cases} \quad (1.3)$$

we have

Theorem 1.4 (Inhomogeneous System)

When $\det(A) \neq 0$, the inhomogeneous system (1.3) has the unique solution

$$\begin{pmatrix} x \\ y \end{pmatrix} = \frac{1}{\det(A)} \begin{pmatrix} de - bf \\ af - ce \end{pmatrix}.$$

When $\det(A) = 0$, (1.3) has a solution iff $ce = af$ and $de = bf$.



2d linear equations system

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

When $a^2 + b^2 \neq 0$, a solution is

$$\begin{pmatrix} x \\ y \end{pmatrix} = \frac{1}{a^2 + b^2} \begin{pmatrix} ae \\ be \end{pmatrix}.$$

When $c^2 + d^2 \neq 0$, a solution is

$$\begin{pmatrix} x \\ y \end{pmatrix} = \frac{1}{c^2 + d^2} \begin{pmatrix} cf \\ df \end{pmatrix}.$$

Any other solution differs from these solutions by a scalar multiple of the homogeneous solution $(x, y) = (-b, a)$.

Exercise 1.5

Prove the Theorems (1.3) and (1.4).



Complex numbers

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

Roughly speaking, the set of all *complex numbers* is the set of all points in \mathbb{R}^2 with different multiplication rule.

Definition 1.9 (Complex numbers)

The complex numbers, \mathbb{C} , is the set

$$\mathbb{C} = \{(x, y) \in \mathbb{R}^2\}$$

with operations

- *Addition:* $(x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2)$.
- *Scalar Multiplication:* $t(x, y) = (tx, ty)$
- *Multiplication:* $(x_1, y_1)(x_2, y_2) = (x_1x_2 - y_1y_2, x_1y_2 + x_2y_1)$.

Then, in \mathbb{C} , we have

- zero: $0 = (0, 0)$.
- opposite or additive inverse: $-(x, y) = (-x, -y)$.
- one: $1 = (1, 0)$.



Example

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

Example 1.11

- $(1, 2) + (3, 4) = (4, 6).$
- $(0, 0) + (1, 2) = (1, 2).$
- $3(1, 2) = (3, 6).$
- $(1, 0)(1, 2) = (1 - 0, 2 + 0) = (1, 2).$
- $(1, 2)(3, 4) = (3 - 8, 4 + 6) = (-5, 10).$
- $(x, 0) + (y, 0) = (x + y, 0).$
- $(x, 0)(y, 0) = (xy, 0).$

Note. By the last two examples, we see that complex numbers with 0 as their second component act like real numbers in addition and multiplication. So, from now on, we set $x = (x, 0).$

Example 1.12

- $0 = (0, 0).$
- $1 = (1, 0).$
- $-1 = (-1, 0).$



Imaginary number

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

Definition 1.10 (Imaginary number)

$$i = (0, 1).$$

Note. Python uses the symbol j for imaginary number.

Theorem 1.5

For each $z = (x, y) \in \mathbb{C}$, we can write

$$z = x + iy.$$

We call x as the real part of z , and y the imaginary part of z .

$$x = \text{Re}(z), \quad y = \text{Im}(z).$$

Proof. $x + iy = (x, 0) + (0, 1)(y, 0) = (x, 0) + (0 - 0, 0 + y) = (x, y).$

Theorem 1.6

$$i^2 = -1.$$

Proof. $i^2 = (0, 1)(0, 1) = (0 - 1, 0 + 0) = (-1, 0) = -1.$



Example

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

Example 1.13

In complex numbers:

- $\sqrt{-1} = i.$
- $\sqrt{-4} = 2i.$
- $(1, 2)(3, 4) = (1 + 2i)(3 + 4i)$
$$= 3 + 4i + 6i + 8i^2$$
$$= 3 + 10i - 8$$
$$= -5 + 10i$$
$$= (-5, 10).$$
- $(1, 2)^3 = (1 + 2i)^3$
$$= (1)^3 + 3(1)^2(2i) + 3(1)(2i)^2 + (2i)^3$$
$$= 1 + 6i + 12i^2 + 8i^3$$
$$= 1 + 6i - 12 - 8i$$
$$= -11 - 2i$$
$$= -(11, 2).$$



Conjugate

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

Definition 1.11 (Conjugate)

For $z = (x, y) \in \mathbb{C}$, the conjugate is

$$\bar{z} = (x, -y) = x - iy \in \mathbb{C}.$$

Some properties.

- $z + \bar{z} = 2\text{Re}(z)$, $z - \bar{z} = 2i\text{Im}(z)$.
- $z\bar{z} = \text{Re}(z)^2 + \text{Im}(z)^2$,

$$\Rightarrow |z| = \sqrt{\text{Re}(z)^2 + \text{Im}(z)^2} = \sqrt{z\bar{z}}$$

$$\Rightarrow |z|^2 = z\bar{z}.$$

Example 1.14

For $z = (4, -3) \in \mathbb{C}$:

- $\bar{z} = (4, 3) = 4 + 3i$,
- $z + \bar{z} = 2 \times 4 = 8$, $z - \bar{z} = 2i \times (-3) = -6i$.
- $z\bar{z} = (4)^2 + (-3)^2 = 16 + 9 = 25 \Rightarrow |z| = \sqrt{25} = 5$.
- $z^2 = (4 - 3i)^2 = 7 - 24i$.
- $|z|^2 = 25$.



Theorem 1.7

For a non-zero $z \in \mathbb{C}$, the inverse of z is

$$z^{-1} = \frac{1}{z} = \frac{\bar{z}}{z\bar{z}} = \frac{\bar{z}}{|z|^2}.$$

Proof. Firstly, if $z = (x, y)$ then $\frac{1}{z} \in \mathbb{C}$, because,

$$\frac{1}{z} = \frac{x - iy}{x^2 + y^2} = \left(\frac{x}{x^2 + y^2}, \frac{-y}{x^2 + y^2} \right) \in \mathbb{C}.$$

Secondly,

$$zz^{-1} = (x + iy) \left(\frac{x - iy}{x^2 + y^2} \right) = \frac{x^2 + y^2}{x^2 + y^2} = 1.$$

Corollary 1.1 (Division)

For $z_1 \in \mathbb{C}$ and $0 \neq z_2 \in \mathbb{C}$

$$\frac{z_1}{z_2} = z_1 z_2^{-1}.$$



Definitions

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

Definition 1.12 (Mean-squared distance)

Let $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N$ be a dataset, say D , in \mathbb{R}^d , and let $\mathbf{x} \in \mathbb{R}^d$. The mean-squared distance of \mathbf{x} to D is

$$MSD(\mathbf{x}) = \frac{1}{N} \sum_{k=1}^N |\mathbf{x}_k - \mathbf{x}|^2.$$

Definition 1.13 (Mean)

Let $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N$ be a dataset in \mathbb{R}^d . The mean or sample mean is

$$\mathbf{m} = \bar{\mathbf{x}}_N = \frac{1}{N} \sum_{k=1}^N \mathbf{x}_k = \frac{\mathbf{x}_1 + \mathbf{x}_2 + \dots + \mathbf{x}_N}{N}.$$

Theorem 1.8 (Point of Best-fit)

The mean is the point of best-fit: The mean minimizes the mean-squared distance to the dataset.

Exercise 1.6

Prove the Theorem (1.8).



Point of Best-fit

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

Example 1.15

```
1 import matplotlib.pyplot as plt
2 import numpy as np
3
4 np.random.seed(1)
5 N = 20
6 rnd = np.random.random
7 dataset = np.array([ [rnd(), rnd()] for _ in range(N) ])
8 # Mean
9 m = np.mean(dataset, axis=0)
10 #Random point
11 p = np.array([rnd(), rnd()])
12
13 plt.grid()
14 X, Y = dataset[:,0], dataset[:,1]
15 plt.scatter(X,Y)
16 for v in dataset:
17     plt.plot([m[0], v[0]], [m[1], v[1]], c='green')
18     plt.plot([p[0], v[0]], [p[1], v[1]], c='red')
19 plt.show()
20
21 # Comparison of MSD of the mean and a random point
22 MSD_m = np.sum(np.abs(dataset-m)**2)/N
23 MSD_p = np.sum(np.abs(dataset-p)**2)/N
24 print(MSD_m, MSD_p) # 0.160478187272121 0.5984208474157081
```



Point of Best-fit

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions




Figure 1.1: MSD for the mean (green) versus MSD for a random point (red).



Tensor product

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

For simplicity, let $\mathbf{u} = (a, b)$ and $\mathbf{v} = (c, d, e)$ be two vectors.

Definition 1.14 (Tensor product)

The tensor product of \mathbf{u} and \mathbf{v} is the matrix

$$\mathbf{u} \otimes \mathbf{v} = \begin{pmatrix} ac & ad & ae \\ bc & bd & be \end{pmatrix} = \begin{pmatrix} c\mathbf{u} & d\mathbf{u} & e\mathbf{u} \end{pmatrix} = \begin{pmatrix} a\mathbf{v} \\ b\mathbf{v} \end{pmatrix}$$

Definition 1.15 (Trace of a matrix)

The trace of a squared matrix A is the sum of the diagonal entries.

Note. For any vectors \mathbf{u}, \mathbf{v} and \mathbf{w} :

- $\mathbf{v} \otimes \mathbf{u} = (\mathbf{u} \otimes \mathbf{v})^t.$

In square case:

- $\det(\mathbf{u} \otimes \mathbf{v}) = 0.$

- $\text{trace}(\mathbf{u} \otimes \mathbf{v}) = \mathbf{u} \cdot \mathbf{v}.$

- $\text{trace}(\mathbf{u} \otimes \mathbf{u}) = |\mathbf{u}|^2.$

- $(\mathbf{u} \otimes \mathbf{v})\mathbf{w} = (\mathbf{v} \cdot \mathbf{w})\mathbf{u}.$



Covariance

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

Let $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N$ be a dataset in \mathbb{R}^d with \mathbf{m} as its mean.

Definition 1.16 (1d Covariance)

When $d = 1$, the covariance q is a scalar

$$q = \frac{1}{N} \sum_{k=1}^N (x_k - m)^2 = MSD(m).$$

In the scalar case, the covariance is called the variance of the scalar dataset.

In general, the covariance is a symmetric $d \times d$ matrix Q . We can center the dataset as

$$\mathbf{v}_1 = \mathbf{x}_1 - \mathbf{m}, \mathbf{v}_2 = \mathbf{x}_2 - \mathbf{m}, \dots, \mathbf{v}_N = \mathbf{x}_N - \mathbf{m}.$$

Then the *covariance matrix* is the $d \times d$ matrix Q as

$$Q = \frac{\mathbf{v}_1 \otimes \mathbf{v}_1 + \mathbf{v}_2 \otimes \mathbf{v}_2 + \dots + \mathbf{v}_N \otimes \mathbf{v}_N}{N}. \quad (1.4)$$



Example

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

Example 1.16

Suppose $N = 5$ and

$$\mathbf{x}_1 = (1, 2), \quad \mathbf{x}_2 = (3, 4), \quad \mathbf{x}_3 = (5, 6), \quad \mathbf{x}_4 = (7, 8), \quad \mathbf{x}_5 = (9, 10).$$

Then $\mathbf{m} = (5, 6)$ and

$$\mathbf{v}_1 = \mathbf{x}_1 - \mathbf{m} = (-4, -4), \quad \mathbf{v}_2 = \mathbf{x}_2 - \mathbf{m} = (-2, -2),$$

$$\mathbf{v}_3 = \mathbf{x}_3 - \mathbf{m} = (0, 0), \quad \mathbf{v}_4 = \mathbf{x}_4 - \mathbf{m} = (2, 2), \quad \mathbf{v}_5 = \mathbf{x}_5 - \mathbf{m} = (4, 4).$$

Since

$$(\pm 4, \pm 4) \otimes (\pm 4, \pm 4) = \begin{pmatrix} 16 & 16 \\ 16 & 16 \end{pmatrix},$$

$$(\pm 2, \pm 2) \otimes (\pm 2, \pm 2) = \begin{pmatrix} 4 & 4 \\ 4 & 4 \end{pmatrix},$$

$$(0, 0) \otimes (0, 0) = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix},$$

then

$$Q = \begin{pmatrix} 8 & 8 \\ 8 & 8 \end{pmatrix}.$$



Example

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

Example 1.17

```
1  import numpy as np
2
3  def tensor(u,v):
4      return np.array([ [ a*b for b in v] for a in u ])
5
6  np.random.seed(1)
7  N = 20
8  rnd = np.random.random
9  dataset = np.array([[rnd(), rnd()] for _ in range(N)])
10 # mean
11 m = np.mean(dataset,axis=0)
12 # center dataset
13 vectors = dataset - m
14 # covariance
15 Q = np.mean([ tensor(v,v) for v in vectors ],axis=0)
16 print(Q)
```



Note. The covariance matrix as written in (1.4) is the *biased covariance matrix*. If the denominator is instead $N - 1$, the matrix is the *unbiased covariance matrix*.

For datasets with large N , it doesn't matter, since N and $N - 1$ are almost equal.

In numpy, the Python covariance constructor is

Example 1.18

```
1  import numpy as np
2
3  np.random.seed(1)
4  N = 20
5  rnd = np.random.random
6  dataset = np.array([[rnd(), rnd()] for _ in range(N)])
7  # covariance
8  Q = np.cov(dataset, bias=True, rowvar=False)
9  print(Q)
```



Total variance

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

Definition 1.17 (Total variance)

From $\text{trace}(\mathbf{u} \otimes \mathbf{u}) = |\mathbf{u}|^2$, if Q is the covariance matrix then

$$\text{trace}(Q) = \frac{1}{N} \sum_{k=1}^N |\mathbf{x}_k - \mathbf{m}|^2. \quad (1.5)$$

We call (1.5) the total variance of the dataset. Thus the total variance equals $\text{MSD}(\mathbf{m})$.

Example 1.19

```
1  import numpy as np
2
3  np.random.seed(1)
4  N = 20
5  rnd = np.random.random
6  dataset = np.array([[rnd(), rnd()] for _ in range(N)])
7  # covariance
8  Q = np.cov(dataset.T, bias=True)
9  print(Q.trace()) # returns 0.16047818727212101
```



Projections

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces


Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

We would like to project a $2d$ dataset onto a line. Let \mathbf{u} be a unit vector (a vector of length one, $|\mathbf{u}| = 1$), and let $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_N$ be a $2d$ dataset, assumed for simplicity to be centered. We wish to project this dataset onto the line through \mathbf{u} . This will result in a $1d$ dataset.



When a vector \mathbf{v} is projected onto the line through \mathbf{u} , the length of the projected vector reads

$$|proj_{\mathbf{u}} \mathbf{v}| = |\mathbf{v}| \cos \theta,$$

where θ is the angle between the vectors \mathbf{v} and \mathbf{u} . Since $|\mathbf{u}| = 1$, this length equals the dot product $\mathbf{v} \cdot \mathbf{u}$. Hence the projected vector is

$$proj_{\mathbf{u}} \mathbf{v} = (\mathbf{v} \cdot \mathbf{u}) \mathbf{u}.$$



Projections

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

Hence,

Definition 1.18 (Reduced dataset)

The projected dataset of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_N$ onto the line through \mathbf{u} is the dataset

$$(\mathbf{v}_1 \cdot \mathbf{u})\mathbf{u}, (\mathbf{v}_2 \cdot \mathbf{u})\mathbf{u}, \dots, (\mathbf{v}_N \cdot \mathbf{u})\mathbf{u}.$$

The projected dataset is in \mathbb{R}^2 . The reduced dataset is

$$(\mathbf{v}_1 \cdot \mathbf{u}), (\mathbf{v}_2 \cdot \mathbf{u}), \dots, (\mathbf{v}_N \cdot \mathbf{u}),$$

which is in \mathbb{R} .

Exercise 1.7

Show that when a $2d$ dataset is centered then the mean of the reduced dataset is 0.

Exercise 1.8

Prove that if Q is the covariance matrix of a $2d$ dataset, then the variance of the projected dataset onto the line through the vector \mathbf{u} equals the quadratic function $\mathbf{u} \cdot Q\mathbf{u}$:

$$q = \frac{1}{N} \sum_{k=1}^N \mathbf{u} \cdot (\mathbf{v}_k \otimes \mathbf{v}_k) \mathbf{u} = \mathbf{u} \cdot Q\mathbf{u}.$$



Covariance ellipse

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

Hence,

Definition 1.19 (Covariance ellipse)

The contour of all points \mathbf{x} satisfying $\mathbf{x} \cdot Q\mathbf{x} = 1$ is the covariance ellipsoid. In two dimensions $d = 2$, this is the covariance ellipse. The contour of all points \mathbf{x} satisfying $\mathbf{x} \cdot Q^{-1}\mathbf{x} = 1$ is the inverse covariance ellipsoid. In two dimensions $d = 2$, this is the inverse covariance ellipse.

In two dimensions $d = 2$, a covariance matrix has the form

$$Q = \begin{pmatrix} a & b \\ b & c \end{pmatrix}.$$

If we write $\mathbf{u} = (x, y)$ for a vector in the plane, the covariance ellipse is

$$\mathbf{u} \cdot Q\mathbf{u} = (x, y) \cdot \begin{pmatrix} a & b \\ b & c \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = ax^2 + 2bxy + cy^2 = 1.$$

The covariance ellipse and inverse covariance ellipses described above are centered at the origin $(0, 0)$. When a dataset has mean \mathbf{m} and covariance Q , the ellipses are drawn centered at \mathbf{m} .

In particular, when $a = c$ and $b = 0$, then $Q = aI$ is a multiple of the identity, the inverse covariance ellipse is the circle of radius \sqrt{a} , and the covariance ellipse is the circle of radius $\frac{1}{\sqrt{a}}$.



Covariance ellipse I

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers


Mean and Covariance

Solutions

Example 1.20

Plot the contour ellipses for

$$Q_1 = \begin{pmatrix} 9 & 0 \\ 0 & 4 \end{pmatrix}, \quad Q_2 = \begin{pmatrix} 9 & 2 \\ 2 & 4 \end{pmatrix}.$$





Covariance ellipse II

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

```
1  import matplotlib.pyplot as plt
2  import numpy as np
3
4  def ellipse(a, b, c, levels, color):
5      L, delta = 4, .1
6      x = np.arange(-L,L,delta)
7      y = np.arange(-L,L,delta)
8      X,Y = np.meshgrid(x, y)
9      plt.contour(X, Y, a*X**2 + 2*b*X*Y + c*Y**2, levels,
                  colors=color)
10
11  # Q1 Covariance entities
12  a, b, c = 9, 0, 4
13
14  # Inverse Covariance entities
15  det = a*c - b**2
16  A, B, C = c/det, -b/det, a/det
17
18  plt.grid()
19  ellipse(a, b, c, [20], 'blue')
20  ellipse(A, B, C, [1], 'red')
21  plt.show()
```



Covariance ellipse III

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

```
22
23 # Q2 Covariance entities
24 a, b, c = 9, 2, 4
25
26 # Inverse Covariance entities
27 det = a*c - b**2
28 A, B, C = c/det, -b/det, a/det
29
30 plt.grid()
31 ellipse(a, b, c, [1], 'blue')
32 ellipse(A, B, C, [1], 'red')
33 plt.show()
```



Standardization

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

Here, we describe how to standardize datasets in \mathbb{R}^2 . *Standardizing* the dataset means to center the dataset and to place the x and y features on the same scale.

Consider the dataset

$\mathbf{x}_1 = (x_1, y_1), \mathbf{x}_2 = (x_2, y_2), \dots, \mathbf{x}_N = (x_N, y_N)$ with mean $\mathbf{m} = (m_x, m_y)$. Then the covariance matrix is

$$Q = \begin{pmatrix} a & b \\ b & c \end{pmatrix},$$

where

$$a = \frac{1}{N} \sum_{k=1}^N (x_k - m_x)^2, \quad b = \frac{1}{N} \sum_{k=1}^N (x_k - m_x)(y_k - m_y),$$

$$c = \frac{1}{N} \sum_{k=1}^N (y_k - m_y)^2.$$



Standardization

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

If a and c differ, the different scales of x 's and y 's distorts the relation between them, and b may not accurately reflect the correlation. To correct for this, we center and re-scale

$$x_1, x_2, \dots, x_N \rightarrow x'_1 = \frac{x_1 - m_x}{\sqrt{a}}, x'_2 = \frac{x_2 - m_x}{\sqrt{a}}, \dots, x'_N = \frac{x_N - m_x}{\sqrt{a}}$$

and

$$y_1, y_2, \dots, y_N \rightarrow y'_1 = \frac{y_1 - m_y}{\sqrt{c}}, y'_2 = \frac{y_2 - m_y}{\sqrt{c}}, \dots, y'_N = \frac{y_N - m_y}{\sqrt{c}}$$

This results in a new dataset

$\mathbf{v}_1 = (x'_1, y'_1), \mathbf{v}_2 = (x'_2, y'_2), \dots, \mathbf{v}_N = (x'_N, y'_N)$ that is centered:

$$\frac{\mathbf{v}_1 + \mathbf{v}_2 + \dots + \mathbf{v}_N}{N} = 0,$$

with each feature standardized to have unit variance,

$$\frac{1}{N} \sum_{k=1}^N x'_k = 1, \quad \frac{1}{N} \sum_{k=1}^N y'_k = 1.$$

This is the *standardized dataset*.



Standardization

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

The covariance matrix of the standardized dataset has the form

$$Q' = \begin{pmatrix} 1 & \rho \\ \rho & 1 \end{pmatrix},$$

where

$$\rho = \frac{1}{N} \sum_{k=1}^N x'_k y'_k = \frac{b}{\sqrt{ac}} = \frac{\sum_{k=1}^N (x_k - m_x)(y_k - m_y)}{\sqrt{\left(\sum_{k=1}^N (x_k - m_x)^2\right) \left(\sum_{k=1}^N (y_k - m_y)^2\right)}}$$

is the *Pearson correlation coefficient* of the dataset. The matrix Q' is the *correlation matrix*, or the *standardized covariance matrix*.

Example 1.21

$$Q = \begin{pmatrix} 9 & 2 \\ 2 & 4 \end{pmatrix} \Rightarrow \rho = \frac{b}{\sqrt{ac}} = \frac{1}{3} \Rightarrow Q' = \begin{pmatrix} 1 & 1/3 \\ 1/3 & 1 \end{pmatrix}.$$



Standardization

Math for Data

Dr. S. M.
Moosavi

Data Sets

Introduction

Averages and Vector
Spaces

Two Dimensions

Complex Numbers

Mean and Covariance

Solutions

From the Cauchy-Schwarz inequality, the correlation coefficient ρ is always between -1 and 1 . When $\rho = \pm 1$, the dataset samples are perfectly correlated and lie on a line passing through the mean.

When $\rho = 1$, the line has slope 1 , and when $\rho = -1$, the line has slope -1 . When $\rho = 0$, the dataset samples are completely uncorrelated and are considered two independent one-dimensional datasets (In standardized case).

In Python numpy, the correlation matrix is returned by

```
1 import numpy as np
2 np.corrcoef(dataset.T)
```

Here again, we input the transpose of the dataset if our default is vectors as rows.

Notice the $1/N$ cancels in the definition of ρ . Because of this, `corrcoef` is the same whether we deal with biased or unbiased covariance matrices.



Outline

Math for Data

Dr. S. M.
Moosavi

Data Sets

Solutions

1 Data Sets

2 Solutions



Chapter 1

Math for Data

Dr. S. M.
Moosavi

Data Sets

Solutions

Exercise 1.1.

```
1 from sklearn import datasets
2 iris = datasets.load_iris(as_frame=True)
3 dataset = iris["frame"]
```

Exercise 1.2.

- 1 Download file <https://s3.amazonaws.com/img-datasets/mnist.npz>
- 2 Move mnist.npz to .keras/datasets/ directory
- 3 Load data

Code 2.1: pixels

```
1 from keras.datasets import mnist
2 import matplotlib.pyplot as plt
3
4 (train_X, train_y), (test_X, test_y) =mnist.load_data()
5
6 pixels = train_X[1]
7
8 plt.grid()
9 for i in range(28):
10     for j in range(28): plt.scatter(i,j, s = pixels[i,j])
11 plt.show()
```




Chapter 1

Math for Data

Dr. S. M.
Moosavi

Data Sets

Solutions

Notice that for the code:

```
1 plt.imshow(pixels, cmap="gray_r")
```

we have




Figure 2.1: True pixels' image

To simulate Figure (2.1), we have to change our Code (2.1) to:



Chapter 1

Math for Data

Dr. S. M.
Moosavi


Data Sets

Solutions

Code 2.2: pixels

```
1 from keras.datasets import mnist
2 import matplotlib.pyplot as plt
3
4 (train_X, train_y), (test_X, test_y) =mnist.load_data()
5
6 pixels = train_X[1]
7
8 plt.grid()
9 plt.gca().invert_yaxis()
10 plt.axis('equal')
11 for i in range(28):
12     for j in range(28): plt.scatter(i,j, s = pixels[j,i])
13 plt.show()
```

The result is:





Chapter 1

Math for Data

Dr. S. M.
Moosavi

Data Sets

Solutions

Exercise 1.3. By Pythagoras' theorem for general triangles (Figure 2.2 (a)) we have

$$c^2 = a^2 + b^2 - 2ab \cos(\theta). \quad (2.1)$$




Figure 2.2: Dot product identity

Next, connect Figure 2.2 (a) and Figure 2.2 (b) by noting $a = |\mathbf{v}_2|$ and $b = |\mathbf{v}_1|$ and $c = |\mathbf{v}_2 - \mathbf{v}_1|$. Then

$$\begin{aligned}
 a^2 + b^2 - 2|\mathbf{v}_1||\mathbf{v}_2|\cos\theta &= a^2 + b^2 - 2ab \cos(\theta) = c^2 = |\mathbf{v}_2 - \mathbf{v}_1|^2 \\
 &= \left(\sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \right)^2 = (x_2 - x_1)^2 + (y_2 - y_1)^2 \\
 &= x_1^2 + y_1^2 - 2(x_1x_2 + y_1y_2) + x_2^2 + y_2^2 = a^2 + b^2 - 2(\mathbf{x}_1\mathbf{x}_2 + \mathbf{y}_1\mathbf{y}_2)
 \end{aligned}$$



Chapter 1

Math for Data

Dr. S. M.
Moosavi

Data Sets

Solutions

Exercise 1.4.

If one of u or v is a zero vector then the result is obvious. Otherwise,

$$\begin{cases} -1 \leq \cos \theta \leq 1 \\ \cos \theta = \frac{u \cdot v}{|u||v|} \end{cases} \Rightarrow -1 \leq \frac{u \cdot v}{|u||v|} \leq 1 \Rightarrow -|u||v| \leq u \cdot v \leq |u||v|.$$

Exercise 1.5.

For the homogeneous system (1.1) we saw that if $\det(A) = 0$ then $(x, y) = (-b, a)$ was a solution. If $\det(A) \neq 0$ the result comes from:

$$\begin{cases} d(ax + by) = 0 \\ b(cx + dy) = 0 \end{cases} \xrightarrow{\text{subtract}} (ad - bc)x = d(ax + by) - b(cx + dy) = 0.$$

and

$$\begin{cases} c(ax + by) = 0 \\ a(cx + dy) = 0 \end{cases} \xrightarrow{\text{subtract}} (bc - ad)y = c(ax + by) - a(cx + dy) = 0.$$

For the inhomogeneous system (1.3) use the same trick.



Chapter 1

Math for Data

Dr. S. M.
Moosavi

Data Sets

Solutions

Exercise 1.6.

By the inner product properties, if $\mathbf{a}, \mathbf{b} \in \mathbb{R}^d$ then:

$$\begin{aligned}
 |\mathbf{a} - \mathbf{b}|^2 &= (\mathbf{a} - \mathbf{b}) \cdot (\mathbf{a} - \mathbf{b}) \\
 &= (\mathbf{a} \cdot \mathbf{a}) - 2\mathbf{a} \cdot \mathbf{b} + (\mathbf{b} \cdot \mathbf{b}) \\
 &= |\mathbf{a}|^2 - 2\mathbf{a} \cdot \mathbf{b} + |\mathbf{b}|^2.
 \end{aligned}$$

Therefore, letting $\mathbf{a} = \mathbf{x}_k - \mathbf{m}$ and $\mathbf{b} = \mathbf{m} - \mathbf{x}$ we have:

$$\begin{aligned}
 MSD(\mathbf{x}) &= \frac{1}{N} \sum_{k=1}^N |\mathbf{x}_k - \mathbf{x}|^2 \\
 &= \frac{1}{N} \sum_{k=1}^N |\mathbf{x}_k - \mathbf{m} - (\mathbf{x} - \mathbf{m})|^2 \\
 &= \frac{1}{N} \sum_{k=1}^N |(\mathbf{x}_k - \mathbf{m}) - (\mathbf{x} - \mathbf{m})|^2 \\
 &= \frac{1}{N} \sum_{k=1}^N (|\mathbf{x}_k - \mathbf{m}|^2 - 2(\mathbf{x}_k - \mathbf{m}) \cdot (\mathbf{x} - \mathbf{m}) + |\mathbf{x} - \mathbf{m}|^2) \\
 &= \frac{1}{N} \sum_{k=1}^N |\mathbf{x}_k - \mathbf{m}|^2 - \frac{2}{N} \sum_{k=1}^N (\mathbf{x}_k - \mathbf{m}) \cdot (\mathbf{x} - \mathbf{m}) + \frac{1}{N} \sum_{k=1}^N |\mathbf{x} - \mathbf{m}|^2.
 \end{aligned}$$



Chapter 1

Math for Data

Dr. S. M.
Moosavi

Data Sets

Solutions

Now the middle term vanishes:

$$\begin{aligned}\frac{2}{N} \sum_{k=1}^N (\mathbf{x}_k - \mathbf{m}) \cdot (\mathbf{x} - \mathbf{m}) &= \frac{2}{N} (\mathbf{x} - \mathbf{m}) \cdot \sum_{k=1}^N (\mathbf{x}_k - \mathbf{m}) \\ &= \frac{2}{N} (\mathbf{x} - \mathbf{m}) \cdot \left(\sum_{k=1}^N \mathbf{x}_k - \sum_{k=1}^N \mathbf{m} \right) \\ &= \frac{2}{N} (\mathbf{x} - \mathbf{m}) \cdot (N\mathbf{m} - N\mathbf{m}) = 0.\end{aligned}$$

Hence,

$$\begin{aligned}MSD(\mathbf{x}) &= \frac{1}{N} \sum_{k=1}^N |\mathbf{x}_k - \mathbf{m}|^2 + \frac{1}{N} \sum_{k=1}^N |\mathbf{x} - \mathbf{m}|^2 \\ &= MSD(\mathbf{m}) + |\mathbf{x} - \mathbf{m}|^2 \\ &\geq MSD(\mathbf{m}).\end{aligned}$$