
Weight at age in the fishery
SC_Chile_PS
N_Chile
Offshore_Trawl
Weight
Years
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1970
1980
1990
2000
2010
2020
1
2
3
4
5
6
7
8
9
10
11
12


Weight at age in the survey
DEPM
Offshore_CPUE
Chile_AcousCS
Chile_AcousN
Chile_CPUE
Weight
Years
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1970
1980
1990
2000
2010
2020
1
2
3
4
5
6
7
8
9
10
11
12


Weight at age by cohort in the fleet
SC_Chile_PS
N_Chile
Offshore_Trawl
Weight
Age
0.0
0.5
1.0
1.5
2
4
6
8
10
12
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022


Weight at age by cohort in the survey
DEPM
Offshore_CPUE
Chile_AcousCS
Chile_AcousN
Chile_CPUE
Weight
Age
0.0
0.5
1.0
1.5
2
4
6
8
10
12
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022

Age composition in fleets




Age composition in fleets
N_Chile
Offshore_Trawl
SC_Chile_PS
2020
2010
2000
1990
1980
2
4
6
8
10
12
1.0
0.8
0.6
0.4
0.2
0.0
age


Age composition in fleets N_Chile
Proportion at age
Age
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
using pack
'fill' must n
using pack
'fill' must n

Age composition in fleets SC_Chile_PS




Age composition in fleets Offshore_Trawl
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
0.0
0.2
0.4
0.6
0.8
2 4 6 8 10
Proportion at age
Age

Age composition in surveys




Age composition in surveys
DEPM
2
4
6
8
10
12
2020
2010
2000
1990
1980
0.40
0.35
0.30
0.25
0.20
0.15
0.10
0.05
0.00
age


Weight in the stock
Weight (kg)
0.5
1.0
1.5
2
4
6
8
10
12
Age


Maturity in the stock
Proportion mature
1.0
0.9
0.8
0.7
0.6
0.5
2
4
6
8
10
12
Age


Total catch
Catch in kt
4000
3000
2000
1000
1970
1980
1990
2000
2010
2020
Years


Total catch by fleet
Catch by fleet in kt
4000
3000
2000
1000
1970
1980
1990
2000
2010
2020
Years
N_Chile
SC_Chile_PS
Offshore_Trawl


Catch residuals by fleet
−0.01
0
0
0.01
0.02
0.02
SC_Chile_PS
Offshore_Trawl
N_Chile
Residuals
1.4
1.2
1.0
0.8
0.6
1970
1980
1990
2000
2010
2020
Years


SC_Chile_PS
N_Chile
Offshore_Trawl
model − obs
year
1970
1980
1990
2000
2010
2020


Absolute residual catch by fleet
−0.16
−0.08
−0.01
0.06
0.13
0.2
SC_Chile_PS
N_Chile
Offshore_Trawl
Absolute residual catch
Years
12
10
8
6
4
2
1980
1990
2000
2010
2020


Age fits N_Chile
Observed
Predicted
Proportion at age
Age
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023


Age fits SC_Chile_PS
Observed
Predicted
Proportion at age
Age
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023


Age fits Offshore_Trawl
Observed
Predicted
2015
2016
2017
2018
2019
2020
2021
2022
2023
Proportion at age
Age
0.0
0.2
0.4
0.6
0.8
2
4
6
8
10
12


Predicted and observed catches by fleet
Observed
Predicted
SC_Chile_PS
N_Chile
Offshore_Trawl
Thousand tonnes
0
1000
2000
3000
4000
1970
1980
1990
2000
2010
2020
Years


Predicted and observed indices
Observed
Predicted
DEPM
Offshore_CPUE
Chile_AcousCS
Chile_AcousN
Chile_CPUE
Normalized index value
Years
1970
1980
1990
2000
2010
2020


Age fits Chile_AcousCS
Observed
Predicted
2001
2002
2003
2004
2005
2006
2007
2008
2009
Proportion at age
Age
0.0
0.1
0.2
0.3
0.4
0.5
2
4
6
8
10
12


Age fits Chile_AcousN
Observed
Predicted
2006
2007
2009
2013
2014
2015
2016
2017
2018
2019
2020
2021
Proportion at age
Age
0.0
0.2
0.4
0.6
0.8
2
4
6
8
10
12


Age fits DEPM
Observed
Predicted
2001
2003
2004
2005
2006
2008
Proportion at age
Age
0.0
0.1
0.2
0.3
2
4
6
8
10
12


Standardized survey residuals
−6.87
−4.57
−2.26
0.04
2.34
4.65
Offshore_CPUE
DEPM
Chile_CPUE
Chile_AcousN
Chile_AcousCS
Log residuals
1970
1980
1990
2000
2010
2020
Years


SD per input series
SD_age_N_Chile
SD_age_Offshore_Trawl
SD_age_SC_Chile_PS
SD_age_Chile_AcousCS
SD_age_Chile_AcousN
SD_age_DEPM
SD
1.0
0.5
1980
1990
2000
2010
2020
Years


Selectivity of the Fishery by Pentad
N_Chile
shore_Tr
C_Chile_
1970's
1975's
1980's
1985's
1990's
1995's
2000's
2005's
2010's
2015's
2020's
2025's
Selectivity
Age


Selectivity of the survey by Pentad
Selectivity
Age
1970's
1975's
1980's
1985's
1990's
1995's
2000's
2005's
2010's
2015's
2020's
2025's
DEPM

F at age


Years
2023
2022
2021
2020
2019
2018
2017
2016
2015
2014
2013
2012
2011
2010
2009
2008
2007
2006
2005
2004
2003
2002
2001
2000
1999
1998
1997
1996
1995
1994
1993
1992
1991
1990
1989
1988
1987
1986
1985
1984
1983
1982
1981
1980
1979
1978
1977
1976
1975
1974
1973
1972
1971
1970
1 2 3 4 5 6 7 8 9 10 11 12
Age
1.6
1.4
1.2
1.0
0.8
0.6
0.4
0.2
0.0


F proportion at age
Proportion of F at age
0.2
0.4
0.6
0.8
1970
1980
1990
2000
2010
2020
Years
1
2
3
4
5
6
7
8
9
10
11
12


N at age
Years
50
40
30
20
10
2
4
6
8
10
12
Age
50000
40000
30000
20000
10000
0


Proportion at age
Proportion of N at age
0.2
0.4
0.6
0.8
1970
1980
1990
2000
2010
2020
Years
1
2
3
4
5
6
7
8
9
10
11
12


Fishery mean age
Observed
Modelled
SC_Chile_PS
N_Chile
Offshore_Trawl
Age
Years
0
2
4
6
8
10
12
1980
1990
2000
2010
2020


Survey mean age
Observed
Modelled
DEPM
Chile_AcousCS
Chile_AcousN
Age
Years
10
8
6
4
2
2005
2010
2015
2020


SSB
Fishing mortality
Recruitment
Landings
data
year
25000
20000
15000
10000
5000
0
1.2
1.0
0.8
0.6
0.4
0.2
0.0
80000
60000
40000
20000
0
5000
4000
3000
2000
1000
0
1970
1980
1990
2000
2010
2020


Total biomass
30000
20000
10000
Fishing mortality
1.0
0.8
0.6
0.4
0.2
Recruitment
60000
40000
20000
Unfished biomass
60000
50000
40000
30000
20000
10000
data
1980
1990
2000
2010
2020
year


Uncertainty of key parameters
CV
0.1
0.2
0.3
0.4
1960
1980
2000
2020
years
R
SSB
TSB


Mature − Immature fish
Mature
Immature
Biomass in kt
25000
20000
15000
10000
5000
0
1970
1980
1990
2000
2010
2020
Years


Stock Recruitment
Simulated
2000 − 2019
Recruitment
Spawning Stock Biomass
0
10000
20000
30000
40000
50000
0
5000
10000
15000
20000
25000
1986
1987
1983
1985
1984
1992
1991
1988
1990
1989
2018
2000
2001
2002
1999
1997
1982
1981
2021
1994
1993
1995
2022
2016
2015
2014
2008
2006
2007
1975
1974
1973
1972
1971
1970
1969
1968
1967
1966
1965
1964
1963
1962
1961
1960
1950


Comparing fished with unfished biomass
Fished
Unfished
60000
50000
40000
30000
20000
10000
Total biomass
1970
1980
1990
2000
2010
2020
Years


SSB prediction
F2023 SQ
F2023 0.75x
F2023 1.25x
FMSY
F2023 0x
Spawning Stock Biomass
20000
15000
10000
5000
0
1960
1980
2000
2020
2040
Years


Catch prediction
F2023 SQ
F2023 0.75x
F2023 1.25x
FMSY
F2023 0x
Catch
4000
3000
2000
1000
0
1980
2000
2020
2040
Years


Yield and spawing stock biomass per recruit
YPR
SpawPR
Spawing biomass / Yield per recruit
Fishing mortality
2500
2000
1500
1000
500
0
2.0
1.5
1.0
0.5
0.0
0.0
0.2
0.4
0.6
0.8
1.0

Kobe plot




Deviation
1
0
−1
Simulated
2000 − 2019


Regime1
mean = −0.257
std = 0.651
ssq = 9.379
ssqT = 9.379
Density
0.6
0.4
0.2
−2
−1
0
1
Deviation


Weight at age in the fishery
FarNorth
Weight
0
1
2
3
1970
1980
1990
2000
2010
2020
Years
1
2
3
4
5
6
7
8
9
10
11
12


Weight at age in the survey
Peru_Acoustic
Peru_CPUE
Weight
0
1
2
3
1970
1980
1990
2000
2010
2020
Years
1
2
3
4
5
6
7
8
9
10
11
12


Weight at age by cohort in the fleet
FarNorth
Weight
Age
0
1
2
3
2
4
6
8
10
12
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022


Weight at age by cohort in the survey
Peru_Acoustic
Peru_CPUE
Weight
Age
0
1
2
3
2
4
6
8
10
12
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022


Weight in the stock
Weight (kg)
0.0
0.5
1.0
1.5
2.0
2.5
2
4
6
8
10
12
Age


Maturity in the stock
Proportion mature
1.0
0.8
0.6
0.4
0.2
0.0
2
4
6
8
10
12
Age

Length composition in fleet FarNorth



Length composition in fleets FarNorth




Total catch
Catch in kt
800
600
400
200
1970
1980
1990
2000
2010
2020
Years


Catch residuals by fleet
−0.02
−0.01
0
0.01
0.02
0.03
FarNorth
1.4
1.2
1.0
0.8
0.6
Residuals
1970
1980
1990
2000
2010
2020
Years


FarNorth
model − obs
year
1970
1980
1990
2000
2010
2020


Residuals catch−at−length by fleet
−0.11
−0.03
0.05
0.14
0.22
0.3
FarNorth
Length
Years
1980
1990
2000
2010
2020
10
20
30
40
50


Length fits FarNorth
Observed
Predicted
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
10 20 30 40 50
0.0 0.1 0.2 0.3 0.4 0.5
Proportion at length
Length


Predicted and observed catches by fleet
Observed
Predicted
FarNorth
Thousand tonnes
0
200
400
600
800
1970
1980
1990
2000
2010
2020
Years


Predicted and observed indices
Observed
Predicted
Peru_Acoustic
Peru_CPUE
Normalized index value
Years
1970
1980
1990
2000
2010
2020


Standardized survey residuals
−2.68
−1.63
−0.58
0.47
1.52
2.58
Peru_CPUE
Peru_Acoustic
Log residuals
1970
1980
1990
2000
2010
2020
Years


SD per input series
SD_length_NA
SD
3.0
2.5
2.0
1.5
1.0
0.5
1980
1990
2000
2010
2020
Years


Selectivity of the Fishery by Pentad
FarNorth 1970's
FarNorth 1975's
FarNorth 1980's
FarNorth 1985's
FarNorth 1990's
FarNorth 1995's
FarNorth 2000's
FarNorth 2005's
FarNorth 2010's
FarNorth 2015's
FarNorth 2020's
FarNorth 2025's
Selectivity
Age
0
1
2
3
4
5
6
2 4 6 8 12 2 4 6 8 12


Selectivity of the survey by Pentad
ru_Acous
eru_CPU
1970's
1975's
1980's
1985's
1990's
1995's
2000's
2005's
2010's
2015's
2020's
2025's
Selectivity
Age

F at age


Years
2023
2022
2021
2020
2019
2018
2017
2016
2015
2014
2013
2012
2011
2010
2009
2008
2007
2006
2005
2004
2003
2002
2001
2000
1999
1998
1997
1996
1995
1994
1993
1992
1991
1990
1989
1988
1987
1986
1985
1984
1983
1982
1981
1980
1979
1978
1977
1976
1975
1974
1973
1972
1971
1970
1 2 3 4 5 6 7 8 9 10 11 12
Age
8
6
4
2
0


F proportion at age
Proportion of F at age
0.8
0.6
0.4
0.2
1970
1980
1990
2000
2010
2020
Years
1
2
3
4
5
6
7
8
9
10
11
12


N at age
Years
50
40
30
20
10
2
4
6
8
10
12
Age
5000
4000
3000
2000
1000
0


Proportion at age
Proportion of N at age
0.8
0.6
0.4
0.2
1970
1980
1990
2000
2010
2020
Years
1
2
3
4
5
6
7
8
9
10
11
12

Fishery mean length
Observed
Modelled
FarNorth
Length (cm)
40
35
30
25
1980
1990
2000
2010
2020
Years


SSB
Fishing mortality
Recruitment
Landings
data
year
6000
5000
4000
3000
2000
1000
0
2.0
1.5
1.0
0.5
0.0
8000
6000
4000
2000
0
1000
800
600
400
200
0
1970
1980
1990
2000
2010
2020


Total biomass
6000
5000
4000
3000
2000
1000
Fishing mortality
1.5
1.0
0.5
Recruitment
6000
4000
2000
Unfished biomass
6000
4000
2000
data
1980
1990
2000
2010
2020
year


Uncertainty of key parameters
CV
0.1
0.2
0.3
0.4
1960
1980
2000
2020
years
R
SSB
TSB


Mature − Immature fish
Mature
Immature
4000
3000
2000
1000
0
Biomass in kt
1970
1980
1990
2000
2010
2020
Years

Stock Recruitment
Simulated
1970 − 1996
2001 − 2016
Recruitment
Spawning Stock Biomass
0
1000
2000
3000
4000
5000
1999
1987
1995
2017
2018
1977
1986
1988
1982
2009
1997
1981
1996
1978
2000
2004
1998
1992
1994
1989
1979
2016
1985
1991
2005
2001
1983
1980
2019
2002
1990
2010
2006
1984
2020
2021
2022
2007


Comparing fished with unfished biomass
Fished
Unfished
Total biomass
6000
5000
4000
3000
2000
1000
1970
1980
1990
2000
2010
2020
Years


SSB prediction
F2023 SQ
F2023 0.75x
F2023 1.25x
FMSY
F2023 0x
Spawning Stock Biomass
3000
2000
1000
0
1960
1980
2000
2020
2040
Years


Catch prediction
F2023 SQ
F2023 0.75x
F2023 1.25x
FMSY
F2023 0x
Catch
0
200
400
600
800
1980
2000
2020
2040
Years


Yield and spawing stock biomass per recruit
YPR
SpawPR
Spawing biomass / Yield per recruit
Fishing mortality
2500
2000
1500
1000
500
0
2.0
1.5
1.0
0.5
0.0
0.0
0.2
0.4
0.6
0.8
1.0


Kobe plot
100
50
0
−50
F/F_msy
0.5
1.0
1.5
2.0
2.5
B/B_msy
1970


Deviation
1
0
−1
−2
Simulated
1970 − 1996
2001 − 2016


Regime1
Regime2
mean = −0
std = 0.3
ssq = 3.2
ssqT = 17
mean = −0.
std = 0.8
ssq = 13.9
ssqT = 17
Density
Deviation
1.5
1.0
0.5
−2
−1
0
1