
CJM Diagnostics − catch FarNorth, age 1
a) Observed and fitted values time series
Observed
Fitted
values
5e−01
1e+01
5e+02
0
10
20
30
40
50
Year
b) Observed vs fitted values
1:1 line
Observed values
5e−01
5e+00
5e+01
5e+02
Fitted values
c) Standardised residuals over time
Standardised Residuals
−0.10
0.00
0.10
Year
d) Tukey−Anscombe plot
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
−2
−1
0
1
2
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
−0.2
0.0
0.1
0.2
5
10
15
Lag (yrs)


CJM Diagnostics − catch FarNorth, age 2
a) Observed and fitted values time series
Observed
Fitted
values
5e−01
1e+01
5e+02
0
10
20
30
40
50
Year
b) Observed vs fitted values
1:1 line
Observed values
5e−01
1e+01
5e+02
5e−01
5e+00
5e+01
5e+02
5e+03
Fitted values
c) Standardised residuals over time
Standardised Residuals
−0.3
−0.1
0.1
0.3
0
10
20
30
40
50
Year
d) Tukey−Anscombe plot
Standardised Residuals
−0.3
−0.1
0.1
0.3
5e−01
5e+00
5e+01
5e+02
5e+03
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
−0.3
−0.1
0.1
0.3
−2
−1
0
1
2
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
−0.4
−0.2
0.0
0.2
5
10
15
Lag (yrs)


CJM Diagnostics − catch FarNorth, age 3
a) Observed and fitted values time series
Observed
Fitted
values [10^3]
1e+01
1e−01
1e−03
0
10
20
30
40
50
Year
b) Observed vs fitted values
1:1 line
Observed values [10^3]
1e+01
1e−01
1e−03
1e−03
1e−02
1e−01
1e+00
1e+01
Fitted values [10^3]
c) Standardised residuals over time
Standardised Residuals
0.2
0.1
0.0
−0.2
0
10
20
30
40
50
Year
d) Tukey−Anscombe plot
Standardised Residuals
0.2
0.1
0.0
−0.2
1e−03
1e−02
1e−01
1e+00
1e+01
Fitted values [10^3]
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
0.2
0.1
0.0
−0.2
−2
−1
0
1
2
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
0.2
0.0
−0.2
−0.4
5
10
15
Lag (yrs)


CJM Diagnostics − catch FarNorth, age 4
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


CJM Diagnostics − catch FarNorth, age 5
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


CJM Diagnostics − catch FarNorth, age 6
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


CJM Diagnostics − catch FarNorth, age 7
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


CJM Diagnostics − catch FarNorth, age 8
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


CJM Diagnostics − catch FarNorth, age 9
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


CJM Diagnostics − catch FarNorth, age 10
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


CJM Diagnostics − catch FarNorth, age 11
a) Observed and fitted values time series
Observed
Fitted
values
1e+01
1e−01
1e−03
0
10
20
30
40
50
Year
b) Observed vs fitted values
1:1 line
Observed values
1e+01
1e−01
1e−03
1e−03
1e−02
1e−01
1e+00
1e+01
1e+02
Fitted values
c) Standardised residuals over time
Standardised Residuals
0.3
0.1
−0.1
−0.3
0
10
20
30
40
50
Year
d) Tukey−Anscombe plot
Standardised Residuals
0.3
0.1
−0.1
−0.3
1e−03
1e−02
1e−01
1e+00
1e+01
1e+02
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
0.3
0.1
−0.1
−0.3
−2
−1
0
1
2
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
0.8
0.6
0.4
0.2
−0.2
5
10
15
Lag (yrs)


CJM Diagnostics − catch FarNorth, age 12
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


CJM Diagnostics − catch N_Chile, age 1
a) Observed and fitted values time series
Observed
Fitted
values [10^3]
Year
b) Observed vs fitted values
1:1 line
Observed values [10^3]
Fitted values [10^3]
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values [10^3]
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


CJM Diagnostics − catch N_Chile, age 2
a) Observed and fitted values time series
Observed
Fitted
values [10^3]
5e+00
1e−01
5e−03
0
10
20
30
40
50
Year
b) Observed vs fitted values
1:1 line
Observed values [10^3]
5e+00
1e−01
5e−03
5e−03
5e−02
5e−01
5e+00
Fitted values [10^3]
c) Standardised residuals over time
Standardised Residuals
0.2
0.1
0.0
−0.1
−0.2
0
10
20
30
40
50
Year
d) Tukey−Anscombe plot
Standardised Residuals
0.2
0.1
0.0
−0.1
−0.2
5e−03
5e−02
5e−01
5e+00
Fitted values [10^3]
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
0.2
0.1
0.0
−0.1
−0.2
−2
−1
0
1
2
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
0.2
0.1
0.0
−0.2
5
10
15
Lag (yrs)


CJM Diagnostics − catch N_Chile, age 3
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


CJM Diagnostics − catch N_Chile, age 4
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


CJM Diagnostics − catch N_Chile, age 5
a) Observed and fitted values time series
Observed
Fitted
values
5e−01
1e+01
5e+02
0
10
20
30
40
50
Year
b) Observed vs fitted values
1:1 line
Observed values
5e−01
1e+01
5e+02
5e−01
5e+00
5e+01
5e+02
5e+03
Fitted values
c) Standardised residuals over time
Standardised Residuals
−0.10
0.00
0.10
0
10
20
30
40
50
Year
d) Tukey−Anscombe plot
Standardised Residuals
−0.10
0.00
0.10
5e−01
5e+00
5e+01
5e+02
5e+03
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
−0.10
0.00
0.10
−2
−1
0
1
2
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
−0.4
−0.2
0.0
0.2
5
10
15
Lag (yrs)


CJM Diagnostics − catch N_Chile, age 6
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


CJM Diagnostics − catch N_Chile, age 7
a) Observed and fitted values time series
Observed
Fitted
values
5e+02
1e+01
5e−01
0
10
20
30
40
50
Year
b) Observed vs fitted values
1:1 line
Observed values
5e+02
1e+01
5e−01
5e−01
5e+00
5e+01
5e+02
Fitted values
c) Standardised residuals over time
Standardised Residuals
0.05
0.00
−0.05
0
10
20
30
40
50
Year
d) Tukey−Anscombe plot
Standardised Residuals
0.05
0.00
−0.05
5e−01
5e+00
5e+01
5e+02
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
0.05
0.00
−0.05
−2
−1
0
1
2
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
0.2
0.1
−0.1
−0.3
5
10
15
Lag (yrs)


CJM Diagnostics − catch N_Chile, age 8
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


CJM Diagnostics − catch N_Chile, age 9
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


CJM Diagnostics − catch N_Chile, age 10
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


CJM Diagnostics − catch N_Chile, age 11
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


CJM Diagnostics − catch N_Chile, age 12
a) Observed and fitted values time series
Observed
Fitted
values
1e−03
1e−01
1e+01
0
10
20
30
40
50
Year
b) Observed vs fitted values
1:1 line
Observed values
1e−03
1e−01
1e+01
1e−03
1e−02
1e−01
1e+00
1e+01
1e+02
Fitted values
c) Standardised residuals over time
Standardised Residuals
−0.2
0.0
0.1
0.2
0
10
20
30
40
50
Year
d) Tukey−Anscombe plot
Standardised Residuals
−0.2
0.0
0.1
0.2
1e−03
1e−02
1e−01
1e+00
1e+01
1e+02
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
−0.2
0.0
0.1
0.2
−2
−1
0
1
2
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
−0.2
0.0
0.2
0.4
0.6
5
10
15
Lag (yrs)


CJM Diagnostics − catch Offshore_Trawl, age 1
a) Observed and fitted values time series
Observed
Fitted
values [10^3]
1e−04
1e−02
1e+00
0
10
20
30
40
50
Year
b) Observed vs fitted values
1:1 line
Observed values [10^3]
1e−04
1e−03
1e−02
1e−01
1e+00
1e+01
Fitted values [10^3]
c) Standardised residuals over time
Standardised Residuals
−0.15
−0.05
0.05
0.15
Year
d) Tukey−Anscombe plot
Fitted values [10^3]
e) Normal Q−Q plot
qqline
1:1 line
−2
−1
0
1
2
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
−0.4
−0.2
0.0
0.2
5
10
15
Lag (yrs)


CJM Diagnostics − catch Offshore_Trawl, age 2
a) Observed and fitted values time series
Observed
Fitted
values [10^3]
1e+00
1e−02
1e−04
0
10
20
30
40
50
Year
b) Observed vs fitted values
1:1 line
Observed values [10^3]
1e+00
1e−02
1e−04
1e−04
1e−03
1e−02
1e−01
1e+00
1e+01
Fitted values [10^3]
c) Standardised residuals over time
Standardised Residuals
0.15
0.05
−0.05
−0.15
0
10
20
30
40
50
Year
d) Tukey−Anscombe plot
Standardised Residuals
0.15
0.05
−0.05
−0.15
1e−04
1e−03
1e−02
1e−01
1e+00
1e+01
Fitted values [10^3]
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
0.15
0.05
−0.05
−0.15
−2
−1
0
1
2
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
0.1
−0.1
−0.3
5
10
15
Lag (yrs)


CJM Diagnostics − catch Offshore_Trawl, age 3
a) Observed and fitted values time series
Observed
Fitted
values
1e−01
1e+01
1e+03
0
10
20
30
40
50
Year
b) Observed vs fitted values
1:1 line
Observed values
1e−01
1e+01
1e+03
1e−01
1e+00
1e+01
1e+02
1e+03
Fitted values
c) Standardised residuals over time
Standardised Residuals
−0.2
−0.1
0.0
0.1
0.2
0
10
20
30
40
50
Year
d) Tukey−Anscombe plot
Standardised Residuals
−0.2
−0.1
0.0
0.1
0.2
1e−01
1e+00
1e+01
1e+02
1e+03
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
−0.2
−0.1
0.0
0.1
0.2
−2
−1
0
1
2
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
−0.3
−0.1
0.1
5
10
15
Lag (yrs)


CJM Diagnostics − catch Offshore_Trawl, age 4
a) Observed and fitted values time series
Observed
Fitted
values
1e−01
1e+01
1e+03
0
10
20
30
40
50
Year
b) Observed vs fitted values
1:1 line
Observed values
1e−01
1e+01
1e+03
1e−01
1e+00
1e+01
1e+02
1e+03
Fitted values
c) Standardised residuals over time
Standardised Residuals
−0.10
0.00
0.05
0.10
0
10
20
30
40
50
Year
d) Tukey−Anscombe plot
Standardised Residuals
−0.10
0.00
0.05
0.10
1e−01
1e+00
1e+01
1e+02
1e+03
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
−0.10
0.00
0.05
0.10
−2
−1
0
1
2
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
−0.6
−0.4
−0.2
0.0
0.2
5
10
15
Lag (yrs)


CJM Diagnostics − catch Offshore_Trawl, age 5
a) Observed and fitted values time series
Observed
Fitted
values
1e−01
1e+01
1e+03
0
10
20
30
40
50
Year
b) Observed vs fitted values
1:1 line
Observed values
1e−01
1e+01
1e+03
1e−01
1e+00
1e+01
1e+02
1e+03
Fitted values
c) Standardised residuals over time
Standardised Residuals
−0.06
−0.02
0.02
0.06
0
10
20
30
40
50
Year
d) Tukey−Anscombe plot
Standardised Residuals
−0.06
−0.02
0.02
0.06
1e−01
1e+00
1e+01
1e+02
1e+03
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
−0.06
−0.02
0.02
0.06
−2
−1
0
1
2
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
−0.2
0.0
0.1
0.2
5
10
15
Lag (yrs)


CJM Diagnostics − catch Offshore_Trawl, age 6
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


CJM Diagnostics − catch Offshore_Trawl, age 7
a) Observed and fitted values time series
Observed
Fitted
values
1e+03
1e+01
1e−01
0
10
20
30
40
50
Year
b) Observed vs fitted values
1:1 line
Observed values
1e+03
1e+01
1e−01
1e−01
1e+00
1e+01
1e+02
1e+03
Fitted values
c) Standardised residuals over time
Standardised Residuals
0.04
0.00
−0.04
0
10
20
30
40
50
Year
d) Tukey−Anscombe plot
Standardised Residuals
0.04
0.00
−0.04
1e−01
1e+00
1e+01
1e+02
1e+03
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
0.04
0.00
−0.04
−2
−1
0
1
2
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
0.1
−0.1
−0.3
5
10
15
Lag (yrs)


CJM Diagnostics − catch Offshore_Trawl, age 8
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


CJM Diagnostics − catch Offshore_Trawl, age 9
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


CJM Diagnostics − catch Offshore_Trawl, age 10
a) Observed and fitted values time series
Observed
Fitted
values
1e−03
1e−01
1e+01
0
10
20
30
40
50
Year
b) Observed vs fitted values
1:1 line
Observed values
1e−03
1e−01
1e+01
1e−03
1e−02
1e−01
1e+00
1e+01
1e+02
Fitted values
c) Standardised residuals over time
Standardised Residuals
−0.2
0.0
0.1
0.2
0
10
20
30
40
50
Year
d) Tukey−Anscombe plot
Standardised Residuals
−0.2
0.0
0.1
0.2
1e−03
1e−02
1e−01
1e+00
1e+01
1e+02
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
−0.2
0.0
0.1
0.2
−2
−1
0
1
2
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
−0.2
0.0
0.2
0.4
0.6
5
10
15
Lag (yrs)


CJM Diagnostics − catch Offshore_Trawl, age 11
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


CJM Diagnostics − catch Offshore_Trawl, age 12
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


CJM Diagnostics − catch SC_Chile_PS, age 1
a) Observed and fitted values time series
Observed
Fitted
values [10^3]
Year
b) Observed vs fitted values
1:1 line
Observed values [10^3]
Fitted values [10^3]
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values [10^3]
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


CJM Diagnostics − catch SC_Chile_PS, age 2
a) Observed and fitted values time series
Observed
Fitted
values [10^3]
5e−03
1e−01
5e+00
0
10
20
30
40
50
Year
b) Observed vs fitted values
1:1 line
Observed values [10^3]
5e−03
1e−01
5e+00
5e−03
5e−02
5e−01
5e+00
Fitted values [10^3]
c) Standardised residuals over time
Standardised Residuals
−0.15
−0.05
0.05
0.15
0
10
20
30
40
50
Year
d) Tukey−Anscombe plot
Standardised Residuals
−0.15
−0.05
0.05
0.15
5e−03
5e−02
5e−01
5e+00
Fitted values [10^3]
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
−0.15
−0.05
0.05
0.15
−2
−1
0
1
2
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
−0.3
−0.1
0.1
5
10
15
Lag (yrs)


CJM Diagnostics − catch SC_Chile_PS, age 3
a) Observed and fitted values time series
Observed
Fitted
values [10^3]
Year
b) Observed vs fitted values
1:1 line
Observed values [10^3]
Fitted values [10^3]
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values [10^3]
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


CJM Diagnostics − catch SC_Chile_PS, age 4
a) Observed and fitted values time series
Observed
Fitted
values [10^3]
5e−03
1e−01
5e+00
0
10
20
30
40
50
Year
b) Observed vs fitted values
1:1 line
Observed values [10^3]
5e−03
5e−02
5e−01
5e+00
Fitted values [10^3]
c) Standardised residuals over time
Standardised Residuals
−0.15
−0.05
0.05
0.15
Year
d) Tukey−Anscombe plot
Fitted values [10^3]
e) Normal Q−Q plot
qqline
1:1 line
Quantiles of the Normal Distribution
−2
−1
0
1
2
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
−0.3
−0.1
0.1
5
10
15
Lag (yrs)


CJM Diagnostics − catch SC_Chile_PS, age 5
a) Observed and fitted values time series
Observed
Fitted
values [10^3]
5e−03
1e−01
5e+00
0
10
20
30
40
50
Year
b) Observed vs fitted values
1:1 line
Observed values [10^3]
5e−03
1e−01
5e+00
5e−03
5e−02
5e−01
5e+00
Fitted values [10^3]
c) Standardised residuals over time
Standardised Residuals
−0.15
−0.05
0.05
0.15
0
10
20
30
40
50
Year
d) Tukey−Anscombe plot
Standardised Residuals
−0.15
−0.05
0.05
0.15
5e−03
5e−02
5e−01
5e+00
Fitted values [10^3]
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
−0.15
−0.05
0.05
0.15
−2
−1
0
1
2
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
−0.3
−0.1
0.1
5
10
15
Lag (yrs)


CJM Diagnostics − catch SC_Chile_PS, age 6
a) Observed and fitted values time series
Observed
Fitted
values [10^3]
5e−03
1e−01
5e+00
0
10
20
30
40
50
Year
b) Observed vs fitted values
1:1 line
Observed values [10^3]
5e−03
5e−02
5e−01
5e+00
Fitted values [10^3]
c) Standardised residuals over time
Standardised Residuals
−0.10
0.00
0.10
Year
d) Tukey−Anscombe plot
Fitted values [10^3]
e) Normal Q−Q plot
qqline
1:1 line
−2
−1
0
1
2
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
−0.3
−0.1
0.1
5
10
15
Lag (yrs)


CJM Diagnostics − catch SC_Chile_PS, age 7
a) Observed and fitted values time series
Observed
Fitted
values [10^3]
1e+01
1e−01
1e−03
0
10
20
30
40
50
Year
b) Observed vs fitted values
1:1 line
Observed values [10^3]
1e+01
1e−01
1e−03
1e−03
1e−02
1e−01
1e+00
1e+01
Fitted values [10^3]
c) Standardised residuals over time
Standardised Residuals
0.10
0.00
−0.10
0
10
20
30
40
50
Year
d) Tukey−Anscombe plot
Standardised Residuals
0.10
0.00
−0.10
1e−03
1e−02
1e−01
1e+00
1e+01
Fitted values [10^3]
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
0.10
0.00
−0.10
−2
−1
0
1
2
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
0.2
0.1
0.0
−0.2
5
10
15
Lag (yrs)


CJM Diagnostics − catch SC_Chile_PS, age 8
a) Observed and fitted values time series
Observed
Fitted
values [10^3]
1e+01
1e−01
1e−03
0
10
20
30
40
50
Year
b) Observed vs fitted values
1:1 line
Observed values [10^3]
1e+01
1e−01
1e−03
1e−03
1e−02
1e−01
1e+00
1e+01
Fitted values [10^3]
c) Standardised residuals over time
Standardised Residuals
0.10
0.00
−0.10
0
10
20
30
40
50
Year
d) Tukey−Anscombe plot
Standardised Residuals
0.10
0.00
−0.10
1e−03
1e−02
1e−01
1e+00
1e+01
Fitted values [10^3]
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
0.10
0.00
−0.10
−2
−1
0
1
2
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
0.4
0.2
0.0
−0.2
5
10
15
Lag (yrs)


CJM Diagnostics − catch SC_Chile_PS, age 9
a) Observed and fitted values time series
Observed
Fitted
values
5e−01
1e+01
5e+02
0
10
20
30
40
50
Year
b) Observed vs fitted values
1:1 line
Observed values
5e−01
1e+01
5e+02
5e−01
5e+00
5e+01
5e+02
Fitted values
c) Standardised residuals over time
Standardised Residuals
−0.6
−0.2
0.2
0.6
0
10
20
30
40
50
Year
d) Tukey−Anscombe plot
Standardised Residuals
−0.6
−0.2
0.2
0.6
5e−01
5e+00
5e+01
5e+02
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
−0.6
−0.2
0.2
0.6
−2
−1
0
1
2
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
−0.2
0.2
0.4
0.6
0.8
5
10
15
Lag (yrs)


CJM Diagnostics − catch SC_Chile_PS, age 10
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


CJM Diagnostics − catch SC_Chile_PS, age 11
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


CJM Diagnostics − catch SC_Chile_PS, age 12
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


CJM Diagnostics − Chile_AcousN, age −1
a) Observed and fitted values time series
Observed
Fitted
values [10^3]
1e+01
1e−01
1e−03
30
35
40
45
50
55
Year
b) Observed vs fitted values
1:1 line
Observed values [10^3]
1e+01
1e−01
1e−03
1e−03
1e−02
1e−01
1e+00
1e+01
Fitted values [10^3]
c) Standardised residuals over time
Standardised Residuals
5
0
−5
30
35
40
45
50
55
Year
d) Tukey−Anscombe plot
Standardised Residuals
5
0
−5
1e−03
1e−02
1e−01
1e+00
1e+01
Fitted values [10^3]
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
5
0
−5
−2
−1
0
1
2
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
0.4
0.2
0.0
−0.4
2
4
6
8
10
12
14
Lag (yrs)


CJM Diagnostics − Chile_CPUE_early, age −1
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


CJM Diagnostics − Chile_CPUE_late, age −1
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


CJM Diagnostics − Offshore_CPUE_early, age −1
a) Observed and fitted values time series
Observed
Fitted
values [10^3]
Year
b) Observed vs fitted values
1:1 line
Observed values [10^3]
Fitted values [10^3]
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values [10^3]
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


CJM Diagnostics − Offshore_CPUE_late, age −1
a) Observed and fitted values time series
Observed
Fitted
values [10^3]
Year
b) Observed vs fitted values
1:1 line
Observed values [10^3]
Fitted values [10^3]
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values [10^3]
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


CJM Diagnostics − Peru_CPUE, age −1
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)