Chilli (*Capsicum annuum* L.)
Solanaceae

Field preparation

Thoroughly prepare the field with the addition of FYM @ 25 t/ ha and form ridges and furrows at a spacing of 60 cm. Apply 2 kg/ha of Azospirillum and 2 kg / ha of Phosphobacteria by mixing with 20 kg of FYM. Irrigate the furrows and transplant 40-45 days old seedlings, with the ball of earth on the ridges.

Spacing

Varieties : 60 x 45 cm
Hybrids : 75 x 60 cm

Weed control

Apply Pendimethalin 1.0 kg *a.i.* / ha or Fluchloralin 1.0 kg *a.i.* / ha as pre-emergece herbicide followed by hand weeding once 30 days after planting.



Chilli spacing

Irrigation
Irrigate is done at weekly intervals.

Layout and planting for drip irrigation and fertigation

- Apply FYM @ 25 t / ha as basal before last ploughing.
- Apply 2 kg / ha of Azospirillum and 2 kg/ha Phosphobacteria by mixing with 20 kg of FYM.
- Apply 75 % total recommended dose of superphosphate i.e. 375 kg / ha as basal.
- Install the drip irrigation with main and sub main pipes and place lateral tubes at an interval of 1.5 m.
- Place the drippers in lateral tubes at an interval of 60 cm and 50 cm spacing with 4 LPH and 3.5 LPH capacities respectively.
- Form raised beds of 120 cm width at an interval of 30 cm and place the laterals at the centre of the each bed.
- Before planting wet the beds using drip system for 8-12 hrs.
- Planting to be done at a spacing of 90 x 60 x 45 cm in the paired row system, using ropes marked at 60 cm spacing.
- Spray Pendimethalin 1.0 kg *a.i.* / ha or Fluchloralin 1.0 kg *a.i* / ha as pre-emergence herbicide at 3rd day after planting.
- Gap filling to be done at 7th day after transplanting.

Updated on October, 2022

© All Rights Reserved. TNAU 2008 - 2022.