

Presentation PROJECT01

Pakala srijith reddy
first year student
ee19btech11041
Dept. of Electrical Engg.,
IIT Hyderabad.
Marpu mythri varshitha
first year student
ee19btech11014
Dept. of Electrical Engg.,
IIT Hyderabad.

August 21, 2019

Problem

- Solution
 - Centre and Radius
 - Points at which tangent meets circle
 - Equation of Tangents

3 Plot

Problem Statement

Draw the tangents through the point

$$\mathbf{P} = \begin{pmatrix} 4 \\ 7 \end{pmatrix}$$

(2.0)

to the circle

$$C: x^T x = 9. (2.0)$$

Finding the Centre and Radius

Let \mathbf{O} be the centre of C. From the above equation of circle given in question,

$$\mathbf{O} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \tag{3.0}$$

also radius of circle c is r $r=\sqrt{9}=3$.

Finding the tangent points

let us name points on circle where tangent drawn from P meets the circle as Q1,Q2 whose solution is let's say Q.since PQ Is perpendicular to QO $^{\circ}$

$$(P-Q)^{T}(Q-O)=0,$$
 (3.0)

$$(P - Q) = k(omat.Q) \tag{3.0}$$

where

$$omat = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \tag{3.0}$$

which implies

$$||P - Q||^2 = K^2 ||omat.Q||^2$$
 (3.0)

since Q is a point on circle C,

$$||Q|| = 3 \tag{3.0}$$

$$k^2 = ||P - Q||^2/9 \tag{3.0}$$

$$||P - Q|| = \sqrt{PO^2 - r^2} \tag{3.0}$$

or
$$k = -\sqrt{56}/3$$
 (3.0)
$$P = (I + (k.omat)).Q$$
 (3.0)

 $k = +\sqrt{56}/3$

$$I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

let's say (I+k.omat)=R then

$$Q = R^{-1}P$$

$$R = \begin{pmatrix} 1 & \sqrt{56/3} \\ -\sqrt{56/3} & 1 \end{pmatrix}$$

$$R = ($$

$$R = \left(\begin{array}{c} 1 \\ 1 \end{array} \right)$$

$$R = \begin{pmatrix} 1 & -\sqrt{56}/3 \\ \sqrt{56}/3 & 1 \end{pmatrix}$$

$$\begin{pmatrix} -\sqrt{56}/3 \\ 1 \end{pmatrix}$$

$$Q1 = \begin{pmatrix} 1 & -\sqrt{56}/3 \\ \sqrt{56}/3 & 1 \end{pmatrix} \begin{pmatrix} 4 \\ 7 \end{pmatrix} / (1 + 56/9)$$

therefore

(3.0)

(3.0)

(3.0)

(3.0)

(3.0)

$$Q1 = 9/65 \begin{pmatrix} 4 - 7\sqrt{56/3} \\ 7 + 4\sqrt{56/3} \end{pmatrix} \tag{3.0}$$

similarly,

$$Q2 = 9/65 \begin{pmatrix} 4 + 7\sqrt{56}/3 \\ 7 - 4\sqrt{56}/3 \end{pmatrix}$$
 (3.0)

Finding the equation of tangents

The equation of tangent can be obtained from direction vector

$$m_1 = (P - Q1) (3.0)$$

The normal vector n_1 is defined as

$$n_1^T m_1 = 0 (3.0)$$

and can be obtained as

$$n_1 = Q1 - O (3.0)$$

$$n_1 = \begin{pmatrix} (392 - 12\sqrt{56})/65\\ -(224 + 21\sqrt{56})/65 \end{pmatrix}$$
 (3.0)

Therefore equation of tangent PQ1 is

$$n_1^T(x - P) = 0 (3.0)$$

$$n_1^T x = n_1^T P \tag{3.0}$$

$$\begin{pmatrix} (392 - 12\sqrt{56})/65 \\ -(224 + 21\sqrt{56})/65 \end{pmatrix}^T x = \begin{pmatrix} (392 - 12\sqrt{56})/65 \\ -(224 + 21\sqrt{56})/65 \end{pmatrix}^T \begin{pmatrix} 4 \\ 7 \end{pmatrix}$$
 (3.0)

$$((392 - 12\sqrt{56}) - (224 + 21\sqrt{56})) x$$
 (3.0)

$$= ((392 - 12\sqrt{56}) - (224 + 21\sqrt{56})) {4 \choose 7}$$
 (3.0)

The equaion of tangent PQ1 is

$$((392 - 12\sqrt{56}) - (224 + 21\sqrt{56})) x = -195\sqrt{56}$$
 (3.0)

similarly,

The equation of tangent can be obtained from direction vector

$$m_2 = (P - Q2) (3.0)$$

The normal vector n_2 is defined as

$$n_2^T m_2 = 0 (3.0)$$

and can be obtained as

$$n_2 = Q2 - O \tag{3.0}$$

$$n_2 = \begin{pmatrix} (392 + 12\sqrt{56})/65 \\ -(224 - 21\sqrt{56})/65 \end{pmatrix}$$
 (3.0)

Therefore equation of tangent PQ2 is

$$n_2^T(x - P) = 0 (3.0)$$

$$n_2^T x = n_2^T P \tag{3.0}$$

$$\begin{pmatrix} (392 + 12\sqrt{56})/65 \\ -(224 - 21\sqrt{56})/65 \end{pmatrix}^T x = \begin{pmatrix} (392 + 12\sqrt{56})/65 \\ -(224 - 21\sqrt{56})/65 \end{pmatrix}^T \begin{pmatrix} 4 \\ 7 \end{pmatrix}$$
 (3.0)

$$((392 + 12\sqrt{56}) - (224 - 21\sqrt{56}))x$$
 (3.0)

$$= ((392 + 12\sqrt{56}) - (224 - 21\sqrt{56})) \begin{pmatrix} 4 \\ 7 \end{pmatrix}$$
 (3.0)

The equaion of tangent PQ2 is

$$((392 + 12\sqrt{56}) - (224 - 21\sqrt{56})) x = +195\sqrt{56}$$
 (3.0)

Plot

https://github.com/SRIJITH01/Srijith/blob/master/srijith.py

plots Fig. 1.




Figure: Tangents To circle.