RAJALAKSHMI ENGINEERING COLLEGE

Question 1 | Correct Mark 1.00 out of 1.00

Given an array `nums` of size `n`, return *the majority element*.

The majority element is the element that appears more than `⌊n / 2⌋` times. You may assume that the majority element always exists in the array.

Example 1:

```
Input: nums = [3,2,3]
Output: 3
```

Example 2:

```
Input: nums = [2,2,1,1,1,2,2]
Output: 2
```

Constraints:

- `n == nums.length`
- `1 <= n <= 5 * 10^4`
- `-2^31 <= nums[i] <= 2^31 - 1`

For example:

Input	Result
3 3 2 3	3
7 2 2 1 1 1 2 2	2

Answer: (penalty regime: 0 %)

```c
#include <stdio.h>
int main() {
    int n;
    scanf("%d",&n);
    int a[n];
    for(int i=0;i<n;i++) {
        scanf("%d",&a[i]);
    }
    for(int i=0;i<n;i++) {
        int count=0;
        for(int j=0;j<n;j++) {
            if(a[j]==a[i]) {
                count++;
            }
        }
        if(count>n/2){
            printf("%d\n",a[i]);
            break;
        }
    }
    return 0;
}
```

Check

	Input	Expected	Got	
✔	3 3 2 3	3	3	✔

Passed all tests! ✔

Correct

Marks for this submission: 1.00/1.00.

Back to Course