

Network Autocorrelation Models

SOC 280: Analysis of Social Network Data

Carter T. Butts

Department of Sociology and
Institute for Mathematical Behavioral Sciences
University of California, Irvine


From Indices to Autocorrelation

- **Last time, we discussed how properties of positions could be related to covariates....**
 - Does being a gatekeeper help your career?
 - Do high-degree organizations make all the decisions?
- **Today: how covariates relate to each other through ties**
 - *Network autocorrelation*: (usually linear) dependence of individual attributes across positions
- **Applicable to many phenomena**
 - E.g., Social influence, organizational isomorphism, work-family spillover, neighborhood quality, economic productivity, etc.


Network Autocorrelation Model

- **Simplest tool for modeling network autocorrelation: the (linear) network autocorrelation model**
 - Builds on standard (OLS) regression
 - Closely analogous to ARMA models in the time series literature
 - Identical to the spatial ARMA models of Cliff and Ord (1973), Anselin (1988), etc. (But we use a different name!)
- **Basic components**
 - Response variable (to be modeled)
 - “Weight” matrices (identical to or derived from original networks)
 - Predictive covariates
 - Parameters measuring nature/extent of dependence, covariate effects


The Classical Regression Model




Adding Network AR Effects




Adding Network MA Effects




Network ARMA Model




AR Vs. MA




AR Vs. MA




AR Vs. MA




Network “Resonance”




Network “Resonance”




Network “Resonance”



Network “Resonance”



Network “Resonance”



The Formal Development

Individual Case Form:

$$y_i = \sum_{j=1}^p X_{ij} \beta_j + \epsilon_i$$

Group (Matrix/Vector) Form:

$$y = X \beta + \epsilon$$

The Formal Development

Individual Case Form:

$$y_i = \sum_{j=1}^p X_{ij} \beta_j + \epsilon_i$$

Response → y_i ← Perturbation

Covariates → X_{ij}

Group (Matrix/Vector) Form:

$$\mathbf{y} = \mathbf{X} \boldsymbol{\beta} + \boldsymbol{\epsilon}$$

The Formal Development

Individual Case Form:

$$y_i = \sum_{j=1}^N \theta W_{ij} y_j + \sum_{j=1}^p X_{ij} \beta_j + \epsilon_i$$

Group (Matrix/Vector) Form:

$$\mathbf{y} = \theta \mathbf{W} \mathbf{y} + \mathbf{X} \boldsymbol{\beta} + \boldsymbol{\epsilon}$$

The Formal Development

Individual Case Form:

$$y_i = \sum_{j=1}^N \theta W_{ij} y_j + \sum_{j=1}^p X_{ij} \beta_j + \epsilon_i$$

Annotations for the Individual Case Form:

- AR Term: Points to the first term $\sum_{j=1}^N \theta W_{ij} y_j$.
- Response: Points to the variable y_i .
- Covariates: Points to the second term $\sum_{j=1}^p X_{ij} \beta_j$.
- Perturbation: Points to the error term ϵ_i .

Group (Matrix/Vector) Form:

$$\mathbf{y} = \boldsymbol{\theta} \mathbf{W} \mathbf{y} + \mathbf{X} \boldsymbol{\beta} + \boldsymbol{\epsilon}$$

The Formal Development

Individual Case Form:

$$y_i = \sum_{j=1}^N \theta_j W_{ij} y_j + \sum_{j=1}^p X_{ij} \beta_j + \epsilon_i$$

$$\epsilon_i = \sum_{j=1}^N \psi_j Z_{ij} \epsilon_j + \nu_i$$

Group (Matrix/Vector) Form:

$$\mathbf{y} = \theta \mathbf{W} \mathbf{y} + \mathbf{X} \boldsymbol{\beta} + \boldsymbol{\epsilon}$$

$$\boldsymbol{\epsilon} = \psi \mathbf{Z} \boldsymbol{\epsilon} + \boldsymbol{\nu}$$

The Formal Development

Individual Case Form:

$$y_i = \sum_{j=1}^N \theta W_{ij} y_j + \sum_{j=1}^p X_{ij} \beta_j + \epsilon_i$$

AR Term → y_i ← Response

Covariates ← X_{ij}

$$\epsilon_i = \sum_{j=1}^N \psi Z_{ij} \epsilon_j + \nu_i$$

MA Term ← Z_{ij}

Total Perturbation → ϵ_i ← Individual Perturbation

Group (Matrix/Vector) Form:

$$\mathbf{y} = \theta \mathbf{W} \mathbf{y} + \mathbf{X} \boldsymbol{\beta} + \boldsymbol{\epsilon}$$

$$\boldsymbol{\epsilon} = \psi \mathbf{Z} \boldsymbol{\epsilon} + \boldsymbol{\nu}$$

Resolving the Resonance: Total Perturbations

$$\epsilon = \psi Z \epsilon + \nu$$

Resolving the Resonance: Total Perturbations

$$\epsilon = \psi Z \epsilon + \nu$$

$$\epsilon - \psi Z \epsilon = \nu$$

Resolving the Resonance: Total Perturbations

$$\epsilon = \psi Z \epsilon + \nu$$

$$\epsilon - \psi Z \epsilon = \nu$$

$$(I - \psi Z) \epsilon = \nu$$

Resolving the Resonance: Total Perturbations

$$\epsilon = \psi Z \epsilon + \nu$$

$$\epsilon - \psi Z \epsilon = \nu$$

$$(I - \psi Z) \epsilon = \nu$$

$$\epsilon = (I - \psi Z)^{-1} \nu$$

Resolving the Resonance: Total Perturbations

$$\epsilon = \psi Z \epsilon + \nu$$

$$\epsilon - \psi Z \epsilon = \nu$$

$$(I - \psi Z) \epsilon = \nu$$

Total
Perturbations

Matrix of “Equilibrium
MA Weights”

Individual
Perturbations

$$\epsilon = (I - \psi Z)^{-1} \nu$$

Resolving the Resonance, II: Responses

$$\mathbf{y} = \theta \mathbf{W} \mathbf{y} + \mathbf{X} \boldsymbol{\beta} + \boldsymbol{\epsilon}$$

Resolving the Resonance, II: Responses

$$\mathbf{y} = \theta \mathbf{W} \mathbf{y} + \mathbf{X} \boldsymbol{\beta} + \boldsymbol{\epsilon}$$

$$\mathbf{y} - \theta \mathbf{W} \mathbf{y} = \mathbf{X} \boldsymbol{\beta} + \boldsymbol{\epsilon}$$

Resolving the Resonance, II: Responses

$$\mathbf{y} = \theta \mathbf{W} \mathbf{y} + \mathbf{X} \boldsymbol{\beta} + \boldsymbol{\epsilon}$$

$$\mathbf{y} - \theta \mathbf{W} \mathbf{y} = \mathbf{X} \boldsymbol{\beta} + \boldsymbol{\epsilon}$$

$$(\mathbf{I} - \theta \mathbf{W})\mathbf{y} = \mathbf{X} \boldsymbol{\beta} + \boldsymbol{\epsilon}$$

Resolving the Resonance, II: Responses

$$\mathbf{y} = \theta \mathbf{W} \mathbf{y} + \mathbf{X} \boldsymbol{\beta} + \boldsymbol{\epsilon}$$

$$\mathbf{y} - \theta \mathbf{W} \mathbf{y} = \mathbf{X} \boldsymbol{\beta} + \boldsymbol{\epsilon}$$

$$(\mathbf{I} - \theta \mathbf{W}) \mathbf{y} = \mathbf{X} \boldsymbol{\beta} + \boldsymbol{\epsilon}$$

$$\mathbf{y} = (\mathbf{I} - \theta \mathbf{W})^{-1} (\mathbf{X} \boldsymbol{\beta} + \boldsymbol{\epsilon})$$

Resolving the Resonance, II: Responses

$$y = \theta W y + X \beta + \epsilon$$

$$y - \theta W y = X \beta + \epsilon$$

$$(I - \theta W)y = X \beta + \epsilon$$

$$y = (I - \theta W)^{-1} (X \beta + \epsilon)$$

Responses → $y = (I - \theta W)^{-1} (X \beta + \epsilon)$

Matrix of “Equilibrium AR Weights” → $(I - \theta W)^{-1}$

Covariates → $(X \beta)$

Total Perturbations → ϵ

Resolving the Resonance, II: Responses

$$\mathbf{y} = \theta \mathbf{W} \mathbf{y} + \mathbf{X} \boldsymbol{\beta} + \boldsymbol{\epsilon}$$

$$\mathbf{y} - \theta \mathbf{W} \mathbf{y} = \mathbf{X} \boldsymbol{\beta} + \boldsymbol{\epsilon}$$

$$(\mathbf{I} - \theta \mathbf{W}) \mathbf{y} = \mathbf{X} \boldsymbol{\beta} + \boldsymbol{\epsilon}$$

$$\mathbf{y} = (\mathbf{I} - \theta \mathbf{W})^{-1} (\mathbf{X} \boldsymbol{\beta} + \boldsymbol{\epsilon})$$

$$\mathbf{y} = (\mathbf{I} - \theta \mathbf{W})^{-1} \left(\mathbf{X} \boldsymbol{\beta} + (\mathbf{I} - \psi \mathbf{Z})^{-1} \mathbf{v} \right)$$

Resolving the Resonance, II: Responses

$$y = (I - \theta W)^{-1} (X\beta + (I - \psi Z)^{-1} \nu)$$

Diagram illustrating the components of the equation:

- Responses** (green) points to the leftmost term y .
- Matrix of “Equilibrium AR Weights”** (teal) points to the term $(I - \theta W)^{-1}$.
- Covariates** (blue) points to the term $X\beta$.
- Matrix of “Equilibrium MA Weights”** (purple) points to the term $(I - \psi Z)^{-1}$.
- Individual Perturbations** (yellow) points to the rightmost term ν .

Using Multiple Weight Matrices

$$\theta \text{ W} \rightarrow \sum_{i=1}^w \theta_i W_i$$

$$\psi \text{ Z} \rightarrow \sum_{i=1}^z \psi_i Z_i$$

Using Multiple Weight Matrices

$$\theta \text{ W} \rightarrow \sum_{i=1}^w \theta_i \text{ W}_i$$

$$\psi \text{ Z} \rightarrow \sum_{i=1}^z \psi_i \text{ Z}_i$$

$$y = \left(I - \sum_{i=1}^w \theta_i \text{ W}_i \right)^{-1} \left(X \beta + \left(I - \sum_{i=1}^z \psi_i \text{ Z}_i \right)^{-1} v \right)$$

Inference with the Network Autocorrelation Model

- **In practice, rarely given parameters**
 - Usually observe y , X , and W and/or Z , want to infer β , θ , ψ
 - Need each $I-W$, $I-Z$ invertible for solution to exist
- **Typical approach: maximum likelihood**
 - Generally take v as iid, $v_i \sim N(0, \sigma^2)$
 - Standard errors based on the inverse information matrix at the MLE
 - Compare models in the usual way (e.g., AIC, BIC)
 - (Could also do this in a Bayesian way, could substitute other forms for v , etc.)

Choosing the Weight Matrix

- An important modeling issue: choosing the form for \mathbf{W} and/or \mathbf{Z}
 - Must satisfy invertibility condition
 - i,j cell must express the dependence of i on j
- Many suggestions given by Leenders (2002)
- Ex: Friedkin-Johnson criteria
 - Based on small-group social influence experiments
 - Induce quasi-convexity in equilibrium attitudes
 - $\mathbf{W}_{ij} \geq 0, \mathbf{W}_{ij} \leq 1$
 - $\sum_j \mathbf{W}_{ij} \leq 1$
 - Diagonal can be > 0

Important Errors to Avoid

- **Using NAMs when you need NLIs (or vice versa)**
 - Am I affected by my *position*, or by the *properties of my neighbors*?
 - Nearly all influence theories imply some sort of autocorrelation; ditto for some theories of competition
- **Confusing correlation with influence**
 - If influence occurs through direct ties, structurally equivalent positions will correlate; NAM will show an SE effect
- **Confusing AR and MA terms**
 - AR terms imply diffusion of response values, MA terms imply diffusion of perturbations
 - Do my neighbors' covariates affect me? If so, this is an AR process! (If not, it's an MA process.)
- **Forgetting selection effects**
 - NAM assumes that weight matrix is fixed
 - If selection is present, NAM results will be misleading