Virial Theorem

Consider stationary bound state $|\psi\rangle$: $\hat{H}|\psi\rangle = E|\psi\rangle$

Virial Theorem

Consider stationary bound state $|\psi\rangle$: $\hat{H}|\psi\rangle = E|\psi\rangle$

and a linear time independent operator, $\hat{A}$

Virial Theorem

Consider stationary bound state $|\psi\rangle$: $\hat{H}|\psi\rangle = E|\psi\rangle$

and a linear time independent operator, $\hat{A}$

Consider $\langle\psi|[\hat{H},\hat{A}]|\psi\rangle = \langle\psi|\hat{H}\hat{A} - \hat{A}\hat{H}|\psi\rangle = \langle\psi|\hat{H}|A\psi\rangle - E\langle\psi|A|\psi\rangle$

$= E\langle\psi|A|\psi\rangle - E\langle\psi|A|\psi\rangle = 0$

Virial Theorem

Consider stationary bound state $|\psi\rangle$: $\hat{H}|\psi\rangle = E|\psi\rangle$

and a linear time independent operator, $\hat{A}$

Consider $\langle\psi|[\hat{H},\hat{A}]|\psi\rangle = \langle\psi|\hat{H}\hat{A} - \hat{A}\hat{H}|\psi\rangle = \langle\psi|\hat{H}|A\psi\rangle - E\langle\psi|A|\psi\rangle$

$= E\langle\psi|A|\psi\rangle - E\langle\psi|A|\psi\rangle = 0$

$\langle[\hat{H},\hat{A}]\rangle = 0$ [manifold $\{\psi\}$ must be one of the bound states]

Virial Theorem

Consider stationary bound state $|\psi\rangle$: $\hat{H}|\psi\rangle = E|\psi\rangle$

and a linear time independent operator, $\hat{A}$

Consider $\langle\psi|[\hat{H},\hat{A}]|\psi\rangle = \langle\psi|\hat{H}\hat{A} - \hat{A}\hat{H}|\psi\rangle = \langle\psi|\hat{H}|A\psi\rangle - E\langle\psi|A|\psi\rangle$

$= E\langle\psi|A|\psi\rangle - E\langle\psi|A|\psi\rangle = 0$

$\langle[\hat{H},\hat{A}]\rangle = 0$ [manifold $\{\psi\}$ must be one of the bound states]

Choose $\hat{A}$ to be $\sum_i \hat{q}_i\hat{p}_i = -i\hbar\sum_i q_i\frac{\partial}{\partial q_i}$

Virial Theorem

Consider stationary bound state $|\psi\rangle$: $\hat{H}|\psi\rangle = E|\psi\rangle$

and a linear time independent operator, $\hat{A}$

Consider $\langle\psi|[\hat{H},\hat{A}]|\psi\rangle = \langle\psi|\hat{H}\hat{A} - \hat{A}\hat{H}|\psi\rangle = \langle\psi|\hat{H}|A\psi\rangle - E\langle\psi|A|\psi\rangle$

$= E\langle\psi|A|\psi\rangle - E\langle\psi|A|\psi\rangle = 0$

$\langle[\hat{H},\hat{A}]\rangle = 0$ [manifold $\{\psi\}$ must be one of the bound states]

Choose $\hat{A}$ to be $\sum_i \hat{q}_i\hat{p}_i = -i\hbar\sum_i q_i\frac{\partial}{\partial q_i}$

$[\hat{H},\ \sum_i q_i\hat{p}_i] = \sum_i [\hat{H}, q_i\hat{p}_i]$

Virial Theorem

Consider stationary bound state $|\psi\rangle$: $\hat{H}|\psi\rangle = E|\psi\rangle$

and a linear time independent operator, $\hat{A}$

Consider $\langle\psi|[\hat{H},\hat{A}]|\psi\rangle = \langle\psi|\hat{H}\hat{A} - \hat{A}\hat{H}|\psi\rangle = \langle\psi|\hat{H}|A\psi\rangle - E\langle\psi|A|\psi\rangle$

$= E\langle\psi|A|\psi\rangle - E\langle\psi|A|\psi\rangle = 0$

$\langle[\hat{H},\hat{A}]\rangle = 0$ [manifold $\{\psi\}$ must be one of the bound states]

Choose $\hat{A}$ to be $\sum_i \hat{q}_i\hat{p}_i = -i\hbar\sum_i q_i\frac{\partial}{\partial q_i}$

$[\hat{H},\ \sum_i q_i\hat{p}_i] = \sum_i[\hat{H}, q_i\hat{p}_i] = \sum_i \hat{q}_i[\hat{H},\hat{p}_i] + \sum_i[\hat{H},\hat{q}_i]\hat{p}_i$

Virial Theorem

Consider stationary bound state $|\psi\rangle$: $\hat{H}|\psi\rangle = E|\psi\rangle$

and a linear time independent operator, $\hat{A}$

Consider $\langle\psi|[\hat{H},\hat{A}]|\psi\rangle = \langle\psi|\hat{H}\hat{A} - \hat{A}\hat{H}|\psi\rangle = \langle\psi|\hat{H}|A\psi\rangle - E\langle\psi|A|\psi\rangle$

$= E\langle\psi|A|\psi\rangle - E\langle\psi|A|\psi\rangle = 0$

$\langle[\hat{H},\hat{A}]\rangle = 0$ [manifold $\{\psi\}$ must be one of the bound states]

Choose $\hat{A}$ to be $\sum_i \hat{q}_i\hat{p}_i = -i\hbar\sum_i q_i\frac{\partial}{\partial q_i}$

$[\hat{H},\ \sum_i q_i\hat{p}_i] = \sum_i[\hat{H}, q_i\hat{p}_i] = \sum_i \hat{q}_i[\hat{H},\hat{p}_i] + \sum_i[\hat{H}, \hat{q}_i]\hat{p}_i$

$= i\hbar\sum_i q_i\frac{\partial V}{\partial q_i} - i\hbar\sum_i \frac{1}{m_i}\hat{p}_i^2$

Virial Theorem

Consider stationary bound state $|\psi\rangle$: $\hat{H}|\psi\rangle = E|\psi\rangle$

and a linear time independent operator, $\hat{A}$

Consider $\langle\psi|[\hat{H},\hat{A}]|\psi\rangle = \langle\psi|\hat{H}\hat{A} - \hat{A}\hat{H}|\psi\rangle = \langle\psi|\hat{H}|A\psi\rangle - E\langle\psi|A|\psi\rangle$

$= E\langle\psi|A|\psi\rangle - E\langle\psi|A|\psi\rangle = 0$

$\langle[\hat{H},\hat{A}]\rangle = 0$ [manifold $\{\psi\}$ must be one of the bound states]

Choose $\hat{A}$ to be $\sum_i \hat{q}_i\hat{p}_i = -i\hbar\sum_i q_i\frac{\partial}{\partial q_i}$

$[\hat{H},\ \sum_i q_i\hat{p}_i] = \sum_i[\hat{H}, q_i\hat{p}_i] = \sum_i \hat{q}_i[\hat{H},\hat{p}_i] + \sum_i[\hat{H},\hat{q}_i]\hat{p}_i$

$= i\hbar\sum_i q_i\frac{\partial V}{\partial q_i} - i\hbar\sum_i \frac{1}{m_i}\hat{p}_i^2 \qquad = i\hbar\sum_i q_i\frac{\partial V}{\partial q_i} - 2i\hbar T$

Virial Theorem

Consider stationary bound state $|\psi\rangle$: $\hat{H}|\psi\rangle = E|\psi\rangle$

and a linear time independent operator, $\hat{A}$

Consider $\langle\psi|[\hat{H},\hat{A}]|\psi\rangle = \langle\psi|\hat{H}\hat{A} - \hat{A}\hat{H}|\psi\rangle = \langle\psi|\hat{H}|A\psi\rangle - E\langle\psi|A|\psi\rangle$

$= E\langle\psi|A|\psi\rangle - E\langle\psi|A|\psi\rangle = 0$

$\langle[\hat{H},\hat{A}]\rangle = 0$ [manifold $\{\psi\}$ must be one of the bound states]

Choose $\hat{A}$ to be $\sum_i \hat{q}_i\hat{p}_i = -i\hbar\sum_i q_i\frac{\partial}{\partial q_i}$

$[\hat{H},\ \sum_i q_i\hat{p}_i] = \sum_i[\hat{H}, q_i\hat{p}_i] = \sum_i \hat{q}_i[\hat{H},\hat{p}_i] + \sum_i[\hat{H},\hat{q}_i]\hat{p}_i$

$= i\hbar\sum_i q_i\frac{\partial V}{\partial q_i} - i\hbar\sum_i \frac{1}{m_i}\hat{p}_i^2 \qquad = i\hbar\sum_i q_i\frac{\partial V}{\partial q_i} - 2i\hbar T$

$= i\hbar(\sum_i q_i\frac{\partial V}{\partial q_i} - 2T)$

Virial Theorem

Consider stationary bound state $|\psi\rangle$: $\hat{H}|\psi\rangle = E|\psi\rangle$

and a linear time independent operator, $\hat{A}$

Consider $\langle\psi|[\hat{H},\hat{A}]|\psi\rangle = \langle\psi|\hat{H}\hat{A} - \hat{A}\hat{H}|\psi\rangle = \langle\psi|\hat{H}|A\psi\rangle - E\langle\psi|A|\psi\rangle$

$= E\langle\psi|A|\psi\rangle - E\langle\psi|A|\psi\rangle = 0$

$\langle[\hat{H},\hat{A}]\rangle = 0$ [manifold $\{\psi\}$ must be one of the bound states]

Choose $\hat{A}$ to be $\sum_i \hat{q}_i\hat{p}_i = -i\hbar\sum_i q_i\frac{\partial}{\partial q_i}$

$[\hat{H},\ \sum_i q_i\hat{p}_i] = \sum_i[\hat{H}, q_i\hat{p}_i] = \sum_i \hat{q}_i[\hat{H},\hat{p}_i] + \sum_i[\hat{H},\hat{q}_i]\hat{p}_i$

$= i\hbar\sum_i q_i\frac{\partial V}{\partial q_i} - i\hbar\sum_i \frac{1}{m_i}\hat{p}_i^2 \qquad = i\hbar\sum_i q_i\frac{\partial V}{\partial q_i} - 2i\hbar T$

$= i\hbar(\sum_i q_i\frac{\partial V}{\partial q_i} - 2T)$

$\therefore$ Hypervirial theorem : $\boxed{\left\langle\sum_i q_i\frac{\partial V}{\partial q_i}\right\rangle = 2\langle T\rangle}$

In particular, if $V = \sum_i V_i$ and $V_i \propto q_i^n$,

In particular, if $V = \sum_i V_i$ and $V_i \propto q_i^n$,

e. g., $V_i = \alpha q_i^n \Longrightarrow \frac{\partial V}{\partial q_i} = n\alpha q_i^{n-1}$

In particular, if $V = \sum_i V_i$ and $V_i \propto q_i^n$,

e. g., $V_i = \alpha q_i^n \Longrightarrow \frac{\partial V}{\partial q_i} = n\alpha q_i^{n-1} \quad \Longrightarrow q_i \frac{\partial V}{\partial q_i} = nV_i$

In particular, if $V = \sum_i V_i$ and $V_i \propto q_i^n$,

e. g., $V_i = \alpha q_i^n \Longrightarrow \frac{\partial V}{\partial q_i} = n\alpha q_i^{n-1} \quad \Longrightarrow q_i \frac{\partial V}{\partial q_i} = nV_i$

$$\boxed{n\langle V \rangle = 2\langle T \rangle} : \text{virial theorem}$$

In particular, if $V = \sum_i V_i$ and $V_i \propto q_i^n$,

e. g., $V_i = \alpha q_i^n \Longrightarrow \frac{\partial V}{\partial q_i} = n\alpha q_i^{n-1} \quad \Longrightarrow q_i \frac{\partial V}{\partial q_i} = nV_i$

$$\boxed{n\langle V \rangle = 2\langle T \rangle} \text{ : virial theorem}$$

or $\langle T \rangle = \frac{n}{2}\langle V \rangle$

In particular, if $V = \sum_i V_i$ and $V_i \propto q_i^n$,

e. g., $V_i = \alpha q_i^n \Longrightarrow \frac{\partial V}{\partial q_i} = n\alpha q_i^{n-1} \quad \Longrightarrow q_i \frac{\partial V}{\partial q_i} = nV_i$

$$\boxed{n\langle V \rangle = 2\langle T \rangle} \text{ : virial theorem}$$

or $\langle T \rangle = \frac{n}{2}\langle V \rangle$

e. g., for coulomb potential, $n = -1 \Longrightarrow \langle T \rangle = -\frac{1}{2}\langle V \rangle$

In particular, if $V = \sum_i V_i$ and $V_i \propto q_i^n$,

e. g., $V_i = \alpha q_i^n \Longrightarrow \frac{\partial V}{\partial q_i} = n\alpha q_i^{n-1} \quad \Longrightarrow q_i \frac{\partial V}{\partial q_i} = nV_i$

$$\boxed{n\langle V \rangle = 2\langle T \rangle} \text{ : virial theorem}$$

or $\langle T \rangle = \frac{n}{2}\langle V \rangle$

e. g., for coulomb potential, $n = -1 \Longrightarrow \langle T \rangle = -\frac{1}{2}\langle V \rangle$

and $\langle E_{\text{total}} \rangle = \langle T \rangle + \langle V \rangle = -\frac{1}{2}\langle V \rangle + \langle V \rangle = \frac{1}{2}\langle V \rangle$

UNCERTAINTY PRINCIPLE : If two operators $\hat{A}$ and $\hat{B}$ do not commute, there exists NO state for which outcome of simultaneous measurement of observables corresponding to $\hat{A}$ and $\hat{B}$ can be known with precision

UNCERTAINTY PRINCIPLE : If two operators $\hat{A}$ and $\hat{B}$ do not commute, there exists NO state for which outcome of simultaneous measurement of observables corresponding to $\hat{A}$ and $\hat{B}$ can be known with precision

Consider an arbitrary state ψ

UNCERTAINTY PRINCIPLE : If two operators $\hat{A}$ and $\hat{B}$ do not commute, there exists NO state for which outcome of simultaneous measurement of observables corresponding to $\hat{A}$ and $\hat{B}$ can be known with precision

Consider an arbitrary state ψ

Form the operator $\hat{A} - i\lambda\hat{B}$; $\hat{A}$ and $\hat{B}$ are Hermitian operators,

UNCERTAINTY PRINCIPLE : If two operators $\hat{A}$ and $\hat{B}$ do not commute, there exists NO state for which outcome of simultaneous measurement of observables corresponding to $\hat{A}$ and $\hat{B}$ can be known with precision

Consider an arbitrary state ψ

Form the operator $\hat{A} - i\lambda\hat{B}$; $\hat{A}$ and $\hat{B}$ are Hermitian operators,

$\langle\psi_1|\hat{A}|\psi_2\rangle = \langle\left(\hat{A}\psi_1\right)|\psi_2\rangle$ [$\because \hat{A}^\dagger = \hat{A}$] and similarly for $\hat{B}$; λ is real but arbitrary

UNCERTAINTY PRINCIPLE : If two operators $\hat{A}$ and $\hat{B}$ do not commute, there exists NO state for which outcome of simultaneous measurement of observables corresponding to $\hat{A}$ and $\hat{B}$ can be known with precision

Consider an arbitrary state ψ

Form the operator $\hat{A} - i\lambda\hat{B}$; $\hat{A}$ and $\hat{B}$ are Hermitian operators,

$\langle\psi_1|\hat{A}|\psi_2\rangle = \langle\left(\hat{A}\psi_1\right)|\psi_2\rangle$ [$\because \hat{A}^\dagger = \hat{A}$] and similarly for $\hat{B}$; λ is real but arbitrary

Then if $(\hat{A} + i\lambda\hat{B})|\psi\rangle = |\phi\rangle$, construct $g = \langle\phi|\phi\rangle \geq 0$

UNCERTAINTY PRINCIPLE : If two operators $\hat{A}$ and $\hat{B}$ do not commute, there exists NO state for which outcome of simultaneous measurement of observables corresponding to $\hat{A}$ and $\hat{B}$ can be known with precision

Consider an arbitrary state ψ

Form the operator $\hat{A} - i\lambda\hat{B}$; $\hat{A}$ and $\hat{B}$ are Hermitian operators,

$\langle\psi_1|\hat{A}|\psi_2\rangle = \langle\left(\hat{A}\psi_1\right)|\psi_2\rangle$ [$\because \hat{A}^\dagger = \hat{A}$] and similarly for $\hat{B}$; λ is real but arbitrary

Then if $(\hat{A} + i\lambda\hat{B})|\psi\rangle = |\phi\rangle$, construct $g = \langle\phi|\phi\rangle \geq 0$

Form $g = \langle\psi|(\hat{A} - i\lambda\hat{B})(\hat{A} + i\lambda\hat{B})|\psi\rangle$

UNCERTAINTY PRINCIPLE : If two operators $\hat{A}$ and $\hat{B}$ do not commute, there exists NO state for which outcome of simultaneous measurement of observables corresponding to $\hat{A}$ and $\hat{B}$ can be known with precision

Consider an arbitrary state ψ

Form the operator $\hat{A} - i\lambda\hat{B}$; $\hat{A}$ and $\hat{B}$ are Hermitian operators,

$\langle\psi_1|\hat{A}|\psi_2\rangle = \langle\left(\hat{A}\psi_1\right)|\psi_2\rangle$ [$\because \hat{A}^\dagger = \hat{A}$] and similarly for $\hat{B}$; λ is real but arbitrary

Then if $(\hat{A} + i\lambda\hat{B})|\psi\rangle = |\phi\rangle$, construct $g = \langle\phi|\phi\rangle \geq 0$

Form $g = \langle\psi|(\hat{A} - i\lambda\hat{B})(\hat{A} + i\lambda\hat{B})|\psi\rangle$
$= \langle A^2\rangle + \lambda^2\langle B^2\rangle + i\lambda\langle\psi|\hat{A}\hat{B} - \hat{B}\hat{A}|\psi\rangle$

UNCERTAINTY PRINCIPLE : If two operators $\hat{A}$ and $\hat{B}$ do not commute, there exists NO state for which outcome of simultaneous measurement of observables corresponding to $\hat{A}$ and $\hat{B}$ can be known with precision

Consider an arbitrary state ψ

Form the operator $\hat{A} - i\lambda\hat{B}$; $\hat{A}$ and $\hat{B}$ are Hermitian operators,

$\langle\psi_1|\hat{A}|\psi_2\rangle = \langle\left(\hat{A}\psi_1\right)|\psi_2\rangle$ [$\because \hat{A}^\dagger = \hat{A}$] and similarly for $\hat{B}$; λ is real but arbitrary

Then if $(\hat{A} + i\lambda\hat{B})|\psi\rangle = |\phi\rangle$, construct $g = \langle\phi|\phi\rangle \geq 0$

Form $g = \langle\psi|(\hat{A} - i\lambda\hat{B})(\hat{A} + i\lambda\hat{B})|\psi\rangle$

$= \langle A^2\rangle + \lambda^2\langle B^2\rangle + i\lambda\langle\psi|\hat{A}\hat{B} - \hat{B}\hat{A}|\psi\rangle \qquad = \langle A^2\rangle + \lambda^2\langle B^2\rangle + \lambda\langle i[\hat{A}, \hat{B}]\rangle$

UNCERTAINTY PRINCIPLE : If two operators $\hat{A}$ and $\hat{B}$ do not commute, there exists NO state for which outcome of simultaneous measurement of observables corresponding to $\hat{A}$ and $\hat{B}$ can be known with precision

Consider an arbitrary state ψ

Form the operator $\hat{A} - i\lambda\hat{B}$; $\hat{A}$ and $\hat{B}$ are Hermitian operators,

$\langle\psi_1|\hat{A}|\psi_2\rangle = \langle\left(\hat{A}\psi_1\right)|\psi_2\rangle$ [$\because \hat{A}^\dagger = \hat{A}$] and similarly for $\hat{B}$; λ is real but arbitrary

Then if $(\hat{A} + i\lambda\hat{B})|\psi\rangle = |\phi\rangle$, construct $g = \langle\phi|\phi\rangle \geq 0$

Form $g = \langle\psi|(\hat{A} - i\lambda\hat{B})(\hat{A} + i\lambda\hat{B})|\psi\rangle$
$= \langle A^2\rangle + \lambda^2\langle B^2\rangle + i\lambda\langle\psi|\hat{A}\hat{B} - \hat{B}\hat{A}|\psi\rangle \qquad = \langle A^2\rangle + \lambda^2\langle B^2\rangle + \lambda\langle i[\hat{A}, \hat{B}]\rangle$

It can be shown that $\lambda\langle i[A, \hat{B}]\rangle$ is Hermitian:

UNCERTAINTY PRINCIPLE : If two operators $\hat{A}$ and $\hat{B}$ do not commute, there exists NO state for which outcome of simultaneous measurement of observables corresponding to $\hat{A}$ and $\hat{B}$ can be known with precision

Consider an arbitrary state ψ

Form the operator $\hat{A} - i\lambda\hat{B}$; $\hat{A}$ and $\hat{B}$ are Hermitian operators,

$\langle\psi_1|\hat{A}|\psi_2\rangle = \langle\left(\hat{A}\psi_1\right)|\psi_2\rangle$ [$\because \hat{A}^\dagger = \hat{A}$] and similarly for $\hat{B}$; λ is real but arbitrary

Then if $(\hat{A} + i\lambda\hat{B})|\psi\rangle = |\phi\rangle$, construct $g = \langle\phi|\phi\rangle \geq 0$

Form $g = \langle\psi|(\hat{A} - i\lambda\hat{B})(\hat{A} + i\lambda\hat{B})|\psi\rangle$
$= \langle A^2\rangle + \lambda^2\langle B^2\rangle + i\lambda\langle\psi|\hat{A}\hat{B} - \hat{B}\hat{A}|\psi\rangle \qquad = \langle A^2\rangle + \lambda^2\langle B^2\rangle + \lambda\langle i[\hat{A}, \hat{B}]\rangle$

It can be shown that $\lambda\langle i[A, \hat{B}]\rangle$ is Hermitian:

$\langle\psi_1|i[A, B]|\psi_2\rangle = \langle\psi_1|i(\hat{A}\hat{B} - \hat{B}\hat{A})|\psi_2\rangle = i\langle\psi_1|\hat{A}\hat{B}|\psi_2\rangle - i\langle\psi_1|\hat{B}\hat{A}|\psi_2\rangle$

UNCERTAINTY PRINCIPLE : If two operators $\hat{A}$ and $\hat{B}$ do not commute, there exists NO state for which outcome of simultaneous measurement of observables corresponding to $\hat{A}$ and $\hat{B}$ can be known with precision

Consider an arbitrary state ψ

Form the operator $\hat{A} - i\lambda\hat{B}$; $\hat{A}$ and $\hat{B}$ are Hermitian operators,

$\langle\psi_1|\hat{A}|\psi_2\rangle = \langle\left(\hat{A}\psi_1\right)|\psi_2\rangle$ [$\because \hat{A}^\dagger = \hat{A}$] and similarly for $\hat{B}$; λ is real but arbitrary

Then if $(\hat{A} + i\lambda\hat{B})|\psi\rangle = |\phi\rangle$, construct $g = \langle\phi|\phi\rangle \geq 0$

Form $g = \langle\psi|(\hat{A} - i\lambda\hat{B})(\hat{A} + i\lambda\hat{B})|\psi\rangle$

$= \langle A^2\rangle + \lambda^2\langle B^2\rangle + i\lambda\langle\psi|\hat{A}\hat{B} - \hat{B}\hat{A}|\psi\rangle \qquad = \langle A^2\rangle + \lambda^2\langle B^2\rangle + \lambda\langle i[\hat{A}, \hat{B}]\rangle$

It can be shown that $\lambda\langle i[A, \hat{B}]\rangle$ is Hermitian:

$\langle\psi_1|i[A, B]|\psi_2\rangle = \langle\psi_1|i(\hat{A}\hat{B} - \hat{B}\hat{A})|\psi_2\rangle = i\langle\psi_1|\hat{A}\hat{B}|\psi_2\rangle - i\langle\psi_1|\hat{B}\hat{A}|\psi_2\rangle$

$= i\langle\hat{A}\psi_1|\hat{B}|\psi_2\rangle - i\langle\hat{B}\psi_1|\hat{A}|\psi_2\rangle = +i\langle\hat{B}\hat{A}\psi_1|\psi_2\rangle - i\langle\hat{A}\hat{B}\psi_1|\psi_2\rangle$

UNCERTAINTY PRINCIPLE : If two operators $\hat{A}$ and $\hat{B}$ do not commute, there exists NO state for which outcome of simultaneous measurement of observables corresponding to $\hat{A}$ and $\hat{B}$ can be known with precision

Consider an arbitrary state ψ

Form the operator $\hat{A} - i\lambda\hat{B}$; $\hat{A}$ and $\hat{B}$ are Hermitian operators,

$\langle\psi_1|\hat{A}|\psi_2\rangle = \langle\left(\hat{A}\psi_1\right)|\psi_2\rangle$ [$\because \hat{A}^\dagger = \hat{A}$] and similarly for $\hat{B}$; λ is real but arbitrary

Then if $(\hat{A} + i\lambda\hat{B})|\psi\rangle = |\phi\rangle$, construct $g = \langle\phi|\phi\rangle \geq 0$

Form $g = \langle\psi|(\hat{A} - i\lambda\hat{B})(\hat{A} + i\lambda\hat{B})|\psi\rangle$
$= \langle A^2\rangle + \lambda^2\langle B^2\rangle + i\lambda\langle\psi|\hat{A}\hat{B} - \hat{B}\hat{A}|\psi\rangle \qquad = \langle A^2\rangle + \lambda^2\langle B^2\rangle + \lambda\langle i[\hat{A}, \hat{B}]\rangle$

It can be shown that $\lambda\langle i[A, \hat{B}]\rangle$ is Hermitian:

$\langle\psi_1|i[A, B]|\psi_2\rangle = \langle\psi_1|i(\hat{A}\hat{B} - \hat{B}\hat{A})|\psi_2\rangle = i\langle\psi_1|\hat{A}\hat{B}|\psi_2\rangle - i\langle\psi_1|\hat{B}\hat{A}|\psi_2\rangle$

$= i\langle\hat{A}\psi_1|\hat{B}|\psi_2\rangle - i\langle\hat{B}\psi_1|\hat{A}|\psi_2\rangle = +i\langle\hat{B}\hat{A}\psi_1|\psi_2\rangle - i\langle\hat{A}\hat{B}\psi_1|\psi_2\rangle$

$= \langle i(\hat{B}\hat{A} - \hat{A}\hat{B})\psi_1|\psi_2\rangle$ [i on the left has the value $i^* = -i$]

UNCERTAINTY PRINCIPLE : If two operators $\hat{A}$ and $\hat{B}$ do not commute, there exists NO state for which outcome of simultaneous measurement of observables corresponding to $\hat{A}$ and $\hat{B}$ can be known with precision

Consider an arbitrary state ψ

Form the operator $\hat{A} - i\lambda\hat{B}$; $\hat{A}$ and $\hat{B}$ are Hermitian operators,

$\langle\psi_1|\hat{A}|\psi_2\rangle = \langle\left(\hat{A}\psi_1\right)|\psi_2\rangle$ [$\because \hat{A}^\dagger = \hat{A}$] and similarly for $\hat{B}$; λ is real but arbitrary

Then if $(\hat{A} + i\lambda\hat{B})|\psi\rangle = |\phi\rangle$, construct $g = \langle\phi|\phi\rangle \geq 0$

Form $g = \langle\psi|(\hat{A} - i\lambda\hat{B})(\hat{A} + i\lambda\hat{B})|\psi\rangle$
$= \langle A^2\rangle + \lambda^2\langle B^2\rangle + i\lambda\langle\psi|\hat{A}\hat{B} - \hat{B}\hat{A}|\psi\rangle \qquad = \langle A^2\rangle + \lambda^2\langle B^2\rangle + \lambda\langle i[\hat{A}, \hat{B}]\rangle$

It can be shown that $\lambda\langle i[A, \hat{B}]\rangle$ is Hermitian:

$\langle\psi_1|i[A, B]|\psi_2\rangle = \langle\psi_1|i(\hat{A}\hat{B} - \hat{B}\hat{A})|\psi_2\rangle = i\langle\psi_1|\hat{A}\hat{B}|\psi_2\rangle - i\langle\psi_1|\hat{B}\hat{A}|\psi_2\rangle$

$= i\langle\hat{A}\psi_1|\hat{B}|\psi_2\rangle - i\langle\hat{B}\psi_1|\hat{A}|\psi_2\rangle = +i\langle\hat{B}\hat{A}\psi_1|\psi_2\rangle - i\langle\hat{A}\hat{B}\psi_1|\psi_2\rangle$

$= \langle i(\hat{B}\hat{A} - \hat{A}\hat{B})\psi_1|\psi_2\rangle$ [i on the left has the value $i^* = -i$]

$= \langle(i[A, B])\,\psi_1|\psi_2\rangle$

$$g = \langle \phi | \phi \rangle \geq 0$$

$g = \langle \phi | \phi \rangle \geq 0$

the equality holds if $|\phi\rangle \equiv 0$

$g = \langle \phi | \phi \rangle \geq 0$

the equality holds if $|\phi\rangle \equiv 0$

$$\Longrightarrow (\hat{A} + i\lambda \hat{B})|\psi\rangle = 0$$

$g = \langle\phi|\phi\rangle \geq 0$

the equality holds if $|\phi\rangle \equiv 0$

$$\Longrightarrow (\hat{A} + i\lambda\hat{B})|\psi\rangle = 0 \qquad \Longrightarrow \hat{A} = -i\lambda\hat{B} \text{ for arbitrary } |\psi\rangle$$

$g = \langle\phi|\phi\rangle \geq 0$

the equality holds if $|\phi\rangle \equiv 0$

$\Longrightarrow (\hat{A} + i\lambda\hat{B})|\psi\rangle = 0 \qquad \Longrightarrow \hat{A} = -i\lambda\hat{B}$ for arbitrary $|\psi\rangle$

let us minimise g with respect to λ

$g = \langle\phi|\phi\rangle \geq 0$

the equality holds if $|\phi\rangle \equiv 0$

$$\implies (\hat{A} + i\lambda\hat{B})|\psi\rangle = 0 \qquad \implies \hat{A} = -i\lambda\hat{B} \text{ for arbitrary } |\psi\rangle$$

let us minimise g with respect to λ

$\frac{\partial g}{\partial \lambda} = 0 \implies \lambda = -\frac{1}{2}\frac{\langle i[\hat{A},\hat{B}]\rangle}{\langle \widehat{B^2}\rangle}$

$g = \langle\phi|\phi\rangle \geq 0$

the equality holds if $|\phi\rangle \equiv 0$

$$\implies (\hat{A} + i\lambda\hat{B})|\psi\rangle = 0 \qquad \implies \hat{A} = -i\lambda\hat{B} \text{ for arbitrary } |\psi\rangle$$

let us minimise g with respect to λ

$$\frac{\partial g}{\partial \lambda} = 0 \implies \lambda = -\frac{1}{2}\frac{\langle i[\hat{A},\hat{B}]\rangle}{\langle \widehat{B^2}\rangle}$$

$$\therefore g_{\text{min}} = \langle \hat{A}^2\rangle - \frac{1}{4}\frac{\langle i[\hat{A},\hat{B}]\rangle^2}{\langle \widehat{B^2}\rangle}$$

$g = \langle\phi|\phi\rangle \geq 0$

the equality holds if $|\phi\rangle \equiv 0$

$$\implies (\hat{A} + i\lambda\hat{B})|\psi\rangle = 0 \qquad \implies \hat{A} = -i\lambda\hat{B} \text{ for arbitrary } |\psi\rangle$$

let us minimise g with respect to λ

$$\frac{\partial g}{\partial \lambda} = 0 \implies \lambda = -\frac{1}{2}\frac{\langle i[\hat{A},\hat{B}]\rangle}{\langle \widehat{B^2}\rangle}$$

$$\therefore g_{\text{min}} = \langle \hat{A}^2\rangle - \frac{1}{4}\frac{\langle i[\hat{A},\hat{B}]\rangle^2}{\langle \widehat{B^2}\rangle}$$

g_{min} is never negative. Even at its minimum value $g_{\text{min}} \geq 0$

$g = \langle\phi|\phi\rangle \geq 0$

the equality holds if $|\phi\rangle \equiv 0$

$\implies (\hat{A} + i\lambda\hat{B})|\psi\rangle = 0 \qquad \implies \hat{A} = -i\lambda\hat{B}$ for arbitrary $|\psi\rangle$

let us minimise g with respect to λ

$\frac{\partial g}{\partial \lambda} = 0 \implies \lambda = -\frac{1}{2}\frac{\langle i[\hat{A},\hat{B}]\rangle}{\langle \widehat{B^2} \rangle}$

$\therefore g_{\text{min}} = \langle \hat{A}^2 \rangle - \frac{1}{4}\frac{\langle i[\hat{A},\hat{B}]\rangle^2}{\langle \widehat{B^2} \rangle}$

g_{min} is never negative. Even at its minimum value $g_{\text{min}} \geq 0$

$\implies \langle \hat{A}^2 \rangle \langle \hat{B}^2 \rangle \geq \frac{1}{4}\langle i[\hat{A},\hat{B}]\rangle^2$

what are the special states of ψ for which equality holds ?

what are the special states of ψ for which equality holds ?

Let ψ refer to these states

what are the special states of ψ for which equality holds ?

Let ψ refer to these states$\Longrightarrow (\hat{A} + i\lambda\hat{B})\hat{\psi} = 0$

what are the special states of ψ for which equality holds ?

Let ψ refer to these states$\Longrightarrow (\hat{A} + i\lambda\hat{B})\hat{\psi} = 0$

we can have a family of such states taking different λs

what are the special states of ψ for which equality holds ?

Let ψ refer to these states$\Longrightarrow (\hat{A} + i\lambda\hat{B})\hat{\psi} = 0$

we can have a family of such states taking different λs

$\langle\psi|\hat{A} + i\lambda\hat{B}|\psi\rangle = 0 = \langle A\rangle + i\lambda\langle B\rangle = 0$

what are the special states of ψ for which equality holds ?

Let ψ refer to these states$\Longrightarrow (\hat{A} + i\lambda\hat{B})\hat{\psi} = 0$

we can have a family of such states taking different λs

$\langle\psi|\hat{A} + i\lambda\hat{B}|\psi\rangle = 0 = \langle A\rangle + i\lambda\langle B\rangle = 0 \Longrightarrow \langle A\rangle = 0 = \langle B\rangle$

what are the special states of ψ for which equality holds ?

Let ψ refer to these states$\Longrightarrow (\hat{A} + i\lambda\hat{B})\hat{\psi} = 0$

we can have a family of such states taking different λs

$\langle\psi|\hat{A} + i\lambda\hat{B}|\psi\rangle = 0 = \langle A\rangle + i\lambda\langle B\rangle = 0 \Longrightarrow \langle A\rangle = 0 = \langle B\rangle$

Consider $\hat{A} = \hat{p_x} - \overline{p_x}$ (mean deviation)

what are the special states of ψ for which equality holds ?

Let ψ refer to these states$\Longrightarrow (\hat{A} + i\lambda\hat{B})\hat{\psi} = 0$

we can have a family of such states taking different λs

$\langle\psi|\hat{A} + i\lambda\hat{B}|\psi\rangle = 0 = \langle A\rangle + i\lambda\langle B\rangle = 0 \Longrightarrow \langle A\rangle = 0 = \langle B\rangle$

Consider $\hat{A} = \hat{p_x} - \overline{p_x}$ (mean deviation)

$$\hat{B} = \hat{x} - \overline{x} \Longrightarrow i[\hat{A}, \hat{B}] = i(-i\hbar) = +\hbar$$

what are the special states of ψ for which equality holds ?

Let ψ refer to these states$\Longrightarrow (\hat{A} + i\lambda\hat{B})\hat{\psi} = 0$

we can have a family of such states taking different λs

$\langle\psi|\hat{A} + i\lambda\hat{B}|\psi\rangle = 0 = \langle A\rangle + i\lambda\langle B\rangle = 0 \Longrightarrow \langle A\rangle = 0 = \langle B\rangle$

Consider $\hat{A} = \hat{p_x} - \overline{p_x}$ (mean deviation)

$$\hat{B} = \hat{x} - \overline{x} \Longrightarrow i[\hat{A}, \hat{B}] = i(-i\hbar) = +\hbar$$

$$\Longrightarrow \langle(\hat{p} - \overline{p})^2\rangle\langle(\hat{x} - \overline{x})^2\rangle \geq \frac{\hbar^2}{4}$$

what are the special states of ψ for which equality holds ?

Let ψ refer to these states$\Longrightarrow (\hat{A} + i\lambda\hat{B})\hat{\psi} = 0$

we can have a family of such states taking different λs

$\langle\psi|\hat{A} + i\lambda\hat{B}|\psi\rangle = 0 = \langle A\rangle + i\lambda\langle B\rangle = 0 \Longrightarrow \langle A\rangle = 0 = \langle B\rangle$

Consider $\hat{A} = \hat{p_x} - \overline{p_x}$ (mean deviation)

$$\hat{B} = \hat{x} - \overline{x} \Longrightarrow i[\hat{A}, \hat{B}] = i(-i\hbar) = +\hbar$$

$$\Longrightarrow \langle(\hat{p} - \overline{p})^2\rangle\langle(\hat{x} - \overline{x})^2\rangle \geq \frac{\hbar^2}{4}$$

$$\Longrightarrow \triangle p_x \triangle x \geq \frac{\hbar}{2},$$

what are the special states of ψ for which equality holds ?

Let ψ refer to these states$\Longrightarrow (\hat{A} + i\lambda\hat{B})\hat{\psi} = 0$

we can have a family of such states taking different λs

$\langle\psi|\hat{A} + i\lambda\hat{B}|\psi\rangle = 0 = \langle A\rangle + i\lambda\langle B\rangle = 0 \Longrightarrow \langle A\rangle = 0 = \langle B\rangle$

Consider $\hat{A} = \hat{p_x} - \overline{p_x}$ (mean deviation)

$\hat{B} = \hat{x} - \overline{x} \Longrightarrow i[\hat{A}, \hat{B}] = i(-i\hbar) = +\hbar$

$\Longrightarrow \langle(\hat{p} - \overline{p})^2\rangle\langle(\hat{x} - \overline{x})^2\rangle \geq \frac{\hbar^2}{4}$

$\Longrightarrow \triangle p_x \triangle x \geq \frac{\hbar}{2}$, where $\triangle p_x \equiv \sigma_{p_x} = \sqrt{(\hat{p} - \overline{p})^2}$ and $\triangle x \equiv \sigma_x = \sqrt{(\hat{x} - \overline{x})^2}$

Proof using Schwarz inequality :

Proof using Schwarz inequality :

Consider for an observable A :

Proof using Schwarz inequality :

Consider for an observable A :$\sigma_A^2 = \langle(\hat{A} - \langle A\rangle)\Psi|(\hat{A} - \langle A\rangle)\Psi\rangle = \langle f|f\rangle$, where $f \equiv \left(\hat{A} - \langle A\rangle\right)\Psi$

Proof using Schwarz inequality :

Consider for an observable A :$\sigma_A^2 = \langle(\hat{A} - \langle A\rangle)\Psi|(\hat{A} - \langle A\rangle)\Psi\rangle = \langle f|f\rangle$, where $f \equiv \left(\hat{A} - \langle A\rangle\right)\Psi$

Likewise, for any other observable, B,

Proof using Schwarz inequality :

Consider for an observable A :$\sigma_A^2 = \langle(\hat{A} - \langle A\rangle)\Psi|(\hat{A} - \langle A\rangle)\Psi\rangle = \langle f|f\rangle$, where $f \equiv \left(\hat{A} - \langle A\rangle\right)\Psi$

Likewise, for any other observable, B, $\sigma_B^2 = \langle g \mid g\rangle$, where $g \equiv \left(\hat{B} - \langle B\rangle\right)\Psi$

Proof using Schwarz inequality :

Consider for an observable A :$\sigma_A^2 = \langle(\hat{A} - \langle A\rangle)\Psi|(\hat{A} - \langle A\rangle)\Psi\rangle = \langle f|f\rangle$, where $f \equiv \left(\hat{A} - \langle A\rangle\right)\Psi$

Likewise, for any other observable, B, $\sigma_B^2 = \langle g \mid g\rangle$, where $g \equiv \left(\hat{B} - \langle B\rangle\right)\Psi$

Schwarz inequality : $\sigma_A^2\sigma_B^2 = \langle f \mid f\rangle\langle g \mid g\rangle \geq |\langle f \mid g\rangle|^2$

Proof using Schwarz inequality :

Consider for an observable A :$\sigma_A^2 = \langle(\hat{A} - \langle A\rangle)\Psi|(\hat{A} - \langle A\rangle)\Psi\rangle = \langle f|f\rangle$, where $f \equiv \left(\hat{A} - \langle A\rangle\right)\Psi$

Likewise, for any other observable, B, $\sigma_B^2 = \langle g \mid g\rangle$, where $g \equiv \left(\hat{B} - \langle B\rangle\right)\Psi$

Schwarz inequality : $\sigma_A^2\sigma_B^2 = \langle f \mid f\rangle\langle g \mid g\rangle \geq |\langle f \mid g\rangle|^2$

For any complex number z,

$|z|^2 = [\mathrm{Re}(z)]^2 + [\mathrm{Im}(z)]^2 \geq [\mathrm{Im}(z)]^2 = \left[\frac{1}{2i}(z - z^*)\right]^2$

Proof using Schwarz inequality :

Consider for an observable A :$\sigma_A^2 = \langle(\hat{A} - \langle A\rangle)\Psi|(\hat{A} - \langle A\rangle)\Psi\rangle = \langle f|f\rangle$, where $f \equiv \left(\hat{A} - \langle A\rangle\right)\Psi$

Likewise, for any other observable, B, $\sigma_B^2 = \langle g \mid g\rangle$, where $g \equiv \left(\hat{B} - \langle B\rangle\right)\Psi$

Schwarz inequality : $\sigma_A^2\sigma_B^2 = \langle f \mid f\rangle\langle g \mid g\rangle \geq |\langle f \mid g\rangle|^2$

For any complex number z,

$|z|^2 = [\mathrm{Re}(z)]^2 + [\mathrm{Im}(z)]^2 \geq [\mathrm{Im}(z)]^2 = \left[\frac{1}{2i}(z - z^*)\right]^2$

$\therefore$ letting $z = \langle f \mid g\rangle$,

Proof using Schwarz inequality :

Consider for an observable A :$\sigma_A^2 = \langle(\hat{A} - \langle A\rangle)\Psi|(\hat{A} - \langle A\rangle)\Psi\rangle = \langle f|f\rangle$, where $f \equiv \left(\hat{A} - \langle A\rangle\right)\Psi$

Likewise, for any other observable, B, $\sigma_B^2 = \langle g \mid g\rangle$, where $g \equiv \left(\hat{B} - \langle B\rangle\right)\Psi$

Schwarz inequality : $\sigma_A^2\sigma_B^2 = \langle f \mid f\rangle\langle g \mid g\rangle \geq |\langle f \mid g\rangle|^2$

For any complex number z,

$|z|^2 = [\mathrm{Re}(z)]^2 + [\mathrm{Im}(z)]^2 \geq [\mathrm{Im}(z)]^2 = \left[\frac{1}{2i}(z - z^*)\right]^2$

$\therefore$ letting $z = \langle f \mid g\rangle$, $\sigma_A^2\sigma_B^2 \geq \left(\frac{1}{2i}\left[\langle f \mid g\rangle - \langle g \mid f\rangle\right]\right)^2$

Exploiting the hermiticity of $\left(\hat{A} - \langle A \rangle\right)$

Exploiting the hermiticity of $\left(\hat{A} - \langle A \rangle\right)$

$$
\begin{aligned}
\langle f \mid g \rangle &= \left\langle \left(\hat{A} - \langle A \rangle\right) \Psi \mid \left(\hat{B} - \langle B \rangle\right) \Psi \right\rangle = \left\langle \Psi \mid \left(\hat{A} - \langle A \rangle\right) \left(\hat{B} - \langle B \rangle\right) \Psi \right\rangle \\
&= \left\langle \Psi \mid \left(\hat{A}\hat{B} - \hat{A}\langle B \rangle - \hat{B}\langle A \rangle + \langle A \rangle\langle B \rangle\right) \Psi \right\rangle \\
&= \left\langle \Psi \mid \hat{A}\hat{B}\Psi \right\rangle - \langle B \rangle \left\langle \Psi \mid \hat{A}\Psi \right\rangle - \langle A \rangle \left\langle \Psi \mid \hat{B}\Psi \right\rangle + \langle A \rangle\langle B \rangle\langle \Psi \mid \Psi \rangle \\
&= \langle \hat{A}\hat{B} \rangle - \langle B \rangle\langle A \rangle - \langle A \rangle\langle B \rangle + \langle A \rangle\langle B \rangle \\
&= \langle \hat{A}\hat{B} \rangle - \langle A \rangle\langle B \rangle
\end{aligned}
$$

Exploiting the hermiticity of $\left(\hat{A} - \langle A \rangle\right)$

$$
\begin{aligned}
\langle f \mid g \rangle &= \left\langle \left(\hat{A} - \langle A \rangle\right) \Psi \mid \left(\hat{B} - \langle B \rangle\right) \Psi \right\rangle = \left\langle \Psi \mid \left(\hat{A} - \langle A \rangle\right) \left(\hat{B} - \langle B \rangle\right) \Psi \right\rangle \\
&= \left\langle \Psi \mid \left(\hat{A}\hat{B} - \hat{A}\langle B \rangle - \hat{B}\langle A \rangle + \langle A \rangle\langle B \rangle\right) \Psi \right\rangle \\
&= \left\langle \Psi \mid \hat{A}\hat{B}\Psi \right\rangle - \langle B \rangle \left\langle \Psi \mid \hat{A}\Psi \right\rangle - \langle A \rangle \left\langle \Psi \mid \hat{B}\Psi \right\rangle + \langle A \rangle\langle B \rangle\langle \Psi \mid \Psi \rangle \\
&= \langle \hat{A}\hat{B} \rangle - \langle B \rangle\langle A \rangle - \langle A \rangle\langle B \rangle + \langle A \rangle\langle B \rangle \\
&= \langle \hat{A}\hat{B} \rangle - \langle A \rangle\langle B \rangle
\end{aligned}
$$

Similarly, $\langle g \mid f \rangle = \langle \hat{B}\hat{A} \rangle - \langle A \rangle\langle B \rangle$

Exploiting the hermiticity of $\left(\hat{A} - \langle A \rangle\right)$

$$
\begin{aligned}
\langle f \mid g \rangle &= \left\langle \left(\hat{A} - \langle A \rangle\right) \Psi \mid \left(\hat{B} - \langle B \rangle\right) \Psi \right\rangle = \left\langle \Psi \mid \left(\hat{A} - \langle A \rangle\right) \left(\hat{B} - \langle B \rangle\right) \Psi \right\rangle \\
&= \left\langle \Psi \mid \left(\hat{A}\hat{B} - \hat{A}\langle B \rangle - \hat{B}\langle A \rangle + \langle A \rangle\langle B \rangle\right) \Psi \right\rangle \\
&= \left\langle \Psi \mid \hat{A}\hat{B}\Psi \right\rangle - \langle B \rangle \left\langle \Psi \mid \hat{A}\Psi \right\rangle - \langle A \rangle \left\langle \Psi \mid \hat{B}\Psi \right\rangle + \langle A \rangle\langle B \rangle\langle \Psi \mid \Psi \rangle \\
&= \langle \hat{A}\hat{B} \rangle - \langle B \rangle\langle A \rangle - \langle A \rangle\langle B \rangle + \langle A \rangle\langle B \rangle \\
&= \langle \hat{A}\hat{B} \rangle - \langle A \rangle\langle B \rangle
\end{aligned}
$$

Similarly, $\langle g \mid f \rangle = \langle \hat{B}\hat{A} \rangle - \langle A \rangle\langle B \rangle$

$$
\langle f \mid g \rangle - \langle g \mid f \rangle = \langle \hat{A}\hat{B} \rangle - \langle \hat{B}\hat{A} \rangle = \langle [\hat{A}, \hat{B}] \rangle \implies \sigma_A^2 \sigma_B^2 \geq \left(\tfrac{1}{2i} \langle [\hat{A}, \hat{B}] \rangle \right)^2
$$

Exploiting the hermiticity of $\left(\hat{A} - \langle A \rangle\right)$

$$\begin{aligned}
\langle f \mid g \rangle &= \left\langle \left(\hat{A} - \langle A \rangle\right) \Psi \mid \left(\hat{B} - \langle B \rangle\right) \Psi \right\rangle = \left\langle \Psi \mid \left(\hat{A} - \langle A \rangle\right) \left(\hat{B} - \langle B \rangle\right) \Psi \right\rangle \\
&= \left\langle \Psi \mid \left(\hat{A}\hat{B} - \hat{A}\langle B \rangle - \hat{B}\langle A \rangle + \langle A \rangle\langle B \rangle\right) \Psi \right\rangle \\
&= \left\langle \Psi \mid \hat{A}\hat{B}\Psi \right\rangle - \langle B \rangle \left\langle \Psi \mid \hat{A}\Psi \right\rangle - \langle A \rangle \left\langle \Psi \mid \hat{B}\Psi \right\rangle + \langle A \rangle\langle B \rangle\langle \Psi \mid \Psi \rangle \\
&= \langle \hat{A}\hat{B} \rangle - \langle B \rangle\langle A \rangle - \langle A \rangle\langle B \rangle + \langle A \rangle\langle B \rangle \\
&= \langle \hat{A}\hat{B} \rangle - \langle A \rangle\langle B \rangle
\end{aligned}$$

Similarly, $\langle g \mid f \rangle = \langle \hat{B}\hat{A} \rangle - \langle A \rangle\langle B \rangle$

$\langle f \mid g \rangle - \langle g \mid f \rangle = \langle \hat{A}\hat{B} \rangle - \langle \hat{B}\hat{A} \rangle = \langle [\hat{A}, \hat{B}] \rangle \implies \sigma_A^2 \sigma_B^2 \geq \left(\frac{1}{2i}\langle [\hat{A}, \hat{B}] \rangle\right)^2$

Use $\hat{A} = x$ and $\hat{B} = -i\hbar\frac{d}{dx}$ and show : $\sigma_x \sigma_p \geq \frac{\hbar}{2}$