


GRAVITATIONAL LENSING

11 – MICROLENSING LIGHT CURVES: BREAKING THE MICROLENSING DEGENERACIES

Massimo Meneghetti
AA 2018-2019

EXAMPLE OF STANDARD LIGHT CURVE

.....



MICROLENSING DEGENERACY AND ITS BREAKERS

- As seen, from the standard light curve we can measure the Einstein crossing time, which is a degenerate combination of the lens mass, distance and velocity
- Thus it is impossible to characterise the lens and the source of a single event through light curve measurement alone...
- ... unless special circumstances are present!

FINITE SOURCE EFFECTS


- So far, we have assumed that source are point-like
- In practice, while the lens transits the surface of an extended source during the course of microlensing, this approximation is no longer valid
- The correct calculation of the source magnification in this case requires to integrate the magnification over the surface of the source:

$$\mu_{FS}(y | \rho_s) = \frac{\int_0^{2\pi} \int_0^{\rho_s} d\varphi dr r \mu \left(\sqrt{(y + r \cos \varphi)^2 + (r \sin \varphi)^2} \right)}{\int_0^{2\pi} \int_0^{\rho_s} d\varphi dr r}$$

$$\rho_s = \frac{\beta_*}{\theta_E}$$

FINITE SOURCE EFFECTS

Hamolli et al (2015)



We can fit the light curve with an additional parameter (ρ_s), and use some empirical relation to measure the source size from the source color and magnitude. For example, Kervella et al. (2004) find:

$$\log(2\beta_*) = 0.0755(V - K) + 0.517 - 0.2K$$

Then, we can infer the Einstein radius.

ASTROMETRIC MEASUREMENT OF THE RELATIVE PROPER MOTION

- Another method to measure the Einstein radius is by measuring the relative proper motion of source and lens.
- This requires to see the lens!
- If we have high resolution imaging with HST or ground based AO, we can measure the position of lens and source at some time after the maximum of the light curve Δt
- If we measure a shift $\Delta\theta$ then $\mu_{rel} = \Delta\theta/\Delta t$ and

$$\theta_E = t_E \times \mu_{rel}$$