


GRAVITATIONAL LENSING

3 – DEFLECTION ANGLE (CONTINUATION) – LENS EQUATION

Massimo Meneghetti
AA 2018-2019

GENERALISATION OF THE DEFLECTION ANGLE FORMULA


$$\hat{\vec{\alpha}}(\vec{\xi}) = \frac{4GM}{c^2 \xi} \vec{e}_\xi = \frac{4GM}{c^2} \frac{\vec{\xi}}{|\xi|^2}$$



Using “Thin screen approximation”

GENERALISATION OF THE DEFLECTION ANGLE FORMULA


$$\hat{\vec{\alpha}}(\vec{\xi}) = \frac{4GM}{c^2} \frac{\vec{\xi} - \vec{\xi}'}{|\vec{\xi} - \vec{\xi}'|^2}$$



Using “Thin screen approximation”

GENERALISATION OF THE DEFLECTION ANGLE FORMULA


$$\hat{\vec{\alpha}}(\vec{\xi}) = \frac{4G}{c^2} \sum_i M_i \frac{\vec{\xi} - \vec{\xi}_i'}{|\vec{\xi} - \vec{\xi}_i'|^2}$$



Using “Thin screen approximation”


DEFLECTION BY AN ENSEMBLE OF POINT MASSES

- Structure formation is often studied using numerical simulations
- Galaxies, galaxy clusters, etc. are described by ensembles of particles
- The calculation of the deflection angle by direct summation of all contributions from each particle has a computational cost $O(N^2)$



DEFLECTION BY AN ENSEMBLE OF POINT MASSES

- Structure formation is often studied using numerical simulations
- Galaxies, galaxy clusters, etc. are described by ensembles of particles
- The calculation of the deflection angle by direct summation of all contributions from each particle has a computational cost $O(N^2)$




DEFLECTION BY AN EXTENDED MASS DISTRIBUTION

- This can be easily generalized to the case of a continuum distribution of mass


$$\Sigma(\vec{\xi}) = \int \rho(\vec{\xi}, z) dz$$

$$d\hat{\vec{\alpha}}(\vec{\xi}) = \frac{4G}{c^2} \frac{\vec{\xi} - \vec{\xi}'}{|\vec{\xi} - \vec{\xi}'|^2} \Sigma(\vec{\xi}') d\xi'^2$$

$$\hat{\vec{\alpha}}(\vec{\xi}) = \frac{4G}{c^2} \int \frac{\vec{\xi} - \vec{\xi}'}{|\vec{\xi} - \vec{\xi}'|^2} \Sigma(\vec{\xi}') d\xi'^2$$



IS ANY DEFLECTION RIGHT?




Some rays are deflected in the right way, others are not!




IS ANY DEFLECTION RIGHT?

.....




Some rays are deflected in the right way, others are not!


LENS EQUATION




LENS EQUATION




LENS EQUATION




LENS EQUATION




LENS EQUATION




LENS EQUATION



LENS EQUATION







$$\hat{\eta} = D_{LS}\hat{\alpha}(\theta)$$

$$D_S \theta = D_S \beta + D_{LS} \hat{\alpha}(\theta) \Rightarrow \beta = \theta - \frac{D_{LS}}{D_S} \hat{\alpha}(\theta)$$

LENS EQUATION



LENS EQUATION



Remember that:

- 1) positions on the lens and source planes are defined by vectors
- 2) the deflection angle itself is a vector

$$\vec{\theta} = \frac{\vec{\xi}}{D_L} \quad \vec{\beta} = \frac{\vec{\eta}}{D_S}$$

$$\vec{\beta} = \vec{\theta} - \frac{D_{LS}}{D_S} \hat{\alpha}(\vec{\theta}) = \vec{\theta} - \vec{\alpha}(\vec{\theta})$$

DIMENSIONLESS NOTATION

Quite often, an alternative way is chosen to write the lens equation: the so called “**dimension-less**” notation.

This implies the choice of a *reference angle (or length)* to scale the source and image positions and the deflection angle:

$$\vec{\theta} = \frac{\vec{\xi}}{D_L} \quad \vec{\beta} = \frac{\vec{\eta}}{D_S} \quad \vec{\alpha}(\vec{\theta}) = \frac{D_{LS}}{D_S} \hat{\vec{\alpha}}(\vec{\theta}) \quad \vec{\beta} = \vec{\theta} - \frac{D_{LS}}{D_S} \hat{\vec{\alpha}}(\vec{\theta}) = \vec{\theta} - \vec{\alpha}(\vec{\theta})$$

$$\theta_0 = \frac{\xi_0}{D_L} = \frac{\eta_0}{D_S}$$

the reference angle subtends the reference scales on the lens and on the source planes

$$\frac{\vec{\theta}}{\theta_0} = \frac{\vec{\beta}}{\theta_0} - \frac{\vec{\alpha}(\vec{\theta})}{\theta_0}$$

dividing both members of the lens equation by the reference angle...

$$\vec{y} = \vec{x} - \vec{\alpha}(\vec{x})$$

$$\vec{\alpha}(\vec{x}) = \frac{\vec{\alpha}(\vec{\theta})}{\theta_0} = \frac{D_L}{\xi_0} \vec{\alpha}(\vec{\theta})$$

LENSING POTENTIAL

$$\hat{\vec{\alpha}} = \frac{2}{c^2} \int_{-\infty}^{+\infty} \vec{\nabla}_{\perp} \Phi dz$$

This formula tells us that the deflection is caused by the projection of the Newtonian gravitational potential on the lens plane.

$$\hat{\Psi}(\vec{\theta}) = \frac{D_{LS}}{D_L D_S} \frac{2}{c^2} \int \Phi(D_L \vec{\theta}, z) dz$$

*We introduce the **effective lensing potential***

LENSING POTENTIAL

$$\hat{\vec{\alpha}} = \frac{2}{c^2} \int_{-\infty}^{+\infty} \vec{\nabla}_{\perp} \Phi dz$$

This formula tells us that the deflection is caused by the projection of the Newtonian gravitational potential on the lens plane.

$$\hat{\Psi}(\vec{\theta}) = \frac{D_{LS}}{D_L D_S} \frac{2}{c^2} \int \Phi(D_L \vec{\theta}, z) dz$$

*We introduce the **effective lensing potential***

1

the lensing potential is the projection of the 3D potential

LENSING POTENTIAL

$$\hat{\vec{\alpha}} = \frac{2}{c^2} \int_{-\infty}^{+\infty} \vec{\nabla}_{\perp} \Phi dz$$

This formula tells us that the deflection is caused by the projection of the Newtonian gravitational potential on the lens plane.

$$\hat{\Psi}(\vec{\theta}) = \frac{D_{LS}}{D_L D_S} \frac{2}{c^2} \int \Phi(D_L \vec{\theta}, z) dz$$

*We introduce the **effective lensing potential***

1

the lensing potential is the projection of the 3D potential

2

the lensing potential scales with distances

OTHER PROPERTIES OF THE LENSING POTENTIAL

$$\vec{\nabla}_{\theta} \hat{\Psi}(\vec{\theta}) = \vec{\alpha}(\vec{\theta})$$

The reduced deflection angle is the gradient of the lensing potential

$$\begin{aligned} \vec{\nabla}_{\theta} \hat{\Psi}(\vec{\theta}) &= D_L \vec{\nabla}_{\perp} \hat{\Psi} = \vec{\nabla}_{\perp} \left(\frac{D_{LS}}{D_S} \frac{2}{c^2} \int \hat{\Phi}(\vec{\theta}, z) dz \right) \\ &= \frac{D_{LS}}{D_S} \frac{2}{c^2} \int \vec{\nabla}_{\perp} \Phi(\vec{\theta}, z) dz \\ &= \vec{\alpha}(\vec{\theta}) \end{aligned}$$

NOTE THAT...

... the same result holds if we use the dimension-less notation:

$$\vec{\nabla}_x = \frac{\xi_0}{D_L} \vec{\nabla}_\theta$$



$$\vec{\nabla}_x \hat{\Psi} = \frac{\xi_0}{D_L} \vec{\nabla}_\theta \hat{\Psi} = \frac{\xi_0}{D_L} \vec{\alpha}$$

By multiplying both sides of the equation by D_L^2/ξ_0^2 we obtain:

$$\frac{D_L^2}{\xi_0^2} \vec{\nabla}_x \hat{\Psi} = \frac{D_L}{\xi_0} \vec{\alpha} \quad \rightarrow \quad \Psi = \frac{D_L^2}{\xi_0^2} \hat{\Psi} \quad \rightarrow \quad \vec{\nabla}_x \Psi(\vec{x}) = \vec{\alpha}(\vec{x})$$

We have introduced the dimensionless counter-part of the lensing potential!

OTHER PROPERTIES OF THE LENSING POTENTIAL

$$\Delta_{\theta} \hat{\Psi}(\vec{\theta}) = 2\kappa(\vec{\theta})$$

The laplacian of the lensing potential is twice the convergence:

$$\kappa(\vec{\theta}) \equiv \frac{\Sigma(\vec{\theta})}{\Sigma_{\text{cr}}} \quad \text{with} \quad \Sigma_{\text{cr}} = \frac{c^2}{4\pi G} \frac{D_{\text{S}}}{D_{\text{L}} D_{\text{LS}}}$$

OTHER PROPERTIES OF THE LENSING POTENTIAL

$$\Delta_{\theta} \hat{\Psi}(\vec{\theta}) = 2\kappa(\vec{\theta})$$

The laplacian of the lensing potential is twice the convergence:

$$\kappa(\vec{\theta}) \equiv \frac{\Sigma(\vec{\theta})}{\Sigma_{\text{cr}}} \quad \text{with} \quad \Sigma_{\text{cr}} = \frac{c^2}{4\pi G} \frac{D_S}{D_L D_{LS}}$$

$$[G] = L^3/M/T^2$$

$$[c^2] = L^2/T^2$$

$$[D_X] = L$$

OTHER PROPERTIES OF THE LENSING POTENTIAL

$$\Delta_{\theta} \hat{\Psi}(\vec{\theta}) = 2\kappa(\vec{\theta})$$

The laplacian of the lensing potential is twice the convergence:

$$\kappa(\vec{\theta}) \equiv \frac{\Sigma(\vec{\theta})}{\Sigma_{\text{cr}}} \quad \text{with} \quad \Sigma_{\text{cr}} = \frac{c^2}{4\pi G} \frac{D_S}{D_L D_{LS}}$$

$$[G] = L^3/M/T^2$$

$$[c^2] = L^2/T^2$$

$$[D_X] = L$$

The critical surface density is a characteristic density to distinguish between strong and weak gravitational lenses!

OTHER PROPERTIES OF THE LENSING POTENTIAL

$$\Delta_{\theta} \hat{\Psi}(\vec{\theta}) = 2\kappa(\vec{\theta})$$

The laplacian of the lensing potential is twice the convergence:

We start from the poisson equation

$$\Delta \Phi = 4\pi G \rho$$

The surface mass density is then:

$$\Sigma(\vec{\theta}) = \frac{1}{4\pi G} \int_{-\infty}^{+\infty} \Delta \Phi dz$$

$$\kappa(\vec{\theta}) = \frac{1}{c^2} \frac{D_L D_{LS}}{D_S} \int_{-\infty}^{+\infty} \Delta \Phi dz$$

Let's introduce the Laplacian operator on the lens plane:

$$\Delta_{\theta} = \frac{\partial^2}{\partial \theta_1^2} + \frac{\partial^2}{\partial \theta_2^2} = D_L^2 \left(\frac{\partial^2}{\partial \xi_1^2} + \frac{\partial^2}{\partial \xi_2^2} \right) = D_L^2 \left(\Delta - \frac{\partial^2}{\partial z^2} \right)$$

Then:

$$\Delta \Phi = \frac{1}{D_L^2} \Delta_{\theta} \Phi + \frac{\partial^2 \Phi}{\partial z^2}$$

OTHER PROPERTIES OF THE LENSING POTENTIAL

With this substitution:

$$\kappa(\vec{\theta}) = \frac{1}{c^2} \frac{D_{LS}}{D_S D_L} \left[\Delta_\theta \int_{-\infty}^{+\infty} \Phi dz + D_L^2 \int_{-\infty}^{+\infty} \frac{\partial^2 \Phi}{\partial z^2} dz \right]$$


where the second term in the sum is zero, if the lens is gravitationally bound!

Given the definition of lensing potential:

$$\kappa(\theta) = \frac{1}{2} \Delta_\theta \hat{\Psi}$$

Note that:

$$\Delta_\theta = D_L^2 \Delta_\xi = \frac{D_L^2}{\xi_0^2} \Delta_x \quad \kappa(\theta) = \frac{1}{2} \Delta_\theta \hat{\Psi} = \frac{1}{2} \frac{\xi_0^2}{D_L^2} \Delta_\theta \Psi$$



$$\kappa(\vec{x}) = \frac{1}{2} \Delta_x \Psi(\vec{x})$$

DIMENSIONLESS NOTATION

From

$$\vec{\hat{\alpha}}(\vec{\xi}) = \frac{4G}{c^2} \int \frac{(\vec{\xi} - \vec{\xi}') \Sigma(\vec{\xi}')}{|\vec{\xi} - \vec{\xi}'|^2} d^2 \xi'$$

we obtain

$$\vec{\alpha}(\vec{x}) = \frac{1}{\pi} \int_{\mathbf{R}^2} d^2 x' \kappa(\vec{x}') \frac{\vec{x} - \vec{x}'}{|\vec{x} - \vec{x}'|}$$

Using

$$\vec{\nabla}_x \Psi(\vec{x}) = \vec{\alpha}(\vec{x})$$

$$\Psi(\vec{x}) = \frac{1}{\pi} \int_{\mathbf{R}^2} \kappa(\vec{x}') \ln |\vec{x} - \vec{x}'| d^2 x'$$

DIMENSIONLESS NOTATION

From

$$\vec{\hat{\alpha}}(\vec{\xi}) = \frac{4G}{c^2} \int \frac{(\vec{\xi} - \vec{\xi}') \Sigma(\vec{\xi}')}{|\vec{\xi} - \vec{\xi}'|^2} d^2 \xi'$$

we obtain

$$\vec{\alpha}(\vec{x}) = \frac{1}{\pi} \int_{\mathbf{R}^2} d^2 x' \kappa(\vec{x}') \frac{\vec{x} - \vec{x}'}{|\vec{x} - \vec{x}'|}$$

Using

$$\vec{\nabla}_x \Psi(\vec{x}) = \vec{\alpha}(\vec{x})$$

$$\Psi(\vec{x}) = \frac{1}{\pi} \int_{\mathbf{R}^2} \kappa(\vec{x}') \ln |\vec{x} - \vec{x}'| d^2 x'$$

Convolution kernels

