

GRAVITATIONAL LENSING

LECTURE 19

Docente: Massimo Meneghetti
AA 2016-2017

EXTENDED LENSES

- Cosmic structures like galaxies and galaxy clusters are characterized by bound mass distributions, which cannot be approximated by point lenses
- Indeed these are *extended lenses*, and their lensing properties are determined by e.g. their surface mass density:

$$\Sigma(\vec{\xi}) = \int \rho(\vec{\xi}, z) \, dz$$

$$\vec{\alpha}(\vec{\xi}) = \frac{4G}{c^2} \int \frac{(\vec{\xi} - \vec{\xi}') \Sigma(\vec{\xi}')}{|\vec{\xi} - \vec{\xi}'|^2} \, d^2\xi'$$

EXTENDED LENSES

- Recall that the surface density is related to the lensing potential by

$$\Delta_{\theta} \Psi(\vec{\theta}) = 2\kappa(\vec{\theta})$$

$$\kappa(\vec{\theta}) \equiv \frac{\Sigma(\vec{\theta})}{\Sigma_{\text{cr}}} \quad \text{with} \quad \Sigma_{\text{cr}} = \frac{c^2}{4\pi G} \frac{D_S}{D_L D_{LS}}$$

WHAT ARE THE RELEVANT PROPERTIES OF THE LENSES?

- The surface density distribution of a lens (and its potential) can be characterized by means of
 - the profile
 - the shape of the iso-density (iso-potential) contours
 - the smoothness
 - the environment where the lens resides
- In this and in the following lessons, we will study how these features determine the ability of a mass distribution to produce lensing effects.
- We will do that by building analytical models with increasing level of complexity.


AXIALLY SYMMETRIC, CIRCULAR LENSES

- Axially symmetric, circular models are the simplest lens models for describing extended mass distributions
- For these lenses $\Sigma(\vec{\xi}) = \Sigma(|\vec{\xi}|)$
- Several quantities relevant for lensing can be derived in a simple manner by using the symmetry properties of the lens.
- One example is the deflection angle...

DEFLECTION ANGLE OF AN AXIALLY SYMMETRIC LENS

$$\vec{\alpha}(\vec{\xi}) = \frac{4G}{c^2} \int \frac{(\vec{\xi} - \vec{\xi}') \Sigma(\xi')}{|\vec{\xi} - \vec{\xi}'|^2} d^2 \xi'$$

$$\begin{aligned} \vec{\xi} - \vec{\xi}' &= (\xi - \xi' \cos \phi, -\xi' \sin \phi) \\ |\vec{\xi} - \vec{\xi}'|^2 &= \xi^2 + \xi'^2 \cos^2 \phi - 2\xi\xi' \cos \phi + \xi'^2 \sin^2 \phi \\ &= \xi^2 + \xi'^2 - 2\xi\xi' \cos \phi \end{aligned}$$



$$\begin{aligned} \hat{\alpha}_1(\vec{\xi}) &= \frac{4G}{c^2} \int_0^\infty d\xi' \xi' \Sigma(\xi') \int_0^{2\pi} d\phi \frac{\xi - \xi' \cos \phi}{\xi^2 + \xi'^2 - 2\xi\xi' \cos \phi} \\ \hat{\alpha}_2(\vec{\xi}) &= \frac{4G}{c^2} \int_0^\infty d\xi' \xi' \Sigma(\xi') \int_0^{2\pi} d\phi \frac{-\xi' \sin \phi}{\xi^2 + \xi'^2 - 2\xi\xi' \cos \phi} \end{aligned}$$


$$\hat{\alpha}(\xi) = \frac{4G}{c^2} \frac{2\pi \int_0^\xi \Sigma(\xi') \xi' d\xi'}{\xi} = \frac{4GM(\xi)}{c^2 \xi}$$

For an axially symmetric lens, the deflection is “radial”: it depends only on the distance from the lens center.

DEFLECTION ANGLE OF AN AXIALLY SYMMETRIC LENS

$$\vec{\alpha}(\vec{\xi}) = \frac{4G}{c^2} \int \frac{(\vec{\xi} - \vec{\xi}') \Sigma(\xi')}{|\vec{\xi} - \vec{\xi}'|^2} d^2 \xi'$$

$$\begin{aligned} \vec{\xi} - \vec{\xi}' &= (\xi - \xi' \cos \phi, -\xi' \sin \phi) \\ |\vec{\xi} - \vec{\xi}'|^2 &= \xi^2 + \xi'^2 \cos^2 \phi - 2\xi\xi' \cos \phi + \xi'^2 \sin^2 \phi \\ &= \xi^2 + \xi'^2 - 2\xi\xi' \cos \phi \end{aligned}$$



$$\begin{aligned} \hat{\alpha}_1(\vec{\xi}) &= \frac{4G}{c^2} \int_0^\infty d\xi' \xi' \Sigma(\xi') \int_0^{2\pi} d\phi \frac{\xi - \xi' \cos \phi}{\xi^2 + \xi'^2 - 2\xi\xi' \cos \phi} \\ \hat{\alpha}_2(\vec{\xi}) &= \frac{4G}{c^2} \int_0^\infty d\xi' \xi' \Sigma(\xi') \int_0^{2\pi} d\phi \frac{-\xi' \sin \phi}{\xi^2 + \xi'^2 - 2\xi\xi' \cos \phi} \end{aligned}$$

$$\frac{2\pi}{\xi}$$

$$\xi' < \xi$$

$$\hat{\alpha}(\xi) = \frac{4G}{c^2} \frac{2\pi \int_0^\xi \Sigma(\xi') \xi' d\xi'}{\xi} = \frac{4GM(\xi)}{c^2 \xi}$$

For an axially symmetric lens, the deflection is “radial”: it depends only on the distance from the lens center.

DEFLECTION ANGLE IN ADIMENSIONAL FORM

$$\hat{\alpha}(\xi) = \frac{4G}{c^2} \frac{2\pi \int_0^\xi \Sigma(\xi') \xi' d\xi'}{\xi} = \frac{4GM(\xi)}{c^2 \xi}$$

*The adimensional,
reduced deflection
angle is:*

$$\begin{aligned} \alpha(x) &= \frac{D_L D_{LS}}{\xi_0 D_S} \hat{\alpha}(\xi_0 x) \\ &= \frac{D_L D_{LS}}{\xi_0 D_S} \frac{4GM(\xi_0 x)}{c^2 \xi} \frac{\pi \xi_0}{\pi \xi_0} \\ &= \frac{M(\xi_0 x)}{\pi \xi_0^2 \Sigma_{\text{cr}}} \frac{1}{x} \equiv \frac{m(x)}{x}, \end{aligned}$$

We have introduced the dimensionless mass:

$$\alpha(x) = \frac{2}{x} \int_0^x x' \kappa(x') dx' \Rightarrow m(x) = 2 \int_0^x x' \kappa(x') dx' .$$

LENS EQUATION

$$\vec{y} = \vec{x} - \vec{\alpha}(\vec{x}) \qquad \vec{\alpha}(\vec{x}) = \frac{m(\vec{x})}{x^2} \vec{x}$$

Given that the deflection angle and x are parallel, so will be y !

$$y = x - \frac{m(x)}{x}$$

CONVERGENCE AND SHEAR

To compute convergence and shear, we need to compute the partial derivatives of the deflection angle:

$$\frac{\partial \alpha_1}{\partial x_1} = \frac{dm}{dx} \frac{x_1^2}{x^3} + m \frac{x_2^2 - x_1^2}{x^4},$$

$$\frac{\partial \alpha_2}{\partial x_2} = \frac{dm}{dx} \frac{x_2^2}{x^3} + m \frac{x_1^2 - x_2^2}{x^4},$$

$$\frac{\partial \alpha_1}{\partial x_2} = \frac{\partial \alpha_2}{\partial x_1} = \frac{dm}{dx} \frac{x_1 x_2}{x^3} - 2m \frac{x_1 x_2}{x^4}$$

CONVERGENCE AND SHEAR

From the partial derivatives of the deflection angle we obtain:

$$\kappa(x) = \frac{1}{2x} \frac{dm(x)}{dx},$$

$$\gamma_1(x) = \frac{1}{2}(x_2^2 - x_1^2) \left(\frac{2m(x)}{x^4} - \frac{dm(x)}{dx} \frac{1}{x^3} \right)$$

$$\gamma_2(x) = x_1 x_2 \left(\frac{dm(x)}{dx} \frac{1}{x^3} - \frac{2m(x)}{x^4} \right).$$

CONVERGENCE AND SHEAR

For an axially symmetric lens the following relation exists between convergence and shear:

$$\gamma(x) = \frac{m(x)}{x^2} - \kappa(x)$$

$$\frac{m(x)}{x^2} = 2\pi \frac{\int_0^x x' \kappa(x') dx'}{\pi x^2} = \bar{\kappa}(x)$$

$$\gamma(x) = \bar{\kappa}(x) - \kappa(x)$$

LENSING JACOBIAN

$$A = I - \frac{m(x)}{x^4} \begin{pmatrix} x_2^2 - x_1^2 & -2x_1x_2 \\ -2x_1x_2 & x_1^2 - x_2^2 \end{pmatrix} - \frac{dm(x)}{dx} \frac{1}{x^3} \begin{pmatrix} x_1^2 & x_1x_2 \\ x_1x_2 & x_2^2 \end{pmatrix}.$$

DETERMINANT OF THE LENSING JACOBIN

$$\begin{aligned}\det A &= \frac{y}{x} \frac{dy}{dx} = \left(1 - \frac{m(x)}{x^2}\right) \left[1 - \frac{d}{dx} \left(\frac{m(x)}{x}\right)\right] \\ &= \left(1 - \frac{m(x)}{x^2}\right) \left(1 + \frac{m(x)}{x^2} - 2\kappa(x)\right) \\ &= \left(1 - \frac{\alpha(x)}{x}\right) \left(1 - \frac{d\alpha(x)}{dx}\right) .\end{aligned}$$

CRITICAL LINES

The critical lines can be found by setting $\det A = 0$:

$$m(x)/x^2 = 1$$

$$d(m(x)/x)/dx = 1$$

The first defines the “tangential” critical line. The second the “radial” critical line.

If they exist, what is the shape of the critical lines for this kind of lenses? What are the caustics?