Modeling Relationship between Alcohol Policy Perception and Alcohol Consumption in '01 Harvard College Alcohol Survey (CAS)

Youngsoo Baek, Michael Christensen, and Yufeng Jiang

Objective


- ▶ Data: 2001 Harvard College Alcohol Study (CAS)
 - ▶ 10904 participants (unknown response rate)
- Investigate the correlation structure between subjective beliefs about campus alcohol policy and objective measures of alcohol consumption
- ► *Section B* for subjective questions, *Section C* for objective questions
- ► Standard survey modeling techniques: factor analysis, structural equations model, item response theory

Data Processing

- Consistent ordering of responses
 - ▶ More stringent alcohol policy beliefs (1) -> Less stringent
 - Less alcohol consumption (1) -> More consumption
- Aggressive pruning of the variables before modeling
- Unreliable responses classified based on
 - Response to A7: A (alone) is not allowed with other responses (family/partner/roommate)
 - Response in Section C: participants who chose 1 in C10 and answered C11–C15, etc.
- WEIGHT01 used as sampling weights (intended for cross-sectinoal studies)

Missing Reponses

 Missing response rate adjusted for questions that only target certain demographic subgroups



Missing Responses for Each Section

Structural Equations Modeling (SEM)

- Survey responses X_i can be grouped together as repeated measurements of a lower-dimensional, latent factors: alcohol beliefs, alcohol consumption attitude, . . .
- ► Factor analysis identifies the loadings Λ of latent variable η_i .

$$X_i = \Lambda \eta_i + \epsilon_i, \ \eta_i \sim N(0,1), \ \epsilon_i \sim N(0,1)$$

- Structural Equations Models extend factor analysis by specifying within-question correlations and regressing η_i on predictors.
 - ► All of our model predictors are directly observed rather than "manifested" by questions: age, gender, . . .
- Causal interpretation is not necessary (though often made!).

Graphical Representation of the Model

...PLOT...

Modeling Challenges

- ► The model is clearly misspecified: Gaussian error assumption is made on ordered response
 - Asymptotic standard errors of factor loading estimators are valid for nonnormal factor analyses (Anderson and Amemiya, 1988)
 - In practice can cause lower goodness-of-fit
- Complete case analysis due to excessive computation in maximizing the full likelihood
- ► (ANYTHING ELSE??)

Main Results

A MATRIX PLOT OF LOADINGS (FACTOR CORRELATION) WILL BE GREAT HERE, RATHER THAN ANY ESTIMATE FIGURES . . .

Model Diagnostics

- ▶ Various statistics to evaluate model fit in practice: TLI, BL89, CFI, RMSEA (Hu and Bentler, 1999)
- ► (INCLUDE RESULTS HERE ...)

Interpretation

Conclusion

- Limitations
 - ► Alternative approaches to account for ordered response
 - ▶ Theory-driven priors may improve fit of more complex models
 - ▶ Need information to correct for estimate biases

Reference

- "Asymptotic Chi-Square Tests for a Large Class of Factor Analysis Models," Anderson, T. W. and Amemiya, Y. The Annals of Statistics, 16(2), 1988.
- "Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives," Hu, L.-T. and Bentler, P. M. Structural Equation Modeling: A Multidisciplinary Journal, 6(1), 1999.