Case Study 1-Group 1

Melody Jiang, Irene Ji, Keru Wu

Department of Statistical Science, Duke University

01/21/2019

Introduction

- ▶ Data: A study by Longnecker et al. (2001), comprised of 2380 observations of pregnant women.
- Goal: Assess how DDE and PCBs relate to risk of premature delivery.

▶ Preterm birth: Gestational Age ≤ 36.

- ▶ Preterm birth: Gestational Age ≤ 36.
- Standardize continuous variables.

- ▶ Preterm birth: Gestational Age ≤ 36.
- Standardize continuous variables.
- Missing data: Multivariate Imputations by Chained Equations (MICE package in R) for covariates. Only one missing in continuous, deleted.

- ▶ Preterm birth: Gestational Age ≤ 36.
- Standardize continuous variables.
- Missing data: Multivariate Imputations by Chained Equations (MICE package in R) for covariates. Only one missing in continuous, deleted.
- ▶ Limit of Detection (LOD): Exists in some PCBs. All LODs are negligible compared to data scale (e.g. 0.01 compared to 0.3)

Collinearity & Dimensionality Reduction

► There are 11 types of PCBs, some of which have high correlation and may be redundant.

Collinearity & Dimensionality Reduction

- ► There are 11 types of PCBs, some of which have high correlation and may be redundant.
- Possible approaches: Simple sum, PCA, Factor Analysis.

Collinearity & Dimensionality Reduction

- ► There are 11 types of PCBs, some of which have high correlation and may be redundant.
- ▶ Possible approaches: Simple sum, PCA, Factor Analysis.
- Check Multicollinearity: Variation Inflation Factor (VIF).

► Linear regression or logistic regression?

- ► Linear regression or logistic regression?
- ► Too simple & Can't fit nonlinear trend.

- ► Linear regression or logistic regression?
- ► Too simple & Can't fit nonlinear trend.
- Domain knowledge?

- ► Linear regression or logistic regression?
- ► Too simple & Can't fit nonlinear trend.
- ► Domain knowledge?
- Chemicals have no effect when concentration is lower than a bound.

- Linear regression or logistic regression?
- ► Too simple & Can't fit nonlinear trend.
- Domain knowledge?
- Chemicals have no effect when concentration is lower than a bound.
- Chemicals have constant effect when concentration is higher than a bound.

- Linear regression or logistic regression?
- ► Too simple & Can't fit nonlinear trend.
- Domain knowledge?
- Chemicals have no effect when concentration is lower than a bound.
- Chemicals have constant effect when concentration is higher than a bound.
- Nonlinear Model

Generalized Additive Model (GAM)

$$g(Y_i) = \beta_0 + \sum_{j=1}^m f_i(x_{ij}) + \sum_{k=1}^l \beta_k z_{ik}$$

- Choice of g: probit or logit.
- x_{.j}s include DDE, PCBs, maternal age, etc.
- z_{.k}s include categorical variables and some confounding variables.

- Frequentist approach may overestimate uncertainty.
- ► Frequentist GAM may produce a non-significant p-value.

- Frequentist approach may overestimate uncertainty.
- Frequentist GAM may produce a non-significant p-value.
- Bayesian Generalized Additive Model

$$g(Y_i) = \beta_0 + \sum_{j=1}^m f_j(x_{ij}) + \sum_{k=1}^l \beta_k z_{ik}$$

▶ Adds priors on the common regression coefficients, priors on the standard deviations of the smooth terms.

► Deal with different centers

- Deal with different centers
- ► Approach 1: Bayesian Hierarchical Model

- Deal with different centers
- ► Approach 1: Bayesian Hierarchical Model
- ► Approach 2: Mixed Effect / Random Effect Model

- Deal with different centers
- ► Approach 1: Bayesian Hierarchical Model
- Approach 2: Mixed Effect / Random Effect Model
- Generalized Additive Mixed Model (GAMM)
- Bayesian GAMM

► Specialized prior may give narrower credible intervals.

- ▶ Specialized prior may give narrower credible intervals.
- ▶ Including Interactions: Bayesian Factor Analysis (Ferrari, F. and Dunson, D.B. 2019)