

## Case Study 1

Emily Gentles    Phuc Nguyen    Joseph Lawson

Jan 21, 2020

## Case Discussion

- ▶ Data obtained from a subset of women enrolled in the CPP during pregnancy

**Goal:** Assess how exposure to DDE and PCBs relates to the risk of premature delivery and the relative severity thereof

# Exploratory Data Analysis








Figure 1: corr-plot


► Data issues: uncertainty, inflation, and missingness














**Severely Preterm Probability vs Maternal Age**



# Analysis

- ▶ Ordinal Logistic Regression with Term, Preterm, and Severely Preterm Gest. Categories
  - ▶ Useful interpretation in terms of risk
  - ▶ Uses naturally ordinal structure
  - ▶ Differentiates between different severities
- ▶ Impute score data with MICE to check usefulness
- ▶ Remove obs with missing PCB value
- ▶ Include blood cholesterol/triglyceride levels, as well as center and SES/Lifestyle metrics
- ▶ Use PCA to handle the various PCA measures.

## Model Comparison F-Tests

- ▶ Model Comparison indicated ( $p=0.97$ ) that the first principle component of the  $pcb\_*$  values is sufficient.
- ▶ Indication ( $p=0.74$ ) against including Score Variables (post imputation)
- ▶ Indication ( $p=0.21$ ) against including Center interactions
- ▶ Indication ( $p=0$ ) for including Center as variable
  - ▶ Indicates heterogeneity in preterm birth risk accross medical centers

- ▶ Indication ( $p=0.46$ ) against PCB-DDE interaction effect
- ▶ Indication ( $p=0.15$ ) (weakly) against Triglyceride interaction with PCE/DDE
  - ▶ Hypothesized that there might be because of fat-solubility of the contaminants of interest
- ▶ Indication ( $p=0.02$ ) for inclusion of quadratic term in maternal age
  - ▶ Makes sense given above chart/intuition of increased risk at extremes of age.
- ▶ Strong indication that the indicator of testing for Albumin is associated with longer gestational period on the margin

- ▶ Control Variables of Significance (marginally) in Final Model:
  - ▶ Center
  - ▶ Triglycerides (g/dL)
  - ▶ Cholesterol (g/dL)
  - ▶ Maternal Age (years)
  - ▶ Albumin (Testing Indicator)
  - ▶ Race

## Final Model

$$GestCategory \sim Center + \log(dde) + \log PCB1 \quad (1)$$

$$+ \log(trigl.) + Poly(MaternalAge, 2) \quad (2)$$

$$+ Smoking + \log(Cholest.) + AlbuminTested \quad (3)$$

$$+ race \quad (4)$$

## Results


- ▶ (log) DDE and PCB are (collectively) significantly associated with preterm delivery likelihood even when adjusting for other factors
- ▶ Interpretation that 1% decrease in DDE is approximately associated with a -0.18% change, and a unit increase in logPCB1 with a -0.06 change, in the log odds of of a gestational age occurrence above a given ordinal threshold.
  - ▶ Note for interpretability that the PCB coefficient is relative to the (standardized/centered) first principle component of the pcb\_\* measurements
- ▶ Testing inclusion of both DDE and PCB vs control indicate ( $p=0.005$ ) that they provide increased explanatory power vs control, but marginally in the full model each is just at the edge of significance, as seen in the following table:

	Coef Est	2.5 %	97.5 %
log(dde)	-0.183	-0.388	0.022
logPCB1	-0.056	-0.113	0.000


- Note the overlap of the confidence intervals with 0.



Fitted Probabilities vs DDE (ug/dL)/PCB (princ comp.) (log scale)



# Sensitivity Analysis



# Sensitivity Analysis

