

An Investigation of Social Factors Influencing Emergency Healthcare Expenditure

Maggie Lundberg, Riya Mohan, Izzy Kjaerulff

Contents

Abstract	1
Background & Significance	2
Methods	2
Data Collection	2
Exploratory Data Analysis	3
References	15

Abstract

This paper provides insight into an open investigation of personal healthcare spending in the United States, with focus on the emergency department (ED). We used data from the Disease Expenditure Project (DEX) at IHME, giving estimates for ED spending split into three major groups: public insurance, private insurance, and out of pocket. The data compiled is used to investigate the existence of a relationship between demographic (sex and age), disease, and expenditure. In the following report, we hypothesize demographic, disease, and type of expenditure from our three included groups are not independent of one another. For example, we expect that results will suggest that public healthcare spending will be higher in older populations when compared to younger populations, whereas private spending will be higher in those younger populations. From the analysis devised in this report, we gather that the data is consistent with the following conclusions: [INSERT CONCLUSIONS HERE]

Background & Significance

Emergency services ensure that individuals can receive timely care for unexpected ailments and injuries, making them a vital component of the healthcare industry. In recent years, however, emergency service spending has seen a significant increase (Scott and Liu 2021), which begs the question of equal accessibility. Expenditure is one of many ways by which to investigate interactions between demographic and healthcare access. [ELABORATE] As a preliminary piece of evaluation of the question of healthcare equity and accessibility by disease and demographic, we have prepared an analysis report of spending habits divided into the payer categories of public insurance, private insurance, and out of pocket. We hope to dive deeper into the relationship between spending habits and demographic through the lens of factors influencing payment models for the emergency department.

Methods


Data Collection

Our data is provided by the Institute of Health Metrics and Evaluation as part of the Disease Expenditure Project (DEX). These Emergency Department (ED) health spending data include estimates for U.S. spending on health care divided into three types of payers: public insurance (including Medicare, Medicaid, and other government programs), private insurance, and out-of-pocket payments. This dataset contains ED spending estimates by aggregate health category, age group, sex, and payer for 2006 through 2016, released in October 2021. Data were gathered from “government budgets, insurance claims, facility records, household surveys, and official US records” (IHME 2021). The data collection and agglomeration is funded by the National Institute on Aging (NIA) and the National Institutes of Health (NIH), and estimates were generated from an underlying data set—the National Emergency Department Sample (NEDS).

The data given includes summaries of identified gender and ages as “Both” and “All Ages” observations, respectively. In order to gauge accurate analysis of this data, we chose to exclude the aforementioned observations to avoid double counting.


It is important to acknowledge that this data set did not specify whether gender observations are based on individual reporting or otherwise observed identification and are limited to male and female. Therefore, the data may not encompass a complete representation of the population.

Normal Distribution of Payer Groups



The normal distribution for public spending, private spending, and out-of-pocket spending all show a severe right skew in the data. Therefore, all three variables do not meet the normal distribution assumption needed for many tests, such as ANOVA; however, this can easily be resolved by applying a log transformation to the data to give a fairly normal distribution of the data.

Normal Distribution of Payer Groups, Ln Applied



These graphs of the log distribution of the various spending means appear to be fairly normal in distribution, which means they meet the requirements to be used in various analyses. In order to convert to log scale, those with mean_all, mean_pub, mean_pri, and mean_oop equal to zero must be excluded.

Exploratory Data Analysis

Gender Our first question in this analysis is if males and females spend a different amount of money on emergency services.

First this t-test looks at overall differences in log mean emergency department spending between males and females

```
##
##  Welch Two Sample t-test
##
## data:  spending_malefemale$lmean_all by spending_malefemale$sex
## t = 1.1519, df = 5865.2, p-value = 0.2494
## alternative hypothesis: true difference in means between group Female and group Male is
## 95 percent confidence interval:
## -0.04706281  0.18118093
## sample estimates:
## mean in group Female    mean in group Male
##          17.88067          17.81361
```

This t-test shows that for mean spending of all emergency services payment types, the p value of 0.1543 (95% CI -0.0315862, 0.1996079) indicates there is not a significant difference between male and female spending.

Next, we perform a t-test on each type of insurance to see if there is a difference in spending between males and females:

```
##
##  Welch Two Sample t-test
##
## data:  spending_malefemale$lmean_pub by spending_malefemale$sex
## t = 1.5223, df = 5842.9, p-value = 0.128
## alternative hypothesis: true difference in means between group Female and group Male is
## 95 percent confidence interval:
## -0.02456331  0.19525433
## sample estimates:
## mean in group Female    mean in group Male
##          17.22124          17.13589
```

The t-test on emergency services spending for people who have public insurance indicates there is not a significant difference between male and female spending, with p value of 0.0697 (95% CI -0.00833746, 0.21532602).

```
##
##  Welch Two Sample t-test
```

```
##
## data: spending_malefemale$lmean_pri by spending_malefemale$sex
## t = 0.47771, df = 5906.8, p-value = 0.6329
## alternative hypothesis: true difference in means between group Female and group Male is
## 95 percent confidence interval:
## -0.09696189 0.15944338
## sample estimates:
## mean in group Female mean in group Male
## 16.63015 16.59891
```

The t-test on emergency services spending for people who have private insurance indicates there is not a significant difference between male and female spending, with p value of 0.4803 (95% CI -0.08283085, 0.17603825).

```
##
## Welch Two Sample t-test
##
## data: spending_malefemale$lmean_oop by spending_malefemale$sex
## t = 0.73927, df = 5876.1, p-value = 0.4598
## alternative hypothesis: true difference in means between group Female and group Male is
## 95 percent confidence interval:
## -0.07578874 0.16755600
## sample estimates:
## mean in group Female mean in group Male
## 14.46838 14.42250
```

The t-test on emergency services spending for people who pay out of pocket indicates there is not a significant difference between male and female spending, with p value of 0.3272 (95% CI -0.0615859, 0.1846904).

The t-tests for each type of insurance indicate that there is not enough evidence to reject the null hypothesis that emergency department spending is the same for males and females who have public insurance, private insurance, or pay out of pocket, leading us to the conclusion that gender does not influence emergency spending in the forms of payment studied here.

Disease In order to determine emergency department spending based on disease type, an ANOVA test is performed due to the data for spending on the log scale being normally distributed, relatively similar variance, and independent.

The null hypothesis for this ANOVA test is that the overall mean of spending are the same for each disease category

```
##          Df Sum Sq Mean Sq F value Pr(>F)
## agg_cause  14  18795   1342.5    639.5 <2e-16 ***
## Residuals 6035  12670     2.1
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Based on the p-value here of <2e-16, these data or more extreme data it is highly unlikely the null hypothesis is true. Therefore, we perform step-down tests using a Holm correction for multiple comparisons.

```
## [1] 95
```

The step-down t tests indicate 92 disease category pairs are different out of 105, indicating most disease categories do differ in the amount of government spending by the emergency department. There is lots of variation!!

```
## # A tibble: 15 x 5
##   term                estimate std.error statistic  p.value
##   <chr>              <dbl>    <dbl>    <dbl>    <dbl>
## 1 (Intercept)        17.5      0.0709    247.      0
## 2 agg_causeCancers   -2.52     0.100   -25.2  6.85e-133
## 3 agg_causeCardiovascular diseases  1.28     0.100    12.8  5.80e- 37
## 4 agg_causeChronic respiratory diseases  1.43     0.102    14.1  1.97e- 44
```


```
## 5 agg_causeCommunicable and nutrition d~ 1.76 0.100 17.6 1.01e- 67
## 6 agg_causeDiabetes and kidney diseases -0.0968 0.100 -0.966 3.34e- 1
## 7 agg_causeDigestive diseases 1.98 0.100 19.8 2.74e- 84
## 8 agg_causeEndocrine disorders -1.12 0.100 -11.2 9.89e- 29
## 9 agg_causeInjuries 2.71 0.100 27.0 6.31e-152
## 10 agg_causeMaternal and neonatal condit~ -3.47 0.121 -28.7 1.92e-170
## 11 agg_causeMusculoskeletal conditions 1.07 0.100 10.6 3.52e- 26
## 12 agg_causeNeurological disorders 0.994 0.100 9.92 5.01e- 23
## 13 agg_causeOther non-communicable disea~ 1.62 0.100 16.1 1.89e- 57
## 14 agg_causePrevention and coordination -3.00 0.100 -29.9 8.05e-184
## 15 agg_causeSkin and other sense organ d~ 1.12 0.100 11.2 9.17e- 29
```

Table 1: Temp

term	estimate	p.value
(Intercept)	17.4802309	0.0000000
agg_causeCancers	-2.5213164	0.0000000
agg_causeCardiovascular diseases	1.2816083	0.0000000
agg_causeChronic respiratory diseases	1.4324606	0.0000000
agg_causeCommunicable and nutrition disorders	1.7648936	0.0000000
agg_causeDiabetes and kidney diseases	-0.0968248	0.3340405
agg_causeDigestive diseases	1.9807183	0.0000000
agg_causeEndocrine disorders	-1.1203911	0.0000000
agg_causeInjuries	2.7084273	0.0000000
agg_causeMaternal and neonatal conditions	-3.4694399	0.0000000
agg_causeMusculoskeletal conditions	1.0657868	0.0000000
agg_causeNeurological disorders	0.9944291	0.0000000
agg_causeOther non-communicable diseases	1.6182974	0.0000000
agg_causePrevention and coordination	-3.0012823	0.0000000
agg_causeSkin and other sense organ disorders	1.1210773	0.0000000

```
glance(meanalldiseasecatfit)$adj.r.squared
```

```
## [1] 0.596407
```



```
## # A tibble: 30 x 5
##   term                                estimate std.error statistic  p.value
##   <chr>                                <dbl>     <dbl>     <dbl>    <dbl>
## 1 (Intercept)                        17.2      0.235     73.1      0
## 2 as.factor(year_id)2016              0.544     0.333      1.63  1.02e- 1
## 3 agg_causeCancers                    -2.54     0.333     -7.64  4.98e-14
## 4 agg_causeCardiovascular diseases    1.38     0.333      4.16  3.37e- 5
## 5 agg_causeChronic respiratory diseases 1.40     0.337      4.17  3.33e- 5
## 6 agg_causeCommunicable and nutrition di~ 1.79     0.333      5.40  8.32e- 8
## 7 agg_causeDiabetes and kidney diseases -0.0843   0.333     -0.254 8.00e- 1
## 8 agg_causeDigestive diseases         1.96     0.333      5.90  4.87e- 9
## 9 agg_causeEndocrine disorders        -1.26     0.333     -3.78  1.68e- 4
## 10 agg_causeInjuries                  2.78     0.333      8.35  2.07e-16
## # ... with 20 more rows
```

Table 2: Temp

term	estimate	p.value
(Intercept)	17.1841515	0.0000000
as.factor(year_id)2016	0.5435012	0.1024311
agg_causeCancers	-2.5387252	0.0000000
agg_causeCardiovascular diseases	1.3846396	0.0000337
agg_causeChronic respiratory diseases	1.4047353	0.0000333
agg_causeCommunicable and nutrition disorders	1.7946444	0.0000001
agg_causeDiabetes and kidney diseases	-0.0843353	0.7998236
agg_causeDigestive diseases	1.9618318	0.0000000

term	estimate	p.value
agg_causeEndocrine disorders	-1.2554208	0.0001683
agg_causeInjuries	2.7768257	0.0000000
agg_causeMaternal and neonatal conditions	-3.2613820	0.0000000
agg_causeMusculoskeletal conditions	1.0383789	0.0018387
agg_causeNeurological disorders	0.9448387	0.0045739
agg_causeOther non-communicable diseases	1.6312963	0.0000011
agg_causePrevention and coordination	-2.8957096	0.0000000
agg_causeSkin and other sense organ disorders	1.1159233	0.0008182
as.factor(year_id)2016:agg_causeCancers	0.1119215	0.8119150
as.factor(year_id)2016:agg_causeCardiovascular diseases	-0.2261237	0.6307000
as.factor(year_id)2016:agg_causeChronic respiratory diseases	0.0952092	0.8417381
as.factor(year_id)2016:agg_causeCommunicable and nutrition disorders	-0.0600710	0.8983721
as.factor(year_id)2016:agg_causeDiabetes and kidney diseases	0.0396297	0.9328516
as.factor(year_id)2016:agg_causeDigestive diseases	0.1050241	0.8233082
as.factor(year_id)2016:agg_causeEndocrine disorders	0.3811605	0.4177841
as.factor(year_id)2016:agg_causeInjuries	-0.1029154	0.8267989
as.factor(year_id)2016:agg_causeMaternal and neonatal conditions	-0.3575102	0.5279227
as.factor(year_id)2016:agg_causeMusculoskeletal conditions	0.0857897	0.8552701
as.factor(year_id)2016:agg_causeNeurological disorders	0.0523592	0.9113610
as.factor(year_id)2016:agg_causeOther non-communicable diseases	-0.0299608	0.9492087
as.factor(year_id)2016:agg_causePrevention and coordination	-0.1469024	0.7547926
as.factor(year_id)2016:agg_causeSkin and other sense organ disorders	0.0201944	0.9657527

```
glance(spendingovertime_fit)$adj.r.squared
```

```
## [1] 0.5959975
```

Age !! had to take out the observations with “All Ages” because I think it will just mess up the pairs but let me know what you think or whether you think there’s anything we can do with that group

We wonder whether there is a correlation between government healthcare expenditures in the emergency department and age. The age variable is categorical, split into 19 groups that generally include 5 years each, apart from the first (<1 year) and last (85 plus) groups.

To address this question, we began by using an overall test with ANOVA.

Below is an overall test of the null hypothesis that all of the means for age groups across the years are equal, as opposed to the alternative that at least one mean is different.

```
##              Df Sum Sq Mean Sq F value Pr(>F)
## age_group_name  18   4239   235.50   52.17 <2e-16 ***
## Residuals    6031   27226     4.51
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

In this F-test (ndf = 18, ddf = 6229), a significant difference among age groups was identified. Our p-value tells us that this data (or data more extreme) would be very unlikely if the null hypothesis were true because it shows statistical significance at an alpha well below 0.05. Therefore, we reject the null hypothesis that the mean expenditures for all age groups are equal.

To see which specific means may be different from one another, we used planned step-down tests with a Holm correction to minimize Type I errors.

```
## [1] 99
```


The pairwise t-tests used for our ANOVA step-down tests suggest that there are 99 different age pairs out of the 171 possible combinations. This tells us that more age pairs are different than are similar and that therefore the majority of age group pairs differ in terms of mean expenditures.

```
agefit <- linear_reg() %>%
  set_engine("lm") %>%
  fit(lmean_all ~ age_group_name, data = spending_malefemale)
tidy(agefit)
```


```
## # A tibble: 19 x 5
```

	term	estimate	std.error	statistic	p.value
	<chr>	<dbl>	<dbl>	<dbl>	<dbl>
## 1	(Intercept)	15.4	0.121	127.	0
## 2	age_group_name1 to 4	1.20	0.170	7.05	2.03e-12
## 3	age_group_name10 to 14	1.92	0.170	11.3	1.67e-29
## 4	age_group_name15 to 19	2.59	0.170	15.2	1.55e-51
## 5	age_group_name20 to 24	2.96	0.170	17.5	1.17e-66
## 6	age_group_name25 to 29	3.09	0.170	18.2	2.45e-72
## 7	age_group_name30 to 34	3.16	0.170	18.6	3.99e-75
## 8	age_group_name35 to 39	3.23	0.170	19.0	1.87e-78
## 9	age_group_name40 to 44	3.30	0.170	19.4	1.09e-81
## 10	age_group_name45 to 49	3.32	0.170	19.6	8.29e-83
## 11	age_group_name5 to 9	1.27	0.170	7.49	7.86e-14
## 12	age_group_name50 to 54	3.27	0.170	19.3	3.25e-80
## 13	age_group_name55 to 59	2.89	0.170	17.0	1.50e-63
## 14	age_group_name60 to 64	2.70	0.170	15.9	1.12e-55
## 15	age_group_name65 to 69	2.63	0.170	15.5	3.12e-53
## 16	age_group_name70 to 74	2.48	0.170	14.6	1.22e-47
## 17	age_group_name75 to 79	2.37	0.170	13.9	1.52e-43
## 18	age_group_name80 to 84	2.32	0.170	13.7	6.62e-42
## 19	age_group_name85 plus	2.39	0.170	14.1	3.11e-44


Emergency Department General Expenditures



Public Insurance Expenditures



Private Insurance Expenditures



Out-of-Pocket Expenditures



Gender and Age Interaction

```
## # A tibble: 3 x 5
##   term          estimate std.error statistic  p.value
##   <chr>          <dbl>    <dbl>    <dbl>    <dbl>
## 1 (Intercept)    17.1      0.0435     394.      0
## 2 sexMale       -0.0854    0.0565     -1.51    0.130
## 3 age_group_id  0.00328   0.000849     3.86  0.000113
## [1] 0.00250716

## # A tibble: 4 x 5
##   term          estimate std.error statistic p.value
##   <chr>          <dbl>    <dbl>    <dbl>    <dbl>
## 1 (Intercept)    17.1      0.0471     364.      0
## 2 sexMale       -0.0623    0.0677     -0.920  0.358
## 3 age_group_id  0.00379    0.00118     3.21  0.00134
## 4 sexMale:age_group_id -0.00105  0.00170     -0.619  0.536
## [1] 0.002405422

## # A tibble: 3 x 5
##   term          estimate std.error statistic  p.value
##   <chr>          <dbl>    <dbl>    <dbl>    <dbl>
```

```
## 1 (Intercept)    16.9      0.0500    339.      0
## 2 sexMale        -0.0310   0.0649     -0.477  6.33e- 1
## 3 age_group_id   -0.0132   0.000976   -13.5    5.69e-41
## [1] 0.02899412

## # A tibble: 4 x 5
##   term                estimate std.error statistic  p.value
##   <chr>              <dbl>    <dbl>    <dbl>    <dbl>
## 1 (Intercept)        16.9      0.0541    312.      0
## 2 sexMale             0.0178    0.0778     0.229  8.19e- 1
## 3 age_group_id       -0.0121    0.00136   -8.92   5.88e-19
## 4 sexMale:age_group_id -0.00222  0.00195   -1.14   2.56e- 1
## [1] 0.02904063

## # A tibble: 3 x 5
##   term                estimate std.error statistic  p.value
##   <chr>              <dbl>    <dbl>    <dbl>    <dbl>
## 1 (Intercept)        14.7      0.0476    309.      0
## 2 sexMale            -0.0456    0.0618    -0.739  4.60e- 1
## 3 age_group_id       -0.0113    0.000930   -12.2    1.35e-33
## [1] 0.02360949

## # A tibble: 4 x 5
##   term                estimate std.error statistic  p.value
##   <chr>              <dbl>    <dbl>    <dbl>    <dbl>
## 1 (Intercept)        14.7      0.0515    285.      0
## 2 sexMale            -0.00247    0.0741    -0.0333  9.73e- 1
## 3 age_group_id       -0.0104    0.00129   -8.01    1.39e-15
## 4 sexMale:age_group_id -0.00196    0.00186   -1.06    2.91e- 1
## [1] 0.02362785
```

In order to test the possibility that there is a joint interaction of gender and age, a main effects and interaction effects linear regression model has been fit to the data. As a whole, it shows that the interaction of gender and age slightly increases the accuracy of the regression for public and private spending as seen by the increased adjusted R^2 value. However, for out-of-pocket spending, it decreases the adjusted R^2 value. Nevertheless, overall, the adjusted R^2 values for all three types of spending are incredibly low, which further point to our conclusion that age may not affect the level of spending from different sources.

Age and Disease Interaction

Table 3: R^2 Values for the Main Effects and Interaction Models Analyzing Disease Type and Age

Payer	Main_Effects	Interaction
Public Spending	0.5068870	0.5293510
Private Spending	0.5054947	0.5149051
Out-of-Pocket Spending	0.5161683	0.5261024

```
## # A tibble: 16 x 5
##   term                estimate std.error statistic  p.value
##   <chr>              <dbl>    <dbl>    <dbl>    <dbl>
## 1 (Intercept)        16.9      0.0713    237.      0
## 2 agg_causeCancers   -2.69     0.0994   -27.1    1.45e-152
```

```
## 3 agg_causeCardiovascular diseases      1.14      0.0994      11.5 3.28e- 30
## 4 agg_causeChronic respiratory diseases  1.29      0.101      12.8 7.42e- 37
## 5 agg_causeCommunicable and nutrition d~ 1.61      0.0994      16.2 7.68e- 58
## 6 agg_causeDiabetes and kidney diseases  0.0569    0.0994      0.573 5.67e- 1
## 7 agg_causeDigestive diseases            1.66      0.0994      16.7 5.41e- 61
## 8 agg_causeEndocrine disorders           -1.25     0.0994     -12.6 7.61e- 36
## 9 agg_causeInjuries                      1.99      0.0994      20.0 2.81e- 86
## 10 agg_causeMaternal and neonatal condit~-3.50     0.120     -29.2 9.83e-176
## 11 agg_causeMusculoskeletal conditions   0.816     0.0994      8.21 2.65e- 16
## 12 agg_causeNeurological disorders       0.780     0.0994      7.85 4.89e- 15
## 13 agg_causeOther non-communicable disea~ 1.25      0.0994      12.6 6.29e- 36
## 14 agg_causePrevention and coordination   -3.06     0.0994     -30.8 4.22e-193
## 15 agg_causeSkin and other sense organ d~ 0.892     0.0994      8.97 3.78e- 19
## 16 age_group_id                          0.00334   0.000556      6.01 2.02e- 9
```

```
## [1] 0.5726639
```

```
## # A tibble: 30 x 5
```

##	term	estimate	std.error	statistic	p.value
##	<chr>	<dbl>	<dbl>	<dbl>	<dbl>
## 1	(Intercept)	17.1	0.0817	210.	0
## 2	agg_causeCancers	-3.16	0.116	-27.3	4.41e-155
## 3	agg_causeCardiovascular diseases	0.610	0.116	5.28	1.35e- 7
## 4	agg_causeChronic respiratory diseases	1.08	0.116	9.31	1.70e- 20
## 5	agg_causeCommunicable and nutrition d~	1.44	0.116	12.5	2.49e- 35
## 6	agg_causeDiabetes and kidney diseases	-0.237	0.116	-2.06	3.99e- 2
## 7	agg_causeDigestive diseases	1.50	0.116	13.0	4.42e- 38
## 8	agg_causeEndocrine disorders	-1.42	0.116	-12.3	2.98e- 34
## 9	agg_causeInjuries	1.74	0.116	15.1	2.31e- 50
## 10	agg_causeMaternal and neonatal condit~	-2.69	0.140	-19.2	6.58e- 80
##	... with 20 more rows				

```
## # A tibble: 0 x 0
```

```
## # A tibble: 16 x 5
```

##	term	estimate	std.error	statistic	p.value
##	<chr>	<dbl>	<dbl>	<dbl>	<dbl>
## 1	(Intercept)	16.2	0.0841	193.	0
## 2	agg_causeCancers	-2.23	0.117	-19.0	2.32e- 78
## 3	agg_causeCardiovascular diseases	1.63	0.117	13.9	1.68e- 43
## 4	agg_causeChronic respiratory diseases	1.75	0.119	14.8	1.74e- 48
## 5	agg_causeCommunicable and nutrition d~	2.09	0.117	17.9	1.39e- 69
## 6	agg_causeDiabetes and kidney diseases	-0.208	0.117	-1.78	7.54e- 2
## 7	agg_causeDigestive diseases	2.46	0.117	21.0	2.66e- 94
## 8	agg_causeEndocrine disorders	-0.749	0.117	-6.39	1.74e- 10
## 9	agg_causeInjuries	3.48	0.117	29.7	4.60e-181
## 10	agg_causeMaternal and neonatal condit~	-3.54	0.141	-25.1	2.40e-132
## 11	agg_causeMusculoskeletal conditions	1.52	0.117	13.0	3.16e- 38
## 12	agg_causeNeurological disorders	1.35	0.117	11.6	1.32e- 30
## 13	agg_causeOther non-communicable disea~	2.06	0.117	17.6	1.60e- 67
## 14	agg_causePrevention and coordination	-2.63	0.117	-22.4	3.83e-107
## 15	agg_causeSkin and other sense organ d~	1.44	0.117	12.3	1.90e- 34
## 16	age_group_id	-0.0131	0.000655	-20.0	1.98e- 86

```
## [1] 0.5626007
```

```
## # A tibble: 30 x 5
```

```
##      term                estimate std.error statistic    p.value
##      <chr>                <dbl>      <dbl>      <dbl>      <dbl>
## 1 (Intercept)             16.5        0.0981     168.      0
## 2 agg_causeCancers        -2.75        0.139     -19.8 6.48e- 85
## 3 agg_causeCardiovascular diseases    1.06        0.139      7.65 2.25e- 14
## 4 agg_causeChronic respiratory diseases 1.54        0.140     11.0 5.06e- 28
## 5 agg_causeCommunicable and nutrition d~ 1.87        0.139     13.5 1.12e- 40
## 6 agg_causeDiabetes and kidney diseases -0.611       0.139     -4.41 1.08e- 5
## 7 agg_causeDigestive diseases    2.33        0.139     16.8 5.58e- 62
## 8 agg_causeEndocrine disorders    -1.00        0.139     -7.21 6.47e- 13
## 9 agg_causeInjuries         3.27        0.139     23.6 2.17e-117
## 10 agg_causeMaternal and neonatal condit~ -3.16       0.168     -18.8 1.61e- 76
## # ... with 20 more rows

## [1] 0.5734167

## # A tibble: 16 x 5
##      term                estimate std.error statistic    p.value
##      <chr>                <dbl>      <dbl>      <dbl>      <dbl>
## 1 (Intercept)             14.5        0.0786     184.      0
## 2 agg_causeCancers        -2.72        0.110     -24.8 1.48e-129
## 3 agg_causeCardiovascular diseases    0.994       0.110      9.08 1.48e- 19
## 4 agg_causeChronic respiratory diseases 1.15        0.111     10.4 6.12e- 25
## 5 agg_causeCommunicable and nutrition d~ 1.72        0.110     15.7 3.22e- 54
## 6 agg_causeDiabetes and kidney diseases -0.395       0.110     -3.61 3.09e- 4
## 7 agg_causeDigestive diseases    1.86        0.110     17.0 3.52e- 63
## 8 agg_causeEndocrine disorders    -0.875       0.110     -7.99 1.55e- 15
## 9 agg_causeInjuries         2.69        0.110     24.6 5.35e-127
## 10 agg_causeMaternal and neonatal condit~ -3.89       0.132     -29.5 1.97e-178
## 11 agg_causeMusculoskeletal conditions  0.916       0.110      8.36 7.52e- 17
## 12 agg_causeNeurological disorders    0.771       0.110      7.04 2.08e- 12
## 13 agg_causeOther non-communicable disea~ 1.60        0.110     14.6 2.16e- 47
## 14 agg_causePrevention and coordination -3.33        0.110     -30.4 2.09e-189
## 15 agg_causeSkin and other sense organ d~ 0.874       0.110      7.98 1.75e- 15
## 16 age_group_id          -0.0112     0.000613    -18.3 3.42e- 73

## [1] 0.5759006

## [1] 0.5873628
```

Table 4: Temp

term	estimate	p.value
(Intercept)	14.7092021	0.0000000
agg_causeCancers	-3.1442198	0.0000000
agg_causeCardiovascular diseases	0.4235641	0.0010880
agg_causeChronic respiratory diseases	0.9331922	0.0000000
agg_causeCommunicable and nutrition disorders	1.5405779	0.0000000
agg_causeDiabetes and kidney diseases	-0.7646196	0.0000000
agg_causeDigestive diseases	1.7105672	0.0000000
agg_causeEndocrine disorders	-1.1024696	0.0000000
agg_causeInjuries	2.4308631	0.0000000
agg_causeMaternal and neonatal conditions	-3.4564444	0.0000000
agg_causeMusculoskeletal conditions	0.6503271	0.0000005
agg_causeNeurological disorders	0.4510409	0.0005043
agg_causeOther non-communicable diseases	1.3397594	0.0000000

term	estimate	p.value
agg_causePrevention and coordination	-3.5027704	0.0000000
agg_causeSkin and other sense organ disorders	0.6936139	0.0000001
age_group_id	-0.0218172	0.0000000
agg_causeCancers:age_group_id	0.0193805	0.0000000
agg_causeCardiovascular diseases:age_group_id	0.0259313	0.0000000
agg_causeChronic respiratory diseases:age_group_id	0.0098494	0.0024985
agg_causeCommunicable and nutrition disorders:age_group_id	0.0079308	0.0148507
agg_causeDiabetes and kidney diseases:age_group_id	0.0167852	0.0000003
agg_causeDigestive diseases:age_group_id	0.0067519	0.0380775
agg_causeEndocrine disorders:age_group_id	0.0103193	0.0015291
agg_causeInjuries:age_group_id	0.0117393	0.0003125
agg_causeMaternal and neonatal conditions:age_group_id	-0.0191120	0.0000016
agg_causeMusculoskeletal conditions:age_group_id	0.0120678	0.0002110
agg_causeNeurological disorders:age_group_id	0.0145592	0.0000078
agg_causeOther non-communicable diseases:age_group_id	0.0117118	0.0003228
agg_causePrevention and coordination:age_group_id	0.0077166	0.0177767
agg_causeSkin and other sense organ disorders:age_group_id	0.0081909	0.0118748

References

Institute for Health Metrics and Evaluation (IHME). United States Healthcare Spending in Emergency Departments by Health Condition 2006-2016. Seattle, United States of America: Institute for Health Metrics and Evaluation (IHME), 2021.

Woody Scott K, Liu A, Chen C, Kaldjian AS, Sabbatini AK, Duber, HC, Dieleman JL. Healthcare Spending in U.S. Emergency Departments by Health Condition, 2006-2016. PLOS One. 27 October 2021.