

# Lecture 5: Basics of Bayesian Hypothesis Testing

Merlise Clyde

September 9, 2021



# Hypothesis Testing

Suppose we have univariate data  $y_i \stackrel{iid}{\sim} \mathcal{N}(\theta, 1)$



# Hypothesis Testing

Suppose we have univariate data  $y_i \stackrel{iid}{\sim} \mathcal{N}(\theta, 1)$

goal is to test  $\mathcal{H}_0 : \theta = 0$ ; vs  $\mathcal{H}_1 : \theta \neq 0$



# Hypothesis Testing

Suppose we have univariate data  $y_i \stackrel{iid}{\sim} \mathcal{N}(\theta, 1)$

goal is to test  $\mathcal{H}_0 : \theta = 0$ ; vs  $\mathcal{H}_1 : \theta \neq 0$

Frequentist testing - likelihood ratio, Wald, score, UMP, confidence regions, etc




# Hypothesis Testing

Suppose we have univariate data  $y_i \stackrel{iid}{\sim} \mathcal{N}(\theta, 1)$

goal is to test  $\mathcal{H}_0 : \theta = 0$ ; vs  $\mathcal{H}_1 : \theta \neq 0$

Frequentist testing - likelihood ratio, Wald, score, UMP, confidence regions, etc

- Need a **test statistic**  $T(y^{(n)})$  (and its sampling distribution)




# Hypothesis Testing

Suppose we have univariate data  $y_i \stackrel{iid}{\sim} \mathcal{N}(\theta, 1)$

goal is to test  $\mathcal{H}_0 : \theta = 0$ ; vs  $\mathcal{H}_1 : \theta \neq 0$

Frequentist testing - likelihood ratio, Wald, score, UMP, confidence regions, etc

- Need a **test statistic**  $T(y^{(n)})$  (and its sampling distribution)



- **p-value:** Calculate the probability of seeing a dataset/test statistics as extreme or more extreme than the observed data with repeated sampling under the null hypothesis

# Errors

if p-value is less than a pre-specified  $\alpha$  then reject  $\mathcal{H}_0$  in favor of  $\mathcal{H}_1$



# Errors

if p-value is less than a pre-specified  $\alpha$  then reject  $\mathcal{H}_0$  in favor of  $\mathcal{H}_1$

- Type I error: falsely concluding in favor of  $\mathcal{H}_1$  when  $\mathcal{H}_0$  is true



# Errors

if p-value is less than a pre-specified  $\alpha$  then reject  $\mathcal{H}_0$  in favor of  $\mathcal{H}_1$

- Type I error: falsely concluding in favor of  $\mathcal{H}_1$  when  $\mathcal{H}_0$  is true
- To maintain a Type I error rate of  $\alpha$ , then we reject  $\mathcal{H}_0$  in favor of  $\mathcal{H}_1$  when  $p < \alpha$



# Errors

if p-value is less than a pre-specified  $\alpha$  then reject  $\mathcal{H}_0$  in favor of  $\mathcal{H}_1$

- Type I error: falsely concluding in favor of  $\mathcal{H}_1$  when  $\mathcal{H}_0$  is true
- To maintain a Type I error rate of  $\alpha$ , then we reject  $\mathcal{H}_0$  in favor of  $\mathcal{H}_1$  when  $p < \alpha$

For this to be a valid frequentist test the p-value must have a uniform distribution under  $\mathcal{H}_0$



# Errors

if p-value is less than a pre-specified  $\alpha$  then reject  $\mathcal{H}_0$  in favor of  $\mathcal{H}_1$

- Type I error: falsely concluding in favor of  $\mathcal{H}_1$  when  $\mathcal{H}_0$  is true
- To maintain a Type I error rate of  $\alpha$ , then we reject  $\mathcal{H}_0$  in favor of  $\mathcal{H}_1$  when  $p < \alpha$

For this to be a valid frequentist test the p-value must have a uniform distribution under  $\mathcal{H}_0$

- Type II error: failing to conclude in favor of  $\mathcal{H}_1$  when  $\mathcal{H}_1$  is true



# Errors

if p-value is less than a pre-specified  $\alpha$  then reject  $H_0$  in favor of  $H_1$

- Type I error: falsely concluding in favor of  $\mathcal{H}_1$  when  $\mathcal{H}_0$  is true
  - To maintain a Type I error rate of  $\alpha$ , then we reject  $\mathcal{H}_0$  in favor of  $\mathcal{H}_1$  when  $p < \alpha$

For this to be a valid frequentist test the p-value must have a uniform distribution under  $H_0$ .

- Type II error: failing to conclude in favor of  $\mathcal{H}_1$  when  $\mathcal{H}_1$  is true
  - $1 - P(\text{Type II error})$  is the **power** of the test



# Errors

if p-value is less than a pre-specified  $\alpha$  then reject  $\mathcal{H}_0$  in favor of  $\mathcal{H}_1$

- Type I error: falsely concluding in favor of  $\mathcal{H}_1$  when  $\mathcal{H}_0$  is true
  - To maintain a Type I error rate of  $\alpha$ , then we reject  $\mathcal{H}_0$  in favor of  $\mathcal{H}_1$  when  $p < \alpha$

For this to be a valid frequentist test the p-value must have a uniform distribution under  $\mathcal{H}_0$ .

- Type II error: failing to conclude in favor of  $\mathcal{H}_1$  when  $\mathcal{H}_1$  is true
  - $1 - P(\text{Type II error})$  is the **power** of the test

**Note:** we *never* conclude in favor of  $\mathcal{H}_0$ . We are looking for enough evidence to reject  $\mathcal{H}_0$ . But if we fail to reject we do not conclude that it is true!



# Bayesian Approach

1. Put a prior on  $\theta$ ,  $\pi(\theta) = \mathcal{N}(\theta_0, 1/\tau_0^2)$ .




# Bayesian Approach

1. Put a prior on  $\theta$ ,  $\pi(\theta) = \mathcal{N}(\theta_0, 1/\tau_0^2)$ .
2. Compute posterior  $\theta | y^{(n)} \sim \mathcal{N}(\theta_n, 1/\tau_n^2)$  for updated parameters  $\theta_n$  and  $\tau_n^2$ .



# Bayesian Approach

1. Put a prior on  $\theta$ ,  $\pi(\theta) = \mathcal{N}(\theta_0, 1/\tau_0^2)$ .
2. Compute posterior  $\theta | y^{(n)} \sim \mathcal{N}(\theta_n, 1/\tau_n^2)$  for updated parameters  $\theta_n$  and  $\tau_n^2$ .



# Informal

## Credible Intervals

1. Compute a 95% CI based on the posterior.



# Informal

## Credible Intervals

1. Compute a 95% CI based on the posterior.
2. Reject  $H_0$  if interval does not contain zero.



# Informal

## Credible Intervals

1. Compute a 95% CI based on the posterior.
2. Reject  $\mathcal{H}_0$  if interval does not contain zero.

## Tail Areas:

1. Compute  $\Pr(\theta > 0 | y^{(n)})$  and  $\Pr(\theta < 0 | y^{(n)})$



# Informal

# Credible Intervals

1. Compute a 95% CI based on the posterior.
  2. Reject  $\mathcal{H}_0$  if interval does not contain zero.

## Tail Areas:

1. Compute  $\Pr(\theta > 0 | y^{(n)})$  and  $\Pr(\theta < 0 | y^{(n)})$
  2. Report minimum of these probabilities as a "Bayesian p-value"

Note: Tail probability is not the same as  $\Pr(\mathcal{H}_0 \mid y^{(n)})$



# Formal Bayesian Hypothesis Testing

Unknowns are  $\mathcal{H}_0$  and  $\mathcal{H}_1$

Put a prior on the actual hypotheses/models, that is, on  
 $\pi(\mathcal{H}_0) = \Pr(\mathcal{H}_0 = \text{True})$  and  $\pi(\mathcal{H}_1) = \Pr(\mathcal{H}_1 = \text{True})$ .



# Formal Bayesian Hypothesis Testing

Unknowns are  $\mathcal{H}_0$  and  $\mathcal{H}_1$

Put a prior on the actual hypotheses/models, that is, on

$\pi(\mathcal{H}_0) = \Pr(\mathcal{H}_0 = \text{True})$  and  $\pi(\mathcal{H}_1) = \Pr(\mathcal{H}_1 = \text{True})$ .

- For example, set  $\pi(\mathcal{H}_0) = 0.5$  and  $\pi(\mathcal{H}_1) = 0.5$ , if *a priori*, we believe the two hypotheses are equally likely.



# Formal Bayesian Hypothesis Testing

Unknowns are  $\mathcal{H}_0$  and  $\mathcal{H}_1$

Put a prior on the actual hypotheses/models, that is, on  
 $\pi(\mathcal{H}_0) = \Pr(\mathcal{H}_0 = \text{True})$  and  $\pi(\mathcal{H}_1) = \Pr(\mathcal{H}_1 = \text{True})$ .

- For example, set  $\pi(\mathcal{H}_0) = 0.5$  and  $\pi(\mathcal{H}_1) = 0.5$ , if *a priori*, we believe the two hypotheses are equally likely. Likelihood of the hypotheses

$$\mathcal{L}(\mathcal{H}_i) \propto p(y^{(n)} | \mathcal{H}_i)$$



# Formal Bayesian Hypothesis Testing

Unknowns are  $\mathcal{H}_0$  and  $\mathcal{H}_1$

Put a prior on the actual hypotheses/models, that is, on  
 $\pi(\mathcal{H}_0) = \Pr(\mathcal{H}_0 = \text{True})$  and  $\pi(\mathcal{H}_1) = \Pr(\mathcal{H}_1 = \text{True})$ .

- For example, set  $\pi(\mathcal{H}_0) = 0.5$  and  $\pi(\mathcal{H}_1) = 0.5$ , if *a priori*, we believe the two hypotheses are equally likely. Likelihood of the hypotheses

$$\mathcal{L}(\mathcal{H}_i) \propto p(y^{(n)} | \mathcal{H}_i)$$

$$p(y^{(n)} | \mathcal{H}_0) = \prod_{i=1}^n (2\pi)^{-1/2} \exp -\frac{1}{2}(y_i - 0)^2$$



# Formal Bayesian Hypothesis Testing

Unknowns are  $\mathcal{H}_0$  and  $\mathcal{H}_1$

Put a prior on the actual hypotheses/models, that is, on  
 $\pi(\mathcal{H}_0) = \Pr(\mathcal{H}_0 = \text{True})$  and  $\pi(\mathcal{H}_1) = \Pr(\mathcal{H}_1 = \text{True})$ .

- For example, set  $\pi(\mathcal{H}_0) = 0.5$  and  $\pi(\mathcal{H}_1) = 0.5$ , if *a priori*, we believe the two hypotheses are equally likely. Likelihood of the hypotheses

$$\mathcal{L}(\mathcal{H}_i) \propto p(y^{(n)} | \mathcal{H}_i)$$

$$p(y^{(n)} | \mathcal{H}_0) = \prod_{i=1}^n (2\pi)^{-1/2} \exp -\frac{1}{2}(y_i - 0)^2$$

$$p(y^{(n)} | \mathcal{H}_1) = \int_{\Theta} p(y^{(n)} | \mathcal{H}_1, \theta) p(\theta | \mathcal{H}_1) d\theta$$



# Bayesian Approach

Priors on parameters under each hypothesis



# Bayesian Approach

## Priors on parameters under each hypothesis

In our simple normal model, the only unknown parameter is  $\theta$



# Bayesian Approach

## Priors on parameters under each hypothesis

In our simple normal model, the only unknown parameter is  $\theta$

- under  $\mathcal{H}_0$ ,  $\theta = 0$  with probability 1



# Bayesian Approach

## Priors on parameters under each hypothesis

In our simple normal model, the only unknown parameter is  $\theta$

- under  $\mathcal{H}_0$ ,  $\theta = 0$  with probability 1
  - under  $\mathcal{H}_0$ ,  $\theta \in \mathbb{R}$  Could take  $\pi(\theta) = \mathcal{N}(\theta_0, 1/\tau_0^2)$ .



# Bayesian Approach

## Priors on parameters under each hypothesis

In our simple normal model, the only unknown parameter is  $\theta$

- under  $\mathcal{H}_0$ ,  $\theta = 0$  with probability 1
  - under  $\mathcal{H}_0$ ,  $\theta \in \mathbb{R}$  Could take  $\pi(\theta) = \mathcal{N}(\theta_0, 1/\tau_0^2)$ .
  - Compute marginal likelihoods for each hypothesis, that is,  $\mathcal{L}(\mathcal{H}_0)$  and  $\mathcal{L}(\mathcal{H}_1)$ .



# Bayesian Approach

Priors on parameters under each hypothesis

In our simple normal model, the only unknown parameter is  $\theta$

- under  $\mathcal{H}_0$ ,  $\theta = 0$  with probability 1
- under  $\mathcal{H}_0$ ,  $\theta \in \mathbb{R}$  Could take  $\pi(\theta) = \mathcal{N}(\theta_0, 1/\tau_0^2)$ .
- Compute marginal likelihoods for each hypothesis, that is,  $\mathcal{L}(\mathcal{H}_0)$  and  $\mathcal{L}(\mathcal{H}_1)$ .
- Obtain posterior probabilities of  $\mathcal{H}_0$  and  $\mathcal{H}_1$  via Bayes Theorem.



# Bayesian Approach - Decisions

Loss function for hypothesis testing

- $\hat{\mathcal{H}}$  is the chosen hypothesis
- $\mathcal{H}_{true}$  is the true hypothesis,  $\mathcal{H}$  for short



# Bayesian Approach - Decisions

Loss function for hypothesis testing

- $\hat{\mathcal{H}}$  is the chosen hypothesis
- $\mathcal{H}_{true}$  is the true hypothesis,  $\mathcal{H}$  for short

Two types of errors:

- Type I error:  $\hat{\mathcal{H}} = 1$  and  $\mathcal{H} = 0$



# Bayesian Approach - Decisions

Loss function for hypothesis testing

- $\hat{\mathcal{H}}$  is the chosen hypothesis
- $\mathcal{H}_{true}$  is the true hypothesis,  $\mathcal{H}$  for short

Two types of errors:

- Type I error:  $\hat{\mathcal{H}} = 1$  and  $\mathcal{H} = 0$
- Type II error:  $\hat{\mathcal{H}} = 0$  and  $\mathcal{H} = 1$



# Bayesian Approach - Decisions

Loss function for hypothesis testing

- $\hat{\mathcal{H}}$  is the chosen hypothesis
- $\mathcal{H}_{true}$  is the true hypothesis,  $\mathcal{H}$  for short

Two types of errors:

- Type I error:  $\hat{\mathcal{H}} = 1$  and  $\mathcal{H} = 0$
- Type II error:  $\hat{\mathcal{H}} = 0$  and  $\mathcal{H} = 1$

Loss function:

$$L(\hat{\mathcal{H}}, \mathcal{H}) = w_1 \mathbf{1}(\hat{\mathcal{H}} = 1, \mathcal{H} = 0) + w_2 \mathbf{1}(\hat{\mathcal{H}} = 0, \mathcal{H} = 1)$$

- $w_1$  weights how bad making a Type I error
- $w_2$  weights how bad making a Type II error



# Loss Function Functions and Decisions

- Relative weights

$$L(\hat{\mathcal{H}}, \mathcal{H}) = \mathbf{1}(\hat{\mathcal{H}} = 1, \mathcal{H} = 0) + w \mathbf{1}(\hat{\mathcal{H}} = 0, \mathcal{H} = 1)$$



# Loss Function Functions and Decisions

- Relative weights

$$L(\hat{\mathcal{H}}, \mathcal{H}) = \mathbb{1}(\hat{\mathcal{H}} = 1, \mathcal{H} = 0) + w \mathbb{1}(\hat{\mathcal{H}} = 0, \mathcal{H} = 1)$$

- Special case  $w = 1$

$$L(\hat{\mathcal{H}}, \mathcal{H}) = \mathbb{1}(\hat{\mathcal{H}} \neq \mathcal{H})$$

- known as 0-1 loss (most common)



# Loss Function Functions and Decisions

- Relative weights

$$L(\hat{\mathcal{H}}, \mathcal{H}) = \mathbb{1}(\hat{\mathcal{H}} = 1, \mathcal{H} = 0) + w \mathbb{1}(\hat{\mathcal{H}} = 0, \mathcal{H} = 1)$$

- Special case  $w = 1$

$$L(\hat{\mathcal{H}}, \mathcal{H}) = \mathbb{1}(\hat{\mathcal{H}} \neq \mathcal{H})$$

- known as 0-1 loss (most common)
- Bayes Risk (Posterior Expected Loss)

$$\mathbb{E}_{\mathcal{H}|y^{(n)}}[L(\hat{\mathcal{H}}, \mathcal{H})] = \mathbb{1}(\hat{\mathcal{H}} = 1)\pi(\mathcal{H}_0 | y^{(n)}) + \mathbb{1}(\hat{\mathcal{H}} = 0)\pi(\mathcal{H}_1 | y^{(n)})$$



# Loss Function Functions and Decisions

- Relative weights

$$L(\hat{\mathcal{H}}, \mathcal{H}) = \mathbb{1}(\hat{\mathcal{H}} = 1, \mathcal{H} = 0) + w \mathbb{1}(\hat{\mathcal{H}} = 0, \mathcal{H} = 1)$$

- Special case  $w = 1$

$$L(\hat{\mathcal{H}}, \mathcal{H}) = \mathbb{1}(\hat{\mathcal{H}} \neq \mathcal{H})$$

- known as 0-1 loss (most common)
- Bayes Risk (Posterior Expected Loss)

$$\mathbb{E}_{\mathcal{H}|y^{(n)}}[L(\hat{\mathcal{H}}, \mathcal{H})] = \mathbb{1}(\hat{\mathcal{H}} = 1)\pi(\mathcal{H}_0 | y^{(n)}) + \mathbb{1}(\hat{\mathcal{H}} = 0)\pi(\mathcal{H}_1 | y^{(n)})$$

- Minimize loss by picking hypothesis with the highest posterior probability



# Bayesian hypothesis testing

- Using Bayes theorem,

$$\pi(\mathcal{H}_1 \mid Y) = \frac{p(y^{(n)} \mid \mathcal{H}_1)\pi(\mathcal{H}_1)}{p(y^{(n)} \mid \mathcal{H}_0)\pi(\mathcal{H}_0) + p(y^{(n)} \mid \mathcal{H}_1)\pi(\mathcal{H}_1)},$$

where  $p(y^{(n)} | \mathcal{H}_0)$  and  $p(y^{(n)} | \mathcal{H}_1)$  are the marginal likelihoods of the hypotheses.



# Bayesian hypothesis testing

- Using Bayes theorem,

$$\pi(\mathcal{H}_1 \mid Y) = \frac{p(y^{(n)} \mid \mathcal{H}_1)\pi(\mathcal{H}_1)}{p(y^{(n)} \mid \mathcal{H}_0)\pi(\mathcal{H}_0) + p(y^{(n)} \mid \mathcal{H}_1)\pi(\mathcal{H}_1)},$$

where  $p(y^{(n)} \mid \mathcal{H}_0)$  and  $p(y^{(n)} \mid \mathcal{H}_1)$  are the marginal likelihoods hypotheses.

- If for example we set  $\pi(\mathcal{H}_0) = 0.5$  and  $\pi(\mathcal{H}_1) = 0.5$  *a priori*, then

$$\begin{aligned}\pi(\mathcal{H}_1 \mid Y) &= \frac{0.5p(y^{(n)} \mid \mathcal{H}_1)}{0.5p(y^{(n)} \mid \mathcal{H}_0) + 0.5p(y^{(n)} \mid \mathcal{H}_1)} \\ &= \frac{p(y^{(n)} \mid \mathcal{H}_1)}{p(y^{(n)} \mid \mathcal{H}_0) + p(y^{(n)} \mid \mathcal{H}_1)} = \frac{1}{\frac{p(y^{(n)} \mid \mathcal{H}_0)}{p(y^{(n)} \mid \mathcal{H}_1)} + 1}.\end{aligned}$$



# Bayesian hypothesis testing

- Using Bayes theorem,

$$\pi(\mathcal{H}_1 | Y) = \frac{p(y^{(n)} | \mathcal{H}_1)\pi(\mathcal{H}_1)}{p(y^{(n)} | \mathcal{H}_0)\pi(\mathcal{H}_0) + p(y^{(n)} | \mathcal{H}_1)\pi(\mathcal{H}_1)},$$

where  $p(y^{(n)} | \mathcal{H}_0)$  and  $p(y^{(n)} | \mathcal{H}_1)$  are the marginal likelihoods hypotheses.

- If for example we set  $\pi(\mathcal{H}_0) = 0.5$  and  $\pi(\mathcal{H}_1) = 0.5$  *a priori*, then

$$\begin{aligned}\pi(\mathcal{H}_1 | Y) &= \frac{0.5p(y^{(n)} | \mathcal{H}_1)}{0.5p(y^{(n)} | \mathcal{H}_0) + 0.5p(y^{(n)} | \mathcal{H}_1)} \\ &= \frac{p(y^{(n)} | \mathcal{H}_1)}{p(y^{(n)} | \mathcal{H}_0) + p(y^{(n)} | \mathcal{H}_1)} = \frac{1}{\frac{p(y^{(n)} | \mathcal{H}_0)}{p(y^{(n)} | \mathcal{H}_1)} + 1}.\end{aligned}$$

- The ratio  $\frac{p(y^{(n)} | \mathcal{H}_0)}{p(y^{(n)} | \mathcal{H}_1)}$  is known as the **Bayes factor** in favor of  $\mathcal{H}_0$ , and often written as  $\mathcal{BF}_{01}$ . Similarly, we can compute  $\mathcal{BF}_{10}$ .



# Bayes factors

- **Bayes factor:** is a ratio of marginal likelihoods and it provides a weight of evidence in the data in favor of one model over another.



# Bayes factors

- **Bayes factor:** is a ratio of marginal likelihoods and it provides a weight of evidence in the data in favor of one model over another.
- It is often used as an alternative to the frequentist p-value.



# Bayes factors

- **Bayes factor:** is a ratio of marginal likelihoods and it provides a weight of evidence in the data in favor of one model over another.
- It is often used as an alternative to the frequentist p-value.
- **Rule of thumb:**  $\mathcal{BF}_{01} > 10$  is strong evidence for  $\mathcal{H}_0$ ;  $\mathcal{BF}_{01} > 100$  is decisive evidence for  $\mathcal{H}_0$ .



# Bayes factors

- **Bayes factor:** is a ratio of marginal likelihoods and it provides a weight of evidence in the data in favor of one model over another.
- It is often used as an alternative to the frequentist p-value.
- **Rule of thumb:**  $\mathcal{BF}_{01} > 10$  is strong evidence for  $\mathcal{H}_0$ ;  $\mathcal{BF}_{01} > 100$  is decisive evidence for  $\mathcal{H}_0$ .
- Notice that for our example,

$$\pi(\mathcal{H}_1 \mid Y) = \frac{1}{\frac{p(y^{(n)} \mid \mathcal{H}_0)}{p(y^{(n)} \mid \mathcal{H}_1)} + 1} = \frac{1}{\mathcal{BF}_{01} + 1}$$

the higher the value of  $\mathcal{BF}_{01}$ , that is, the weight of evidence in the data in favor of  $\mathcal{H}_0$ , the lower the marginal posterior probability that  $\mathcal{H}_1$  is true.



# Bayes factors

- **Bayes factor:** is a ratio of marginal likelihoods and it provides a weight of evidence in the data in favor of one model over another.
- It is often used as an alternative to the frequentist p-value.
- **Rule of thumb:**  $\mathcal{BF}_{01} > 10$  is strong evidence for  $\mathcal{H}_0$ ;  $\mathcal{BF}_{01} > 100$  is decisive evidence for  $\mathcal{H}_0$ .
- Notice that for our example,

$$\pi(\mathcal{H}_1 \mid Y) = \frac{1}{\frac{p(y^{(n)} \mid \mathcal{H}_0)}{p(y^{(n)} \mid \mathcal{H}_1)} + 1} = \frac{1}{\mathcal{BF}_{01} + 1}$$

the higher the value of  $\mathcal{BF}_{01}$ , that is, the weight of evidence in the data in favor of  $\mathcal{H}_0$ , the lower the marginal posterior probability that  $\mathcal{H}_1$  is true.

- That is, here, as  $\mathcal{BF}_{01} \uparrow$ ,  $\pi(\mathcal{H}_1 \mid Y) \downarrow$ .



# Bayes factors

- Let's look at another way to think of Bayes factors. First, recall that

$$\pi(\mathcal{H}_1 \mid Y) = \frac{p(y^{(n)} \mid \mathcal{H}_1)\pi(\mathcal{H}_1)}{p(y^{(n)} \mid \mathcal{H}_0)\pi(\mathcal{H}_0) + p(y^{(n)} \mid \mathcal{H}_1)\pi(\mathcal{H}_1)},$$

so that

$$\begin{aligned} \frac{\pi(\mathcal{H}_0|Y)}{\pi(\mathcal{H}_1|Y)} &= \frac{p(y^{(n)}|\mathcal{H}_0)\pi(\mathcal{H}_0)}{p(y^{(n)}|\mathcal{H}_0)\pi(\mathcal{H}_0) + p(y^{(n)}|\mathcal{H}_1)\pi(\mathcal{H}_1)} \div \frac{p(y^{(n)}|\mathcal{H}_1)\pi(\mathcal{H}_1)}{p(y^{(n)}|\mathcal{H}_0)\pi(\mathcal{H}_0) + p(y^{(n)}|\mathcal{H}_1)\pi(\mathcal{H}_1)} \\ &= \frac{p(y^{(n)}|\mathcal{H}_0)\pi(\mathcal{H}_0)}{p(y^{(n)}|\mathcal{H}_0)\pi(\mathcal{H}_0) + p(y^{(n)}|\mathcal{H}_1)\pi(\mathcal{H}_1)} \times \frac{p(y^{(n)}|\mathcal{H}_0)\pi(\mathcal{H}_0) + p(y^{(n)}|\mathcal{H}_1)\pi(\mathcal{H}_1)}{p(y^{(n)}|\mathcal{H}_1)\pi(\mathcal{H}_1)} \\ \therefore \underbrace{\frac{\pi(\mathcal{H}_0|Y)}{\pi(\mathcal{H}_1|Y)}}_{\text{posterior odds}} &= \underbrace{\frac{\pi(\mathcal{H}_0)}{\pi(\mathcal{H}_1)}}_{\text{prior odds}} \times \underbrace{\frac{p(y^{(n)}|\mathcal{H}_0)}{p(y^{(n)}|\mathcal{H}_1)}}_{\text{Bayes factor } \mathcal{BF}_{01}} \end{aligned}$$



# Bayes factors

- Let's look at another way to think of Bayes factors. First, recall that

$$\pi(\mathcal{H}_1 | Y) = \frac{p(y^{(n)} | \mathcal{H}_1)\pi(\mathcal{H}_1)}{p(y^{(n)} | \mathcal{H}_0)\pi(\mathcal{H}_0) + p(y^{(n)} | \mathcal{H}_1)\pi(\mathcal{H}_1)},$$

so that


$$\begin{aligned}\frac{\pi(\mathcal{H}_0 | Y)}{\pi(\mathcal{H}_1 | Y)} &= \frac{p(y^{(n)} | \mathcal{H}_0)\pi(\mathcal{H}_0)}{p(y^{(n)} | \mathcal{H}_0)\pi(\mathcal{H}_0) + p(y^{(n)} | \mathcal{H}_1)\pi(\mathcal{H}_1)} \div \frac{p(y^{(n)} | \mathcal{H}_1)\pi(\mathcal{H}_1)}{p(y^{(n)} | \mathcal{H}_0)\pi(\mathcal{H}_0) + p(y^{(n)} | \mathcal{H}_1)\pi(\mathcal{H}_1)} \\ &= \underbrace{\frac{p(y^{(n)} | \mathcal{H}_0)\pi(\mathcal{H}_0)}{p(y^{(n)} | \mathcal{H}_0)\pi(\mathcal{H}_0) + p(y^{(n)} | \mathcal{H}_1)\pi(\mathcal{H}_1)}}_{\text{posterior odds}} \times \underbrace{\frac{p(y^{(n)} | \mathcal{H}_0)\pi(\mathcal{H}_0) + p(y^{(n)} | \mathcal{H}_1)\pi(\mathcal{H}_1)}{p(y^{(n)} | \mathcal{H}_1)\pi(\mathcal{H}_1)}}_{\text{prior odds}} \\ \therefore \underbrace{\frac{\pi(\mathcal{H}_0 | Y)}{\pi(\mathcal{H}_1 | Y)}}_{\text{posterior odds}} &= \underbrace{\frac{\pi(\mathcal{H}_0)}{\pi(\mathcal{H}_1)}}_{\text{prior odds}} \times \underbrace{\frac{p(y^{(n)} | \mathcal{H}_0)}{p(y^{(n)} | \mathcal{H}_1)}}_{\text{Bayes factor } \mathcal{BF}_{01}}\end{aligned}$$

- Therefore, the Bayes factor can be thought of as the factor by which our prior odds change (towards the posterior odds) in the light of the data.



# Likelihoods & Evidence

Maximized Likelihood




p-value = 0.05



# Marginal Likelihoods & Evidence

Maximized Likelihood



$$\mathcal{BF}_{10} = 1.73$$

# Candidate's Formula (Besag 1989)

Alternative expression for Bayes Factor

$$\frac{p(y^{(n)} \mid \mathcal{H}_1)}{p(y^{(n)} \mid \mathcal{H}_0)} = \frac{\pi_\theta(0 \mid \mathcal{H}_1)}{\pi_\theta(0 \mid y^{(n)}, \mathcal{H}_1)}$$



# Candidate's Formula (Besag 1989)

Alternative expression for Bayes Factor

$$\frac{p(y^{(n)} \mid \mathcal{H}_1)}{p(y^{(n)} \mid \mathcal{H}_0)} = \frac{\pi_\theta(0 \mid \mathcal{H}_1)}{\pi_\theta(0 \mid y^{(n)}, \mathcal{H}_1)}$$

- ratio of the prior to posterior densities for  $\theta$  evaluated at zero



# Candidate's Formula (Besag 1989)

Alternative expression for Bayes Factor

$$\frac{p(y^{(n)} \mid \mathcal{H}_1)}{p(y^{(n)} \mid \mathcal{H}_0)} = \frac{\pi_\theta(0 \mid \mathcal{H}_1)}{\pi_\theta(0 \mid y^{(n)}, \mathcal{H}_1)}$$

- ratio of the prior to posterior densities for  $\theta$  evaluated at zero
- Savage-Dickey Ratio



# Prior

Plots were based on a  $\theta \mid \mathcal{H}_1 \sim N(0, 1)$



# Prior

Plots were based on a  $\theta | \mathcal{H}_1 \sim N(0, 1)$

- centered at value for  $\theta$  under  $\mathcal{H}_0$  (goes back to Jeffreys)



# Prior

Plots were based on a  $\theta | \mathcal{H}_1 \sim N(0, 1)$

- centered at value for  $\theta$  under  $\mathcal{H}_0$  (goes back to Jeffreys)
- "unit information prior" equivalent to a prior sample size is 1



# Prior

Plots were based on a  $\theta \mid \mathcal{H}_1 \sim N(0, 1)$

- centered at value for  $\theta$  under  $\mathcal{H}_0$  (goes back to Jeffreys)
  - "unit information prior" equivalent to a prior sample size is 1
  - What happens if  $n \rightarrow \infty$ ?




# Prior

Plots were based on a  $\theta | \mathcal{H}_1 \sim N(0, 1)$

- centered at value for  $\theta$  under  $\mathcal{H}_0$  (goes back to Jeffreys)
- "unit information prior" equivalent to a prior sample size is 1
- What happens if  $n \rightarrow \infty$ ?
- What happens of  $\tau_0 \rightarrow 0$  ?




# Precision




- $\tau_0 = 1/1000$

# Precision



- $\tau_0 = 1/1000$
- Posterior Probability of  $\mathcal{H}_0 = 0.9361$


# Precision



- $\tau_0 = 1/1000$
- Posterior Probability of  $\mathcal{H}_0 = 0.9361$
- As  $\tau_0 \rightarrow 0$  the posterior probability of  $\mathcal{H}_1$  goes to 0!



# Precision



- $\tau_0 = 1/1000$
- Posterior Probability of  $\mathcal{H}_0 = 0.9361$
- As  $\tau_0 \rightarrow 0$  the posterior probability of  $\mathcal{H}_1$  goes to 0!

**Bartlett's Paradox** - the paradox is that a seemingly non-informative prior for  $\theta$  is very informative about  $\mathcal{H}$ !

